Author Query Form

Journal code: EPJQT

Transmittal No.: EPJQT147

Typesetter's queries and/or remarks

Proof page/line	Details required	Author's response
	Author names are tagged as Given name(s) and Family name. Please confirm if these parts have been identified correctly and are presented in the right order. Note that response to this query is mandatory. Please note that the numbers of addresses were changed (31 addresses were duplicated in the file, 39 was missing).	
47/2178, 2196, 2200, 2202	City names are required in the addresses (70, 117, 128, 130). Please provide them.	
52/2443	Please check url address, which is not found.	

Many thanks for your assistance

Metadata of the article that will be visualized in Online First

Journal Name	EPJ Quantum Technology	
Article Title	Cold atoms in space: community workshop summary and proposed road-map	
Copyright holder	The Author(s) This will be the copyright line in the final PDF.	
Author	Family name	Alonso
	Particle	
	Given Name	Iván
	Suffix	
	Division	Higher Polytechnic School
	Organization	University of the Balearic Islands
	Address	Valldemossa Road, Palma de Mallorca, 07122, Spain
	E-mail	
Author	Family name	Alpigiani
	Particle	
	Given Name	Cristiano
	Suffix	
	Division	
	Organization	University of Washington
	Address	Physics-Astronomy Building, 15th Ave NE, Seattle, WA, 98195-1560, USA
	E-mail	
Author	Family name	Altschul
	Particle	
	Given Name	Brett
	Suffix	
	Division	
	Organization	University of South Carolina
	Address	Main Street, Columbia, SC, 29208, USA
	E-mail	
Author	Family name	Araújo
	Particle	
	Given Name	Henrique
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Arduini
	Particle	
	Given Name	Gianluigi
	Suffix	
	Division	
	Organization	CERN
	Address	CH-1211, Geneva, 23, Switzerland
	E-mail	
Author	Family name	Arlt
	Particle	
	Given Name	Jan
	Suffix	

	Division	
	Organization	Aarhus University
	Address	Ny Munkegade, 8000, Aarhus C, Denmark
	E-mail	
Author	Family name	Badurina
	Particle	
	Given Name	Leonardo
	Suffix	
	Division	
	Organization	King′s College London
	Address	Strand, London, WC2R 2LS, UK
	E-mail	
Author	Family name	Balaž
	Particle	
	Given Name	Antun
	Suffix	
	Division	
	Organization	University of Belgrade
	Address	Pregrevica 118, 11080, Belgrade, Serbia
	E-mail	
Author	Family name	Bandarupally
	Particle	
	Given Name	Satvika
	Suffix	
	Division	Dipartimento di Fisica e Astronomia
	Organization	Università di Firenze
	Address	Via Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	European Laboratory for Non-Linear Spectroscopy (LENS)
	Organization	Università degli Studi di Firenze
	Address	Via Giovanni Sansone, 1, 50019 Sesto Fiorentino, Firenze, Italy
	E-mail	
Author	Family name	Barish
	Particle	
	Given Name	Barry
	Given Name	C.
	Suffix	
	Division	LIGO
	Organization	California Institute of Technology
	Address	1201 E California, Pasadena, CA, 91125, USA
	E-mail	
Author	Family name	Barone
	Particle	
	Given Name	Michele
	Suffix	
	Division	Institute of Nuclear and Particle Physics
	Organization	NCSR Demokritos
	Address	Agia Paraskevi, 15310, Greece
	E-mail	
Author	Family name	Barsanti
	Particle	

	Given Name	Michele
	Suffix	
	Division	
	Organization	University of Pisa
	Address	Largo Lazzarino, Pisa, 56122, Italy
	E-mail	
Author	Family name	Bass
	Particle	
	Given Name	Steven
	Suffix	
	Division	
	Organization	Kitzbühel Centre for Physics
	Address	Kitzbühel, Austria
	E-mail	
Author	Family name	Bassi
	Particle	
	Given Name	Angelo
	Suffix	
	Division	
	Organization	University of Trieste
	Address	Strada Costiera 11, 34151, Trieste, Italy
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Trieste Section, Via Valerio 2, 34127, Trieste, Italy
	E-mail	
Author	Family name	Battelier
	Particle	
	Given Name	Baptiste
	Suffix	
	Division	LP2N, Laboratoire Photonique, Numérique et Nanosciences
	Organization	Université Bordeaux-IOGS-CNRS: UMR 5298
	Address	F-33400, Talence, France
	E-mail	
Author	Family name	Baynham
	Particle	
	Given Name	Charles
	Given Name	F.
	Given Name	A.
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Beaufils
	Particle	
	Given Name	Quentin
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE
	Address	61 avenue de l′Observatoire, 75014, Paris, France

	E-mail	
Author	Family name	Belić
	Particle	
	Given Name	Aleksandar
	Suffix	
	Division	
	Organization	University of Belgrade
	Address	Pregrevica 118, 11080, Belgrade, Serbia
	E-mail	
Author	Family name	Bergé
	Particle	
	Given Name	Joel
	Suffix	
	Division	DPHY, ONERA
	Organization	Université Paris-Saclay
	Address	Chemin de la Hunière—BP80100, F-91123, Palaiseau, France
	E-mail	
Author	Family name	Bernabeu
	Particle	
	Given Name	Jose
	Suffix	
	Division	Department of Theoretical Physics
	Organization	University of Valencia
	Address	Joint Centre Univ. Valencia-CSIC, 46100, Burjassot-Valencia, Spain
	Division	IFIC
	Organization	University of Valencia
	Address	Joint Centre Univ. Valencia-CSIC, 46100, Burjassot-Valencia, Spain
	E-mail	
Author	Family name	Bertoldi
	Particle	
	Given Name	Andrea
	Suffix	
	Division	LP2N, Laboratoire Photonique, Numérique et Nanosciences
	Organization	Université Bordeaux-IOGS-CNRS: UMR 5298
	Address	F-33400, Talence, France
	E-mail	
Author	Family name	Bingham
	Particle	
	Given Name	Robert
	Suffix	
	Division	
	Organization	Rutherford Appleton Laboratory
	Address	Harwell Campus, Didcot, OX11 0QX, UK
	Division	
	Organization	University of Strathclyde
	Address	Glasgow, G4 0NG, UK
	E-mail	
Author	Family name	Bize
	Particle	

	Given Name	Sébastien
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE
	Address	61 avenue de l′Observatoire, 75014, Paris, France
	E-mail	
Author	Family name	Blas
	Particle	
	Given Name	Diego
	Suffix	
	Division	Grup de Física Teòrica, Departament de Física
	Organization	Universitat Autònoma de Barcelona
	Address	08193, Bellaterra, Spain
	Division	Institut de Fisica d′Altes Energies
	Organization	The Barcelona Institute of Science and Technology
	Address	Campus UAB, 08193, Bellaterra (Barcelona), Spain
	E-mail	
Author	Family name	Bongs
	Particle	
	Given Name	Kai
	Suffix	
	Division	
	Organization	University of Birmingham
	Address	Birmingham, B15 2TT, UK
	E-mail	
Author	Family name	Bouyer
	Particle	
	Given Name	Philippe
	Suffix	
	Division	LP2N, Laboratoire Photonique, Numérique et Nanosciences
	Organization	Université Bordeaux-IOGS-CNRS: UMR 5298
	Address	F-33400, Talence, France
	E-mail	
Author	Family name	Braitenberg
	Particle	
	Given Name	Carla
	Suffix	
	Division	
	Organization	University of Trieste
	Address	Strada Costiera 11, 34151, Trieste, Italy
	E-mail	
Author	Family name	Brand
	Particle	
	Given Name	Christian
	Suffix	
	Division	German Aerospace Center (DRL)
	Organization	Institute of Quantum Technologies
	Address	Wilhelm-Runge-Straße 10, 89081, Ulm, Germany
	E-mail	
Author	Family name	Braxmaier

	Particle	
	Given Name	Claus
	Suffix	
	Division	Institute of Microelectronics
	Organization	Ulm University
	Address	Albert-Einstein-Allee 43, 89081, Ulm, Germany
	Division	German Aerospace Center (DRL)
	Organization	Institute of Quantum Technologies
	Address	Wilhelm-Runge-Straße 10, 89081, Ulm, Germany
	E-mail	
Author	Family name	Bresson
	Particle	
	Given Name	Alexandre
	Suffix	
	Division	DPHY, ONERA
	Organization	Université Paris-Saclay
	Address	Chemin de la Hunière—BP80100, F-91123, Palaiseau, France
	E-mail	
Corresponding Author	Family name	Buchmueller
	Particle	
	Given Name	Oliver
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	Division	Department of Physics, Clarendon Laboratory
	Organization	University of Oxford
	Address	South Parks Road, Oxford, OX1 3PU, UK
	E-mail	Oliver.Buchmueller@cern.ch
Author	Family name	Budker
	Particle	
	Given Name	Dmitry
	Suffix	
	Division	Helmholtz-Institut Mainz
	Organization	Johannes Gutenberg-Universität Mainz
	Address	GSI Helmholtzzentrum für Schwerionenforschung, 5128, Mainz, Germany
	Division	Department of Physics
	Organization	University of California
	Address	Berkeley, CA, 94720, USA
	E-mail	
Author	Family name	Bugalho
	Particle	
	Given Name	Luís
	Suffix	
	Division	Instituto de Telecomunicações
	Organization	Instituto Superior Técnico
	Address	Av. Rovisco Pais, Torre Norte, Lisboa, 1049-001, Portugal
	E-mail	
Author	Family name	Burdin
	Particle	

	Given Name	Sergey
	Suffix	
	Division	
	Organization	University of Liverpool
	Address	Liverpool, L69 7ZE, UK
	E-mail	
Author	Family name	Cacciapuoti
	Particle	
	Given Name	Luigi
	Suffix	
	Division	
	Organization	European Space Agency
	Address	Keplerlaan 1, PO Box 299, NL-2200, AG Noordwijk, The Netherlands
	E-mail	
Author	Family name	Callegari
	Particle	
	Given Name	Simone
	Suffix	
	Division	
	Organization	Physikalisch-Technische Bundesanstalt
	Address	Bundesallee 100, Braunschweig, 38116, Germany
	E-mail	
Author	Family name	Calmet
	Particle	
	Given Name	Xavier
	Suffix	
	Division	
	Organization	University of Sussex
	Address	Brighton, BN1 9QH, UK
	E-mail	
Author	Family name	Calonico
	Particle	
	Given Name	Davide
	Suffix	
	Division	
	Organization	Istituto Nazionale di Ricerca Metrologica
	Address	Strada delle Cacce 91, Turin, 10035, Italy
	E-mail	
Author	Family name	Canuel
	Particle	
	Given Name	Benjamin
	Suffix	
	Division	LP2N, Laboratoire Photonique, Numérique et Nanosciences
	Organization	Université Bordeaux-IOGS-CNRS: UMR 5298
	Address	F-33400, Talence, France
	E-mail	
Author	Family name	Caramete
	Particle	
	Given Name	Laurentiu-Ioan
	Suffix	

	Division	
	Organization	Institute of Space Science
	Address	409 Atomistilor Street, Bucharest, Magurele, Ilfov, 077125, Romania
	E-mail	
Author	Family name	Carraz
	Particle	
	Given Name	Olivier
	Suffix	
	Division	
	Organization	European Space Agency
	Address	Keplerlaan 1, PO Box 299, NL-2200, AG Noordwijk, The Netherlands
	E-mail	
Author	Family name	Cassettari
	Particle	
	Given Name	Donatella
	Suffix	
	Division	
	Organization	University of St Andrews
	Address	North Haugh, St Andrews, KY16 9SS, UK
	E-mail	
Author	Family name	Chakraborty
	Particle	
	Given Name	Pratik
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Chattopadhyay
	Particle	
	Given Name	Swapan
	Suffix	
	Division	
	Organization	Fermilab
	Address	Batavia, Illinois, USA
	Division	
	Organization	Stanford University
	Address	Stanford, CA, 94305, USA
	Division	Department of Physics
	Organization	University of California
	Address	Berkeley, CA, 94720, USA
	E-mail	
Author	Family name	Chauhan
	Particle	
	Given Name	Upasna
	Suffix	
	Division	
	Organization	Indian Institute of Technology
	Address	New Delhi, India

	E-mail	
Author	Family name	Chen
	Particle	
	Given Name	Xuzong
	Suffix	
	Division	
	Organization	Peking University
	Address	Beijing, 100871, China
	E-mail	
Author	Family name	Chen
	Particle	
	Given Name	Yu-Ao
	Suffix	
	Division	Hefei National Laboratory for Physical Sciences at the Microscale and Department of Modern Physics
	Organization	University of Science and Technology of China
	Address	Hefei, 230026, China
	Division	Shanghai Branch CAS Center for Excellence in Quantum Information and Quantum Physics
	Organization	University of Science and Technology of China
	Address	Shanghai, 201315, China
	Division	
	Organization	Shanghai Research Center for Quantum Sciences
	Address	Shanghai, 201315, China
	E-mail	
Author	Family name	Chiofalo
	Particle	
	Given Name	Maria
	Given Name	Luisa
	Suffix	
	Division	
	Organization	University of Pisa
	Address	Largo Lazzarino, Pisa, 56122, Italy
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Sezione di Pisa, Largo Pontecorvo 3, I-56125, Pisa, Italy
	E-mail	
Author	Family name	Coleman
	Particle	
	Given Name	Jonathon
	Suffix	
	Division	
	Organization	University of Liverpool
	Address	Liverpool, L69 7ZE, UK
	E-mail	
Author	Family name	Corgier
	Particle	
	Given Name	Robin
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE

	Address	61 avenue de l′Observatoire, 75014, Paris, France
	E-mail	
Author	Family name	Cotter
	Particle	
	Given Name	J.
	Given Name	P.
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Michael Cruise
	Particle	
	Given Name	A.
	Suffix	
	Division	
	Organization	University of Birmingham
	Address	Birmingham, B15 2TT, UK
	E-mail	
Author	Family name	Cui
	Particle	
	Given Name	Yanou
	Suffix	
	Division	Department of Physics and Astronomy
	Organization	University of California
	Address	Riverside, CA, 92521, USA
	E-mail	
Author	Family name	Davies
	Particle	
	Given Name	Gavin
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	De Roeck
	Particle	
	Given Name	Albert
	Suffix	
	Division	
	Organization	Antwerp University
	Address	B 2610, Wilrijk, Belgium
	Division	
	Organization	CERN
	Address	CH-1211, Geneva, 23, Switzerland
	E-mail	
Author	Family name	Demarteau
	Particle	
	Given Name	Marcel
	Suffix	

	Division	
	Organization	Oak Ridge National Laboratory
	Address	1 Bethel Valley Road, Oak Ridge, TN, 37830, USA
	E-mail	
Author	Family name	Derevianko
	Particle	
	Given Name	Andrei
	Suffix	
	Division	
	Organization	University of Nevada
	Address	Reno, Nevada, 89557, USA
	E-mail	
Author	Family name	Di Clemente
	Particle	
	Given Name	Marco
	Suffix	
	Division	
	Organization	Italian Space Agency
	Address	Via del Politecnico, Rome, 00133, Italy
	E-mail	
Author	Family name	Djordjevic
	Particle	
	Given Name	Goran
	Given Name	S.
	Suffix	
	Division	
	Organization	University of Nis
	Address	Visegradska 33, Nis, Serbia
	E-mail	
Author	Family name	Donadi
	Particle	
	Given Name	Sandro
	Suffix	
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Trieste Section, Via Valerio 2, 34127, Trieste, Italy
	E-mail	
Author	Family name	Doré
	Particle	
	Given Name	Olivier
	Suffix	
	Division	Jet Propulsion Laboratory
	Organization	California Institute of Technology
	Address	Pasadena, CA, 91109, USA
	E-mail	
Author	Family name	Dornan
	Particle	
	Given Name	Peter
	Suffix	
	Division	

	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Doser
	Particle	
	Given Name	Michael
	Suffix	
	Division	
	Organization	CERN
	Address	CH-1211, Geneva, 23, Switzerland
	E-mail	
Author	Family name	Drougakis
	Particle	
	Given Name	Giannis
	Suffix	
	Division	
	Organization	Foundation for Research and Technology-Hellas
	Address	100 N. Plastira Street, Vassilika Vouton, GR 70013, Heraklion, Crete, Greece
	E-mail	
Author	Family name	Dunningham
	Particle	
	Given Name	Jacob
	Suffix	
	Division	
	Organization	University of Sussex
	Address	Brighton, BN1 9QH, UK
	E-mail	
Author	Family name	Easo
	Particle	
	Given Name	Sajan
	Suffix	
	Division	
	Organization	Rutherford Appleton Laboratory
	Address	Harwell Campus, Didcot, OX11 0QX, UK
	E-mail	
Author	Family name	Eby
	Particle	
	Given Name	Joshua
	Suffix	
	Division	The University of Tokyo Institutes for Advanced Study
	Organization	The University of Tokyo
	Address	Kashiwa, Chiba, 277-8583, Japan
	E-mail	
Author	Family name	Elertas
	Particle	
	Given Name	Gedminas
	Suffix	
	Division	
	Organization	University of Liverpool
	Address	Liverpool, L69 7ZE, UK

	E-mail	
Corresponding Author	Family name	Ellis
	Particle	
	Given Name	John
	Suffix	
	Division	
	Organization	King′s College London
	Address	Strand, London, WC2R 2LS, UK
	Division	
	Organization	CERN
	Address	CH-1211, Geneva, 23, Switzerland
	E-mail	John.Ellis@cern.ch
Author	Family name	Evans
	Particle	
	Given Name	David
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Examilioti
	Particle	
	Given Name	Pandora
	Suffix	
	Division	
	Organization	Foundation for Research and Technology-Hellas
	Address	100 N. Plastira Street, Vassilika Vouton, GR 70013, Heraklion, Crete, Greece
	E-mail	
Author	Family name	Fadeev
	Particle	
	Given Name	Pavel
	Suffix	
	Division	Helmholtz-Institut Mainz
	Organization	Johannes Gutenberg-Universität Mainz
	Address	GSI Helmholtzzentrum für Schwerionenforschung, 5128, Mainz, Germany
	E-mail	
Author	Family name	Fanì
	Particle	
	Given Name	Mattia
	Suffix	
	Division	
	Organization	Los Alamos National Laboratory
	Address	Los Alamos, New Mexico, USA
	E-mail	
Author	Family name	Fassi
	Particle	
	Given Name	Farida
	Suffix	
	Division	Faculty of Science

	Organization	Mohammed V University in Rabat
	Address	4 Avenue Ibn Battouta, B.P. 1014 RP, Rabat, Morocco
	E-mail	
Author	Family name	Fattori
	Particle	
	Given Name	Marco
	Suffix	
	Division	Dipartimento di Fisica e Astronomia
	Organization	Università di Firenze
	Address	Via Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy
	E-mail	
Author	Family name	Fedderke
	Particle	
	Given Name	Michael
	Given Name	A.
	Suffix	
	Division	
	Organization	The Johns Hopkins University
	Address	3400 N Charles St, Baltimore, MD, 21218, USA
	E-mail	
Author	Family name	Felea
	Particle	
	Given Name	Daniel
	Suffix	
	Division	
	Organization	Institute of Space Science
	Address	409 Atomistilor Street, Bucharest, Magurele, Ilfov, 077125, Romania
	E-mail	
Author	Family name	Feng
	Particle	
	Given Name	Chen-Hao
	Suffix	
	Division	LP2N, Laboratoire Photonique, Numérique et Nanosciences
	Organization	Université Bordeaux-IOGS-CNRS: UMR 5298
	Address	F-33400, Talence, France
	E-mail	
Author	Family name	Ferreras
	Particle	
	Given Name	Jorge
	Suffix	
	Division	
	Organization	Rutherford Appleton Laboratory
	Address	Harwell Campus, Didcot, OX11 0QX, UK
	E-mail	
Author	Family name	Flack
	Particle	
	Given Name	Robert
	Suffix	
	Division	
	Organization	University College London

	Address	Gower Street, London, WC1E 6BT, UK
	E-mail	
Author	Family name	Flambaum
	Particle	
	Given Name	Victor
	Given Name	V.
	Suffix	
	Division	School of Physics
	Organization	University of New South Wales
	Address	Sydney, 2052, Australia
	E-mail	
Author	Family name	Forsberg
	Particle	
	Given Name	René
	Suffix	
	Division	
	Organization	Technical University of Denmark
	Address	Kongens Lyngby, 2800, Denmark
	E-mail	
Author	Family name	Fromhold
	Particle	
	Given Name	Mark
	Suffix	
	Division	
	Organization	University of Nottingham
	Address	Nottingham, NG7 2RD, UK
	E-mail	
Author	Family name	Gaaloul
	Particle	
	Given Name	Naceur
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Garraway
	Particle	
	Given Name	Barry
	Given Name	M.
	Suffix	
	Division	
	Organization	University of Sussex
	Address	Brighton, BN1 9QH, UK
	E-mail	
Author	Family name	Georgousi
	Particle	
	Given Name	Maria
	Suffix	
	Division	
	Organization	Foundation for Research and Technology-Hellas

	Address	100 N. Plastira Street, Vassilika Vouton, GR 70013, Heraklion, Crete, Greece
	E-mail	
Author	Family name	Geraci
	Particle	
	Given Name	Andrew
	Suffix	
	Division	
	Organization	Northwestern University
	Address	Evanston, USA
	E-mail	
Author	Family name	Gibble
	Particle	
	Given Name	Kurt
	Suffix	
	Division	
	Organization	The Pennsylvania State University
	Address	University Park, PA, 16802, USA
	E-mail	
Author	Family name	Gibson
	Particle	
	Given Name	Valerie
	Suffix	
	Division	
	Organization	University of Cambridge
	Address	J.J. Thomson Avenue, Cambridge, CB3 0HE, UK
	E-mail	
Author	Family name	Gill
	Particle	
	Given Name	Patrick
	Suffix	
	Division	
	Organization	National Physical Laboratory
	Address	Hampton Road, Teddington, TW11 0LW, UK
	E-mail	
Author	Family name	Giudice
	Particle	
	Given Name	Gian
	Given Name	F.
	Suffix	
	Division	
	Organization	CERN
	Address	CH-1211, Geneva, 23, Switzerland
	E-mail	
Author	Family name	Goldwin
	Particle	
	Given Name	Jon
	Suffix	
	Division	
	Organization	University of Birmingham
	Address	Birmingham, B15 2TT, UK

	E-mail	
Author	Family name	Gould
	Particle	
	Given Name	Oliver
	Suffix	
	Division	
	Organization	University of Nottingham
	Address	Nottingham, NG7 2RD, UK
	E-mail	
Author	Family name	Grachov
	Particle	
	Given Name	Oleg
	Suffix	
	Division	Department of Physics and Astronomy
	Organization	Wayne State University
	Address	Detroit, MI, 48202, USA
	E-mail	
Author	Family name	Graham
	Particle	
	Given Name	Peter
	Given Name	W.
	Suffix	
	Division	
	Organization	Stanford University
	Address	Stanford, CA, 94305, USA
	E-mail	
Author	Family name	Grasso
	Particle	
	Given Name	Dario
	Suffix	
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Sezione di Pisa, Largo Pontecorvo 3, I-56125, Pisa, Italy
	E-mail	
Author	Family name	Griffin
	Particle	
	Given Name	Paul
	Given Name	F.
	Suffix	
	Division	
	Organization	University of Strathclyde
	Address	Glasgow, G4 0NG, UK
	E-mail	
Author	Family name	Guerlin
	Particle	
	Given Name	Christine
	Suffix	
	Division	Laboratoire Kastler Brossel
	Organization	Sorbonne Université, CNRS, ENS-Université PSL, Collège de France
	Address	???, France

	E-mail	
Author	Family name	Gündoğan
	Particle	
	Given Name	Mustafa
	Suffix	
	Division	
	Organization	Humboldt-Universität zu Berlin
	Address	Newtonstraße 15, Berlin, 12489, Germany
	E-mail	
Author	Family name	Gupta
	Particle	
	Given Name	Ratnesh
	Given Name	K.
	Suffix	
	Division	
	Organization	Okinawa Institute of Science and Technology
	Address	Okinawa, Japan
	E-mail	
Author	Family name	Haehnelt
	Particle	
	Given Name	Martin
	Suffix	
	Division	
	Organization	University of Cambridge
	Address	J.J. Thomson Avenue, Cambridge, CB3 0HE, UK
	E-mail	
Author	Family name	Hanımeli
	Particle	
	Given Name	Ekim
	Given Name	T.
	Suffix	
	Division	
	Organization	Universität Bremen
	Address	Am Fallturm 2, 28359, Bremen, Germany
	E-mail	
Author	Family name	Hawkins
	Particle	
	Given Name	Leonie
	Suffix	
	Division	
	Organization	University of Liverpool
	Address	Liverpool, L69 7ZE, UK
	E-mail	
Author	Family name	Hees
	Particle	
	Given Name	Aurélien
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE
	Address	61 avenue de l ′ Observatoire, 75014, Paris, France

	E-mail	
Author	Family name	Henderson
	Particle	
	Given Name	Victoria
	Given Name	A.
	Suffix	
	Division	
	Organization	Humboldt-Universität zu Berlin
	Address	Newtonstraße 15, Berlin, 12489, Germany
	E-mail	
Author	Family name	Herr
	Particle	
	Given Name	Waldemar
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Herrmann
	Particle	
	Given Name	Sven
	Suffix	
	Division	
	Organization	Universität Bremen
	Address	Am Fallturm 2, 28359, Bremen, Germany
	E-mail	
Author	Family name	Hird
	Particle	
	Given Name	Thomas
	Suffix	
	Division	Department of Physics, Clarendon Laboratory
	Organization	University of Oxford
	Address	South Parks Road, Oxford, OX1 3PU, UK
	E-mail	
Author	Family name	Hobson
	Particle	
	Given Name	Richard
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Hock
	Particle	
	Given Name	Vincent
	Suffix	
	Division	
	Organization	Universität Bremen
	Address	Am Fallturm 2, 28359, Bremen, Germany
	E-mail	

Author	Family name	Hogan
	Particle	
	Given Name	Jason
	Given Name	M.
	Suffix	
	Division	
	Organization	Stanford University
	Address	Stanford, CA, 94305, USA
	E-mail	
Author	Family name	Holst
	Particle	
	Given Name	Bodil
	Suffix	
	Division	
	Organization	University of Bergen
	Address	Allégaten 55, 5007, Bergen, Norway
	E-mail	
Author	Family name	Holynski
	Particle	
	Given Name	Michael
	Suffix	
	Division	
	Organization	University of Birmingham
	Address	Birmingham, B15 2TT, UK
	E-mail	
Author	Family name	Israelsson
	Particle	
	Given Name	Ulf
	Suffix	
	Division	Jet Propulsion Laboratory
	Organization	California Institute of Technology
	Address	Pasadena, CA, 91109, USA
	E-mail	
Author	Family name	Jeglič
	Particle	
	Given Name	Peter
	Suffix	
	Division	
	Organization	Jožef Stefan Institute
	Address	Jamova 39, 1000, Ljubljana, Slovenia
	E-mail	
Author	Family name	Jetzer
	Particle	
	Given Name	Philippe
	Suffix	
	Division	
	Organization	University of Zurich
	Address	Winterthurerstrasse 190, 8057, Zurich, Switzerland
	E-mail	
Author	Family name	Juzeliūnas

	Particle	
	Given Name	Gediminas
	Suffix	
	Division	
	Organization	Vilnius University
	Address	Saulėtekio 3, Vilnius, LT-10257, Lithuania
	E-mail	
Author	Family name	Kaltenbaek
	Particle	
	Given Name	Rainer
	Suffix	
	Division	
	Organization	University of Ljubljana
	Address	Jadranska ulica 19, 1000, Ljubljana, Slovenia
	E-mail	
Author	Family name	Kamenik
	Particle	
	Given Name	Jernej
	Given Name	F.
	Suffix	
	Division	
	Organization	University of Ljubljana
	Address	Jadranska ulica 19, 1000, Ljubljana, Slovenia
	E-mail	
Author	Family name	Kehagias
	Particle	
	Given Name	Alex
	Suffix	
	Division	Physics Division
	Organization	National Technical University of Athens
	Address	Zografou, Athens, 15780, Greece
	E-mail	
Author	Family name	Kirova
	Particle	
	Given Name	Teodora
	Suffix	
	Division	
	Organization	University of Latvia
	Address	Riga, LV-1004, Latvia
	E-mail	
Author	Family name	Kiss-Toth
	Particle	
	Given Name	Marton
	Suffix	
	Division	
	Organization	Teledyne e2v
	Address	106 Waterhouse Lane, Chelmsford, CM1 2QU, UK
	E-mail	
Author	Family name	Koke
	Particle	

	Given Name	Sebastian
	Suffix	
	Division	
	Organization	Physikalisch-Technische Bundesanstalt
	Address	Bundesallee 100, Braunschweig, 38116, Germany
	E-mail	
Author	Family name	Kolkowitz
	Particle	
	Given Name	Shimon
	Suffix	
	Division	
	Organization	University of Wisconsin
	Address	Madison, WI, 53706, USA
	E-mail	
Author	Family name	Kornakov
	Particle	
	Given Name	Georgy
	Suffix	
	Division	Faculty of Physics
	Organization	Warsaw University of Technology
	Address	ul. Koszykowa 75, 00-662, Warsaw, Poland
	E-mail	
Author	Family name	Kovachy
	Particle	
	Given Name	Tim
	Suffix	
	Division	
	Organization	Northwestern University
	Address	Evanston, USA
	E-mail	
Author	Family name	Krutzik
	Particle	
	Given Name	Markus
	Suffix	
	Division	
	Organization	Humboldt-Universität zu Berlin
	Address	Newtonstraße 15, Berlin, 12489, Germany
	E-mail	
Author	Family name	Kumar
	Particle	
	Given Name	Mukesh
	Suffix	
	Division	
	Organization	University of the Witwatersrand
	Address	Johannesburg, Wits, 2050, South Africa
	E-mail	
Author	Family name	Kumar
	Particle	
	Given Name	Pradeep
	Suffix	

	Division	
	Organization	Indian Institute of Science Education and Research
	Address	Bhopal, 462066, India
	E-mail	
Author	Family name	Lämmerzahl
	Particle	
	Given Name	Claus
	Suffix	
	Division	
	Organization	Universität Bremen
	Address	Am Fallturm 2, 28359, Bremen, Germany
	E-mail	
Author	Family name	Landsberg
	Particle	
	Given Name	Greg
	Suffix	
	Division	Department of Physics
	Organization	Brown University
	Address	Providence, RI, 02912, USA
	E-mail	
Author	Family name	Le Poncin-Lafitte
	Particle	
	Given Name	Christophe
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE
	Address	61 avenue de l′Observatoire, 75014, Paris, France
	E-mail	
Author	Family name	Leibrandt
	Particle	
	Given Name	David
	Given Name	R.
	Suffix	
	Division	
	Organization	National Institute of Standards and Technology
	Address	325 Broadway Street, Boulder, Colorado, 80305, USA
	E-mail	
Author	Family name	Lévèque
	Particle	
	Given Name	Thomas
	Suffix	
	Division	
	Organization	Centre National d′Etudes Spatiales
	Address	18 Avenue Edouard Belin, 31400, Toulouse, France
	E-mail	
Author	Family name	Lewicki
	Particle	
	Given Name	Marek
	Suffix	
	Division	

	Organization	University of Warsaw
	Address	ul. Pasteura 5, 02-093, Warsaw, Poland
	E-mail	
Author	Family name	Li
	Particle	
	Given Name	Rui
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Lipniacka
	Particle	
	Given Name	Anna
	Suffix	
	Division	
	Organization	University of Bergen
	Address	Allégaten 55, 5007, Bergen, Norway
	E-mail	
Author	Family name	Lisdat
	Particle	
	Given Name	Christian
	Suffix	
	Division	
	Organization	Physikalisch-Technische Bundesanstalt
	Address	Bundesallee 100, Braunschweig, 38116, Germany
	E-mail	
Author	Family name	Liu
	Particle	
	Given Name	Mia
	Suffix	
	Division	
	Organization	Purdue University
	Address	West Lafayette, IN, 47907, USA
	E-mail	
Author	Family name	Lopez-Gonzalez
	Particle	
	Given Name	J.
	Given Name	L.
	Suffix	
	Division	
	Organization	Autonomous University of Aguascalientes
	Address	Av. Universidad 940, Ciudad Universitaria, Aguascalientes, C. P. 20131, Mexico
	E-mail	
Author	Family name	Loriani
	Particle	
	Given Name	Sina
	Suffix	
	Division	
	Organization	Potsdam Institute for Climate Impact Research

	Address	Member of the Leibniz Association, 14473, Potsdam, Germany
	E-mail	
Author	Family name	Louko
	Particle	
	Given Name	Jorma
	Suffix	
	Division	
	Organization	University of Nottingham
	Address	Nottingham, NG7 2RD, UK
	E-mail	
Author	Family name	Luciano
	Particle	
	Given Name	Giuseppe
	Given Name	Gaetano
	Suffix	
	Division	
	Organization	Università degli Studi di Salerno
	Address	Via Giovanni Paolo II 132, I-84084, Fisciano (SA), Italy
	E-mail	
Author	Family name	Lundblad
	Particle	
	Given Name	Nathan
	Suffix	
	Division	
	Organization	Bates College
	Address	Lewiston, ME, 04240, USA
	E-mail	
Author	Family name	Maddox
	Particle	
	Given Name	Steve
	Suffix	
	Division	
	Organization	Teledyne e2v
	Address	106 Waterhouse Lane, Chelmsford, CM1 2QU, UK
	E-mail	
Author	Family name	Mahmoud
	Particle	
	Given Name	M.
	Given Name	A.
	Suffix	
	Division	
	Organization	Fayoum University
	Address	El-Fayoum, Egypt
	E-mail	
Author	Family name	Maleknejad
	Particle	
	Given Name	Azadeh
	Suffix	
	Division	
	Organization	CERN

	Address	CH-1211, Geneva, 23, Switzerland
	E-mail	
Author	Family name	March-Russell
	Particle	
	Given Name	John
	Suffix	
	Division	Department of Physics, Clarendon Laboratory
	Organization	University of Oxford
	Address	South Parks Road, Oxford, OX1 3PU, UK
	E-mail	
Author	Family name	Massonnet
	Particle	
	Given Name	Didier
	Suffix	
	Division	
	Organization	Centre National d ′ Etudes Spatiales
	Address	18 Avenue Edouard Belin, 31400, Toulouse, France
	E-mail	
Author	Family name	McCabe
	Particle	
	Given Name	Christopher
	Suffix	
	Division	
	Organization	King ′ s College London
	Address	Strand, London, WC2R 2LS, UK
	E-mail	
Author	Family name	Meister
	Particle	
	Given Name	Matthias
	Suffix	
	Division	German Aerospace Center (DRL)
	Organization	Institute of Quantum Technologies
	Address	Wilhelm-Runge-Straße 10, 89081, Ulm, Germany
	E-mail	
Author	Family name	Mežnaršič
	Particle	
	Given Name	Tadej
	Suffix	
	Division	
	Organization	Jožef Stefan Institute
	Address	Jamova 39, 1000, Ljubljana, Slovenia
	E-mail	
Author	Family name	Micalizio
	Particle	
	Given Name	Salvatore
	Suffix	
	Division	
	Organization	Istituto Nazionale di Ricerca Metrologica
	Address	Strada delle Cacce 91, Turin, 10035, Italy
	E-mail	

Author	Family name	Migliaccio
	Particle	
	Given Name	Federica
	Suffix	
	Division	
	Organization	Politecnico di Milano
	Address	Piazza Leonardo da Vinci 32, 20133, Milano, Italy
	E-mail	
Author	Family name	Millington
	Particle	
	Given Name	Peter
	Suffix	
	Division	
	Organization	University of Nottingham
	Address	Nottingham, NG7 2RD, UK
	E-mail	
Author	Family name	Milosevic
	Particle	
	Given Name	Milan
	Suffix	
	Division	
	Organization	University of Nis
	Address	Visegradska 33, Nis, Serbia
	E-mail	
Author	Family name	Mitchell
	Particle	
	Given Name	Jeremiah
	Suffix	
	Division	
	Organization	University of Cambridge
	Address	J.J. Thomson Avenue, Cambridge, CB3 0HE, UK
	E-mail	
Author	Family name	Morley
	Particle	
	Given Name	Gavin
	Given Name	W.
	Suffix	
	Division	
	Organization	University of Warwick
	Address	Coventry, CV4 7AL, UK
	E-mail	
Author	Family name	Müller
	Particle	
	Given Name	Jürgen
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Murphy

	Particle	
	Given Name	Eamonn
	Suffix	
	Division	
	Organization	European Space Agency
	Address	Keplerlaan 1, PO Box 299, NL-2200, AG Noordwijk, The Netherlands
	E-mail	
Author	Family name	Müstecaplıoğlu
	Particle	
	Given Name	Özgür
	Given Name	E.
	Suffix	
	Division	Department of Physics
	Organization	Koç University
	Address	İstanbul, Sarıyer, 34450, Turkey
	E-mail	
Author	Family name	O′Shea
	Particle	
	Given Name	Val
	Suffix	
	Division	
	Organization	University of Glasgow
	Address	Glasgow, G12 8QQ, UK
	E-mail	
Author	Family name	Oi
	Particle	
	Given Name	Daniel
	Given Name	K.
	Given Name	L.
	Suffix	
	Division	
	Organization	University of Strathclyde
	Address	Glasgow, G4 0NG, UK
	E-mail	
Author	Family name	Olson
	Particle	
	Given Name	Judith
	Suffix	
	Division	
	Organization	ColdQuanta
	Address	Boulder, Colorado, USA
	E-mail	
Author	Family name	Pal
	Particle	
	Given Name	Debapriya
	Suffix	
	Division	
	Organization	AMOLF
	Address	Amsterdam, The Netherlands
	E-mail	

Author	Family name	Papazoglou
	Particle	
	Given Name	Dimitris
	Given Name	G.
	Suffix	
	Division	
	Organization	University of Crete
	Address	P.O. Box 2208, 71003, Heraklion, Greece
	E-mail	
Author	Family name	Pasatembou
	Particle	
	Given Name	Elizabeth
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Paternostro
	Particle	
	Given Name	Mauro
	Suffix	
	Division	
	Organization	Queen′s University
	Address	Belfast, BT7 1NN, UK
	E-mail	
Author	Family name	Pawlowski
	Particle	
	Given Name	Krzysztof
	Suffix	
	Division	
	Organization	Center for Theoretical Physics PAS
	Address	Aleja Lotnikow 32/46, 02-668, Warsaw, Poland
	E-mail	
Author	Family name	Pelucchi
	Particle	
	Given Name	Emanuele
	Suffix	
	Division	Tyndall National Institute
	Organization	University College Cork
	Address	Cork, Ireland
	E-mail	
Author	Family name	Pereira dos Santos
	Particle	
	Given Name	Franck
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE
	Address	61 avenue de l′Observatoire, 75014, Paris, France
	E-mail	
Author	Family name	Peters

	Particle	
	Given Name	Achim
	Suffix	
	Division	
	Organization	Humboldt-Universität zu Berlin
	Address	Newtonstraße 15, Berlin, 12489, Germany
	E-mail	
Author	Family name	Pikovski
	Particle	
	Given Name	Igor
	Suffix	
	Division	Stevens Institute of Technology
	Organization	Castle Point on the Hudson
	Address	Hoboken, NJ, 07030, USA
	Division	
	Organization	Stockholm University
	Address	AlbaNova, SE-10691, Stockholm, Sweden
	E-mail	
Author	Family name	Pilaftsis
	Particle	
	Given Name	Apostolos
	Suffix	
	Division	Department of Physics and Astronomy
	Organization	University of Manchester
	Address	Manchester, M13 9PL, UK
	E-mail	
Author	Family name	Pinto
	Particle	
	Given Name	Alexandra
	Suffix	
	Division	Hoursec
	Organization	Energy Efficient AI On-Chip
	Address	Uetlibergstrasse 111B, Zurich, 8045, Switzerland
	E-mail	
Author	Family name	Prevedelli
	Particle	
	Given Name	Marco
	Suffix	
	Division	
	Organization	University of Bologna
	Address	Viale Berti-Pichat 6/2, Bologna, I-40126, Italy
	E-mail	
Author	Family name	Puthiya-Veettil
	Particle	
	Given Name	Vishnupriya
	Suffix	
	Division	
	Organization	Foundation for Research and Technology-Hellas
	Address	100 N. Plastira Street, Vassilika Vouton, GR 70013, Heraklion, Crete, Greece
	E-mail	

Author	Family name	Quenby
	Particle	
	Given Name	John
	Suffix	
	Division	
	Organization	Imperial College London
	Address	Prince Consort Road, London, SW7 2AZ, UK
	E-mail	
Author	Family name	Rafelski
	Particle	
	Given Name	Johann
	Suffix	
	Division	Department of Physics
	Organization	The University of Arizona
	Address	Tucson, AZ, 85721-0081, USA
	E-mail	
Author	Family name	Rasel
	Particle	
	Given Name	Ernst
	Given Name	M.
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Ravensbergen
	Particle	
	Given Name	Cornelis
	Suffix	
	Division	
	Organization	ColdQuanta
	Address	Boulder, Colorado, USA
	E-mail	
Author	Family name	Reguzzoni
	Particle	
	Given Name	Mirko
	Suffix	
	Division	
	Organization	Politecnico di Milano
	Address	Piazza Leonardo da Vinci 32, 20133, Milano, Italy
	Division	
	Organization	Antwerp University
	Address	B 2610, Wilrijk, Belgium
	E-mail	
Author	Family name	Richaud
	Particle	
	Given Name	Andrea
	Suffix	
	Division	
	Organization	Scuola Internazionale Superiore di Studi Avanzati

	Address	Via Bonomea 265, I-34136, Trieste, Italy
	E-mail	
Author	Family name	Riou
	Particle	
	Given Name	Isabelle
	Suffix	
	Division	
	Organization	Teledyne e2v
	Address	106 Waterhouse Lane, Chelmsford, CM1 2QU, UK
	E-mail	
Author	Family name	Rothacher
	Particle	
	Given Name	Markus
	Suffix	
	Division	
	Organization	ETH Zurich
	Address	Robert-Gnehm-Weg 15, CH-8093, Zurich, Switzerland
	E-mail	
Author	Family name	Roura
	Particle	
	Given Name	Albert
	Suffix	
	Division	German Aerospace Center (DRL)
	Organization	Institute of Quantum Technologies
	Address	Wilhelm-Runge-Straße 10, 89081, Ulm, Germany
	E-mail	
Author	Family name	Ruschhaupt
	Particle	
	Given Name	Andreas
	Suffix	
	Division	Tyndall National Institute
	Organization	University College Cork
	Address	Cork, Ireland
	E-mail	
Author	Family name	Sabulsky
	Particle	
	Given Name	Dylan
	Given Name	O.
	Suffix	
	Division	LP2N, Laboratoire Photonique, Numérique et Nanosciences
	Organization	Université Bordeaux-IOGS-CNRS: UMR 5298
	Address	F-33400, Talence, France
	E-mail	
Author	Family name	Safronova
	Particle	
	Given Name	Marianna
	Suffix	
	Division	
	Organization	University of Delaware
	Address	Newark, Delaware, 19716, USA

	E-mail	
Author	Family name	Saltas
	Particle	
	Given Name	Ippocratis
	Given Name	D.
	Suffix	
	Division	
	Organization	Institute of Physics of the Czech Academy of Sciences
	Address	Na Slovance 2, 182 21, Praha, 8, Czechia
	E-mail	
Author	Family name	Salvi
	Particle	
	Given Name	Leonardo
	Suffix	
	Division	Dipartimento di Fisica e Astronomia
	Organization	Università di Firenze
	Address	Via Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	European Laboratory for Non-Linear Spectroscopy (LENS)
	Organization	Università degli Studi di Firenze
	Address	Via Giovanni Sansone, 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Sezione di Firenze, via Sansone 1, Sesto Fiorentino, Firenze, Italy
	E-mail	
Author	Family name	Sameed
	Particle	
	Given Name	Muhammed
	Suffix	
	Division	Department of Physics and Astronomy
	Organization	University of Manchester
	Address	Manchester, M13 9PL, UK
	E-mail	
Author	Family name	Saurabh
	Particle	
	Given Name	Pandey
	Suffix	
	Division	The University of Tokyo Institutes for Advanced Study
	Organization	The University of Tokyo
	Address	Kashiwa, Chiba, 277-8583, Japan
	E-mail	
Author	Family name	Schäffer
	Particle	
	Given Name	Stefan
	Suffix	
	Division	Van der Waals-Zeeman Institute
	Organization	University of Amsterdam
	Address	Science Park 904, 1098XH, ???, The Netherlands
	E-mail	
Author	Family name	Schiller
	Particle	

	Given Name	Stephan
	Suffix	
	Division	
	Organization	Heinrich-Heine-Universität
	Address	Düsseldorf, Germany
	E-mail	
Author	Family name	Schilling
	Particle	
	Given Name	Manuel
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Schkolnik
	Particle	
	Given Name	Vladimir
	Suffix	
	Division	
	Organization	Humboldt-Universität zu Berlin
	Address	Newtonstraße 15, Berlin, 12489, Germany
	E-mail	
Author	Family name	Schlippert
	Particle	
	Given Name	Dennis
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Schmidt
	Particle	
	Given Name	Piet
	Given Name	O.
	Suffix	
	Division	
	Organization	Physikalisch-Technische Bundesanstalt
	Address	Bundesallee 100, Braunschweig, 38116, Germany
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Schnatz
	Particle	
	Given Name	Harald
	Suffix	
	Division	
	Organization	Physikalisch-Technische Bundesanstalt
	Address	Bundesallee 100, Braunschweig, 38116, Germany
	E-mail	

Author	Family name	Schneider
	Particle	
	Given Name	Jean
	Suffix	
	Division	Laboratoire Univers et Théorie
	Organization	Observatoire de Paris-PSL
	Address	92190, Meudon, France
	E-mail	
Author	Family name	Schneider
	Particle	
	Given Name	Ulrich
	Suffix	
	Division	
	Organization	University of Cambridge
	Address	J.J. Thomson Avenue, Cambridge, CB3 0HE, UK
	E-mail	
Author	Family name	Schreck
	Particle	
	Given Name	Florian
	Suffix	
	Division	Van der Waals-Zeeman Institute
	Organization	University of Amsterdam
	Address	Science Park 904, 1098XH, ???, The Netherlands
	E-mail	
Author	Family name	Schubert
	Particle	
	Given Name	Christian
	Suffix	
	Division	
	Organization	Leibniz Universität Hannover
	Address	Welfengarten 1, Hannover, 30167, Germany
	E-mail	
Author	Family name	Shayeghi
	Particle	
	Given Name	Armin
	Suffix	
	Division	
	Organization	University of Vienna
	Address	Boltzmanngasse 5, A-1090, Vienna, Austria
	E-mail	
Author	Family name	Sherrill
	Particle	
	Given Name	Nathaniel
	Suffix	
	Division	
	Organization	University of Sussex
	Address	Brighton, BN1 9QH, UK
	E-mail	
Author	Family name	Shipsey
	Particle	

	Given Name	Ian
	Suffix	
	Division	Department of Physics, Clarendon Laboratory
	Organization	University of Oxford
	Address	South Parks Road, Oxford, OX1 3PU, UK
	E-mail	
Author	Family name	Signorini
	Particle	
	Given Name	Carla
	Suffix	
	Division	
	Organization	University of Pisa
	Address	Largo Lazzarino, Pisa, 56122, Italy
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Sezione di Pisa, Largo Pontecorvo 3, I-56125, Pisa, Italy
	E-mail	
Author	Family name	Singh
	Particle	
	Given Name	Rajeev
	Suffix	
	Division	
	Organization	Polish Academy of Sciences
	Address	PL 31-342, Krakow, Poland
	E-mail	
Author	Family name	Singh
	Particle	
	Given Name	Yeshpal
	Suffix	
	Division	
	Organization	University of Birmingham
	Address	Birmingham, B15 2TT, UK
	E-mail	
Author	Family name	Skordis
	Particle	
	Given Name	Constantinos
	Suffix	
	Division	CEICO−FZU
	Organization	Institute of Physics of the Czech Academy of Sciences
	Address	Na Slovance 1999/2, 182 21, Prague, Czechia
	E-mail	
Author	Family name	Smerzi
	Particle	
	Given Name	Augusto
	Suffix	
	Division	
	Organization	INO-CNR
	Address	Largo Enrico Fermi 2, 50125, Firenze, Italy
	Division	European Laboratory for Non-Linear Spectroscopy (LENS)
	Organization	Università degli Studi di Firenze

	Address	Via Giovanni Sansone, 1, 50019 Sesto Fiorentino, Firenze, Italy
	E-mail	
Author	Family name	Sopuerta
	Particle	
	Given Name	Carlos
	Given Name	F.
	Suffix	
	Division	
	Organization	Institut de Ciències de l′Espai (ICE, CSIC)
	Address	Campus UAB, Carrer de Can Magrans s/n, 08193, Cerdanyola del Vallès, Spain
	Division	
	Organization	Institut d′Estudis Espacials de Catalunya (IEEC)
	Address	Edifici Nexus, Carrer del Gran Capità 2–4, despatx 201, 08034, Barcelona, Spain
	E-mail	
Author	Family name	Sorrentino
	Particle	
	Given Name	Fiodor
	Suffix	
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Sezione di Genova, via Dodecaneso 33, 16146, Genova, Italy
	E-mail	
Author	Family name	Sphicas
	Particle	
	Given Name	Paraskevas
	Suffix	
	Division	
	Organization	National and Kapodistrian University of Athens
	Address	Athens, Greece
	Division	
	Organization	CERN
	Address	CH-1211, Geneva, 23, Switzerland
	E-mail	
Author	Family name	Stadnik
	Particle	
	Given Name	Yevgeny
	Given Name	V.
	Suffix	
	Division	
	Organization	The University of Sydney
	Address	???, NSW, 2006, Australia
	E-mail	
Author	Family name	Stefanescu
	Particle	
	Given Name	Petruta
	Suffix	
	Division	
	Organization	Institute of Space Science
	Address	409 Atomistilor Street, Bucharest, Magurele, Ilfov, 077125,

		Romania
	E-mail	
Author	Family name	Tarallo
	Particle	
	Given Name	Marco
	Given Name	G.
	Suffix	
	Division	
	Organization	Istituto Nazionale di Ricerca Metrologica
	Address	Strada delle Cacce 91, Turin, 10035, Italy
	E-mail	
Author	Family name	Tentindo
	Particle	
	Given Name	Silvia
	Suffix	
	Division	High Energy Physics Group, 513 Keen Building, Dept. of Physics
	Organization	Florida State University
	Address	77 Chieflan Way, Tallahassee, FL, 32306, USA
	E-mail	
Author	Family name	Tino
	Particle	
	Given Name	Guglielmo
	Given Name	M.
	Suffix	
	Division	Dipartimento di Fisica e Astronomia
	Organization	Università di Firenze
	Address	Via Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	European Laboratory for Non-Linear Spectroscopy (LENS)
	Organization	Università degli Studi di Firenze
	Address	Via Giovanni Sansone, 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Sezione di Firenze, via Sansone 1, Sesto Fiorentino, Firenze, Italy
	Division	
	Organization	INO-CNR
	Address	Largo Enrico Fermi 2, 50125, Firenze, Italy
	E-mail	
Author	Family name	Tinsley
	Particle	
	Given Name	Jonathan
	Given Name	N.
	Suffix	
	Division	Dipartimento di Fisica e Astronomia
	Organization	Università di Firenze
	Address	Via Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	European Laboratory for Non-Linear Spectroscopy (LENS)
	Organization	Università degli Studi di Firenze
	Address	Via Giovanni Sansone, 1, 50019 Sesto Fiorentino, Firenze, Italy
	E-mail	
Author	Family name	Tornatore
	Particle	

	Given Name	Vincenza
	Suffix	
	Division	
	Organization	Politecnico di Milano
	Address	Piazza Leonardo da Vinci 32, 20133, Milano, Italy
	E-mail	
Author	Family name	Treutlein
	Particle	
	Given Name	Philipp
	Suffix	
	Division	Department of Physics
	Organization	University of Basel
	Address	???, Switzerland
	E-mail	
Author	Family name	Trombettoni
	Particle	
	Given Name	Andrea
	Suffix	
	Division	
	Organization	University of Trieste
	Address	Strada Costiera 11, 34151, Trieste, Italy
	E-mail	
Author	Family name	Tsai
	Particle	
	Given Name	Yu-Dai
	Suffix	
	Division	Department of Physics and Astronomy
	Organization	University of California
	Address	Irvine, CA, 92697-4575, USA
	E-mail	
Author	Family name	Tuckey
	Particle	
	Given Name	Philip
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE
	Address	61 avenue de l′Observatoire, 75014, Paris, France
	E-mail	
Author	Family name	Uchida
	Particle	
	Given Name	Melissa
	Given Name	A.
	Suffix	
	Division	
	Organization	University of Cambridge
	Address	J.J. Thomson Avenue, Cambridge, CB3 0HE, UK
	E-mail	
Author	Family name	Valenzuela
	Particle	
	Given Name	Tristan

	Suffix	
	Division	
	Organization	Rutherford Appleton Laboratory
	Address	Harwell Campus, Didcot, OX11 0QX, UK
	E-mail	
Author	Family name	Van Den Bossche
	Particle	
	Given Name	Mathias
	Suffix	
	Division	
	Organization	Thales Alenia Space
	Address	26 av. J.-F. Champollion, 31100, Toulouse, France
	E-mail	
Author	Family name	Vaskonen
	Particle	
	Given Name	Ville
	Suffix	
	Division	Institut de Fisica d′Altes Energies
	Organization	The Barcelona Institute of Science and Technology
	Address	Campus UAB, 08193, Bellaterra (Barcelona), Spain
	E-mail	
Author	Family name	Verma
	Particle	
	Given Name	Gunjan
	Suffix	
	Division	Dipartimento di Fisica e Astronomia
	Organization	Università di Firenze
	Address	Via Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	European Laboratory for Non-Linear Spectroscopy (LENS)
	Organization	Università degli Studi di Firenze
	Address	Via Giovanni Sansone, 1, 50019 Sesto Fiorentino, Firenze, Italy
	Division	
	Organization	Istituto Nazionale di Fisica Nucleare
	Address	Trieste Section, Via Valerio 2, 34127, Trieste, Italy
	E-mail	
Author	Family name	Vetrano
	Particle	
	Given Name	Flavio
	Suffix	
	Division	DiSPeA, Virgo Group
	Organization	University of Urbino
	Address	via S. Chiara, Urbino, I61029, Italy
	E-mail	
Author	Family name	Vogt
	Particle	
	Given Name	Christian
	Suffix	
	Division	
	Organization	Universität Bremen
	Address	Am Fallturm 2, 28359, Bremen, Germany

	E-mail	
Author	Family name	von Klitzing
	Particle	
	Given Name	Wolf
	Suffix	
	Division	
	Organization	Foundation for Research and Technology-Hellas
	Address	100 N. Plastira Street, Vassilika Vouton, GR 70013, Heraklion, Crete, Greece
	E-mail	
Author	Family name	Waller
	Particle	
	Given Name	Pierre
	Suffix	
	Division	
	Organization	European Space Agency
	Address	Keplerlaan 1, PO Box 299, NL-2200, AG Noordwijk, The Netherlands
	E-mail	
Author	Family name	Walser
	Particle	
	Given Name	Reinhold
	Suffix	
	Division	Institute for Applied Physics
	Organization	Technical University of Darmstadt
	Address	Hochschulstrasse 4A, D-64289, Darmstadt, Germany
	E-mail	
Author	Family name	Wille
	Particle	
	Given Name	Eric
	Suffix	
	Division	
	Organization	European Space Agency
	Address	Keplerlaan 1, PO Box 299, NL-2200, AG Noordwijk, The Netherlands
	E-mail	
Author	Family name	Williams
	Particle	
	Given Name	Jason
	Suffix	
	Division	Jet Propulsion Laboratory
	Organization	California Institute of Technology
	Address	Pasadena, CA, 91109, USA
	E-mail	
Author	Family name	Windpassinger
	Particle	
	Given Name	Patrick
	Suffix	
	Division	EQOQI, QUANTUM, Institute of Physics
	Organization	Johannes Gutenberg-University Mainz
	Address	55122, Mainz, Germany

	E-mail	
Author	Family name	Wittrock
	Particle	
	Given Name	Ulrich
	Suffix	
	Division	Photonics Laboratory, Engineering Physics
	Organization	Münster University of Applied Sciences
	Address	Stegerwaldstr. 39, 48565, Steinfurt, Münster, Germany
	E-mail	
Author	Family name	Wolf
	Particle	
	Given Name	Peter
	Suffix	
	Division	SYRTE, Observatoire de Paris
	Organization	Université PSL, Sorbonne Université, LNE
	Address	61 avenue de l ′ Observatoire, 75014, Paris, France
	E-mail	
Author	Family name	Woltmann
	Particle	
	Given Name	Marian
	Suffix	
	Division	
	Organization	Universität Bremen
	Address	Am Fallturm 2, 28359, Bremen, Germany
	E-mail	
Author	Family name	Wörner
	Particle	
	Given Name	Lisa
	Suffix	
	Division	German Aerospace Center (DRL)
	Organization	Institute of Quantum Technologies
	Address	Wilhelm-Runge-Straße 10, 89081, Ulm, Germany
	E-mail	
Author	Family name	Xuereb
	Particle	
	Given Name	André
	Suffix	
	Division	
	Organization	University of Malta
	Address	Msida, MSD, 2080, Malta
	E-mail	
Author	Family name	Yahia
	Particle	
	Given Name	Mohamed
	Suffix	
	Division	Faculty of Engineering and Natural Sciences
	Organization	International University of Sarajevo
	Address	Hrasnička cesta 15, 71210, Ilidža, Sarajevo, Bosnia-Herzegovina
	E-mail	
Author	Family name	Yazgan

	Particle	
	Given Name	Efe
	Suffix	
	Division	Department of Physics
	Organization	National Taiwan University
	Address	No. 1 Sect. 4 Roosevelt Road, Taipei, 10617, Taiwan
	E-mail	
Author	Family name	Yu
	Particle	
	Given Name	Nan
	Suffix	
	Division	Jet Propulsion Laboratory
	Organization	California Institute of Technology
	Address	Pasadena, CA, 91109, USA
	E-mail	
Author	Family name	Zahzam
	Particle	
	Given Name	Nassim
	Suffix	
	Division	DPHY, ONERA
	Organization	Université Paris-Saclay
	Address	Chemin de la Hunière—BP80100, F-91123, Palaiseau, France
	E-mail	
Author	Family name	Zambrini Cruzeiro
	Particle	
	Given Name	Emmanuel
	Suffix	
	Division	Instituto de Telecomunicações
	Organization	Instituto Superior Técnico
	Address	Av. Rovisco Pais, Torre Norte, Lisboa, 1049-001, Portugal
	E-mail	
Author	Family name	Zhan
	Particle	
	Given Name	Mingsheng
	Suffix	
	Division	Wuhan Institute of Physics and Mathematics, Innovation Academy for Precision Measurement Science and Technology
	Organization	Chinese Academy of Sciences
	Address	Wuhan, 430071, China
	E-mail	
Author	Family name	Zou
	Particle	
	Given Name	Xinhao
	Suffix	
	Division	LP2N, Laboratoire Photonique, Numérique et Nanosciences
	Organization	Université Bordeaux-IOGS-CNRS: UMR 5298
	Address	F-33400, Talence, France
	E-mail	
Author	Family name	Zupan
	Particle	
	Given Name	Jure

	Suffix	
	Division	
	Organization	University of Cincinnati
	Address	Cincinnati, Ohio, 45221, USA
	E-mail	
Author	Family name	Zupanič
	Particle	
	Given Name	Erik
	Suffix	
	Division	
	Organization	Jožef Stefan Institute
	Address	Jamova 39, 1000, Ljubljana, Slovenia
	E-mail	

Schedule	Received	12 April 2022
	Revised	
	Accepted	8 October 2022

Abstract

We summarize the discussions at a virtual Community Workshop on Cold Atoms in Space concerning the status of cold atom technologies, the prospective scientific and societal opportunities offered by their deployment in space, and the developments needed before cold atoms could be operated in space. The cold atom technologies discussed include atomic clocks, quantum gravimeters and accelerometers, and atom interferometers. Prospective applications include metrology, geodesy and measurement of terrestrial mass change due to, e.g., climate change, and fundamental science experiments such as tests of the equivalence principle, searches for dark matter, measurements of gravitational waves and tests of quantum mechanics. We review the current status of cold atom technologies and outline the requirements for their space qualification, including the development paths and the corresponding technical milestones, and identifying possible pathfinder missions to pave the way for missions to exploit the full potential of cold atoms in space. Finally, we present a first draft of a possible road-map for achieving these goals, that we propose for discussion by the interested cold atom, Earth Observation, fundamental physics and other prospective scientific user communities, together with the European Space Agency (ESA) and national space and research funding agencies.

Keywords

Footnotes

Angelo Bassi, Kai Bongs, Philippe Bouyer, Oliver Buchmueller, Luigi Cacciapuoti, Olivier Carraz, Maria Luisa Chiofalo, A. Michael Cruise, Albert De Roeck, Michael Doser, John Ellis, René Forsberg, Naceur Gaaloul, Richard Hobson, Sebastian Koke, Thomas Lévèque, Christian Lisdat, Federica Migliaccio, Eamonn Murphy, Ernst M. Rasel, Stephan Schiller, Christian Schubert, Carla Signorini, Guglielmo M. Tino, Wolf von Klitzing, Eric Wille, Peter Wolf and Lisa Wörner contributed equally to this work.

Alonso et al. *EPJ Quantum Technology* _####################_
https://doi.org/10.1140/epjqt/s40507-022-00147-w

EPJ.org

EPJ Quantum Technology
a SpringerOpen Journal

REVIEW Open Access

AUTHOR'S PROOF

Cold atoms in space: community workshop summary and proposed road-map

Iván Alonso[1], Cristiano Alpigiani[2], Brett Altschul[3], Henrique Araújo[4], Gianluigi Arduini[5], Jan Arlt[6], Leonardo Badurina[7], Antun Balaž[8], Satvika Bandarupally[9,10], Barry C. Barish[11], Michele Barone[12], Michele Barsanti[13], Steven Bass[14], Angelo Bassi[15,16†], Baptiste Battelier[17], Charles F.A. Baynham[4], Quentin Beaufils[18], Aleksandar Belić[8], Joel Bergé[19], Jose Bernabeu[20,21], Andrea Bertoldi[17], Robert Bingham[22,23], Sébastien Bize[18], Diego Blas[24,25], Kai Bongs[26†], Philippe Bouyer[17†], Carla Braitenberg[15], Christian Brand[27], Claus Braxmaier[28,27], Alexandre Bresson[19], Oliver Buchmueller[4,29†], Dmitry Budker[30,31], Luís Bugalho[32], Sergey Burdin[33], Luigi Cacciapuoti[34†], Simone Callegari[35], Xavier Calmet[36], Davide Calonico[37], Benjamin Canuel[17], Laurentiu-Ioan Caramete[38], Olivier Carraz[34†], Donatella Cassettari[39], Pratik Chakraborty[40], Swapan Chattopadhyay[41,42,31], Upasna Chauhan[43], Xuzong Chen[44], Yu-Ao Chen[45,46,47], Maria Luisa Chiofalo[13,48†], Jonathon Coleman[33], Robin Corgier[18], J.P. Cotter[4], A. Michael Cruise[26†], Yanou Cui[49], Gavin Davies[4], Albert De Roeck[50,5†], Marcel Demarteau[51], Andrei Derevianko[52], Marco Di Clemente[53], Goran S. Djordjevic[54], Sandro Donadi[16], Olivier Doré[55], Peter Dornan[4], Michael Doser[5†], Giannis Drougakis[56], Jacob Dunningham[36], Sajan Easo[22], Joshua Eby[57], Gedminas Elertas[33], John Ellis[7,5†], David Evans[4], Pandora Examilioti[56], Pavel Fadeev[30], Mattia Fanì[58], Farida Fassi[59], Marco Fattori[9], Michael A. Fedderke[60], Daniel Felea[38], Chen-Hao Feng[17], Jorge Ferreras[22], Robert Flack[61], Victor V. Flambaum[62], René Forsberg[63†], Mark Fromhold[64], Naceur Gaaloul[40†], Barry M. Garraway[36], Maria Georgousi[56], Andrew Geraci[65], Kurt Gibble[66], Valerie Gibson[67], Patrick Gill[68], Gian F. Giudice[5], Jon Goldwin[26], Oliver Gould[64], Oleg Grachov[69], Peter W. Graham[42], Dario Grasso[48], Paul F. Griffin[23], Christine Guerlin[70], Mustafa Gündoğan[71], Ratnesh K. Gupta[72], Martin Haehnelt[67], Ekim T. Hanımeli[73], Leonie Hawkins[33], Aurélien Hees[18], Victoria A. Henderson[71], Waldemar Herr[40], Sven Herrmann[73], Thomas Hird[29], Richard Hobson[4†], Vincent Hock[73], Jason M. Hogan[42], Bodil Holst[74], Michael Holynski[26], Ulf Israelsson[55], Peter Jeglič[75], Philippe Jetzer[76], Gediminas Juzeliūnas[77], Rainer Kaltenbaek[78], Jernej F. Kamenik[78], Alex Kehagias[79], Teodora Kirova[80], Marton Kiss-Toth[81], Sebastian Koke[35†], Shimon Kolkowitz[82], Georgy Kornakov[83], Tim Kovachy[65], Markus Krutzik[71], Mukesh Kumar[84], Pradeep Kumar[85], Claus Lämmerzahl[73], Greg Landsberg[86], Christophe Le Poncin-Lafitte[18], David R. Leibrandt[87], Thomas Lévèque[88†], Marek Lewicki[89], Rui Li[40], Anna Lipniacka[74], Christian Lisdat[35†], Mia Liu[90], J.L. Lopez-Gonzalez[91], Sina Loriani[92], Jorma Louko[64], Giuseppe Gaetano Luciano[93], Nathan Lundblad[94], Steve Maddox[81], M.A. Mahmoud[95], Azadeh Maleknejad[5], John March-Russell[29], Didier Massonnet[88], Christopher McCabe[7], Matthias Meister[27], Tadej Mežnaršič[75], Salvatore Micalizio[37], Federica Migliaccio[96†], Peter Millington[64], Milan Milosevic[54], Jeremiah Mitchell[67], Gavin W. Morley[97], Jürgen Müller[40], Eamonn Murphy[34†], Özgür E. Müstecaplıoğlu[98], Val O'Shea[99], Daniel K.L. Oi[23], Judith Olson[100], Debapriya Pal[101], Dimitris G. Papazoglou[102], Elizabeth Pasatembou[4], Mauro Paternostro[103], Krzysztof Pawlowski[104], Emanuele Pelucchi[105], Franck Pereira dos Santos[18], Achim Peters[71], Igor Pikovski[106,107],

UNCORRECTED PROOF

© The Author(s) . **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 2/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

Apostolos Pilaftsis[108], Alexandra Pinto[109], Marco Prevedelli[110], Vishnupriya Puthiya-Veettil[56], John Quenby[4], Johann Rafelski[111], Ernst M. Rasel[40†], Cornelis Ravensbergen[100], Mirko Reguzzoni[96,50], Andrea Richaud[112], Isabelle Riou[81], Markus Rothacher[113], Albert Roura[27], Andreas Ruschhaupt[105], Dylan O. Sabulsky[17], Marianna Safronova[114], Ippocratis D. Saltas[115], Leonardo Salvi[9,10,116], Muhammed Sameed[108], Pandey Saurabh[57], Stefan Schäffer[117], Stephan Schiller[118†], Manuel Schilling[40], Vladimir Schkolnik[71], Dennis Schlippert[40], Piet O. Schmidt[35,40], Harald Schnatz[35], Jean Schneider[119], Ulrich Schneider[67], Florian Schreck[117], Christian Schubert[40†], Armin Shayeghi[120], Nathaniel Sherrill[36], Ian Shipsey[29], Carla Signorini[13,48†], Rajeev Singh[121], Yeshpal Singh[26], Constantinos Skordis[122], Augusto Smerzi[123,10], Carlos F. Sopuerta[124,125], Fiodor Sorrentino[126], Paraskevas Sphicas[127,5], Yevgeny V. Stadnik[128], Petruta Stefanescu[38], Marco G. Tarallo[37], Silvia Tentindo[129], Guglielmo M. Tino[9,10,116,123†], Jonathan N. Tinsley[9,10], Vincenza Tornatore[96], Philipp Treutlein[130], Andrea Trombettoni[15], Yu-Dai Tsai[131], Philip Tuckey[18], Melissa A. Uchida[67], Tristan Valenzuela[22], Mathias Van Den Bossche[132], Ville Vaskonen[133], Gunjan Verma[9,10,16], Flavio Vetrano[134], Christian Vogt[73], Wolf von Klitzing[56†], Pierre Waller[34], Reinhold Walser[135], Eric Wille[34†], Jason Williams[55], Patrick Windpassinger[136], Ulrich Wittrock[137], Peter Wolf[18†], Marian Woltmann[73], Lisa Wörner[27†], André Xuereb[138], Mohamed Yahia[139], Efe Yazgan[140], Nan Yu[55], Nassim Zahzam[19], Emmanuel Zambrini Cruzeiro[32], Mingsheng Zhan[141], Xinhao Zou[17], Jure Zupan[142] and Erik Zupanič[75]

*Correspondence: Oliver.Buchmueller@cern.ch; John.Ellis@cern.ch
[4]Imperial College London, Prince Consort Road, London, SW7 2AZ, UK
[7]King's College London, Strand, London, WC2R 2LS, UK
Full list of author information is available at the end of the article
†Section Editor and/or Workshop Organiser

Abstract

We summarize the discussions at a virtual Community Workshop on Cold Atoms in Space concerning the status of cold atom technologies, the prospective scientific and societal opportunities offered by their deployment in space, and the developments needed before cold atoms could be operated in space. The cold atom technologies discussed include atomic clocks, quantum gravimeters and accelerometers, and atom interferometers. Prospective applications include metrology, geodesy and measurement of terrestrial mass change due to, e.g., climate change, and fundamental science experiments such as tests of the equivalence principle, searches for dark matter, measurements of gravitational waves and tests of quantum mechanics. We review the current status of cold atom technologies and outline the requirements for their space qualification, including the development paths and the corresponding technical milestones, and identifying possible pathfinder missions to pave the way for missions to exploit the full potential of cold atoms in space. Finally, we present a first draft of a possible road-map for achieving these goals, that we propose for discussion by the interested cold atom, Earth Observation, fundamental physics and other prospective scientific user communities, together with the European Space Agency (ESA) and national space and research funding agencies.

UNCORRECTED PROOF

1 Preface

This document contains a summary of the *Community Workshop on Cold Atoms in Space* [1] that was held virtually on September 23 and 24, 2021. The purpose of this community workshop was to discuss objectives for a cold atom quantum technology development programme coordinated at the Europe-wide level, and to outline a possible community road-map and milestones to demonstrate the readiness of cold atom technologies in space, as proposed in the recommendations of the Voyage 2050 Senior Science Committee (SSC) [2], and in synergy with European Union (EU) programmes.

The SSC was set up by the ESA Director of Science to advise on the space science programme for the period 2030–2050, and drew attention to the potential of cold atom tech-

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 3/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 1 Statistical analysis of the 331 registered to participate in the Community Workshop on Cold Atoms in Space, held virtually on Sept. 23rd and 24th, 2021, and/or to follow the Community Road-map process. Upper left: Breakdown of registered participants into the prospective User community and those working on Cold Atom technology. Upper right: Breakdown of registered participants by self-declared primary research area. Lower left: Breakdown of registered participants by self-declared secondary research area. Lower right: Breakdown of registered participants by geographical location of institute where they are affiliated

UNCORRECTED

nology in fundamental physics and planetary science as well as in navigation, timekeeping and Earth Observation (EO) [2]. The SSC set out a plausible programme of technology development in the Voyage 2050 report that would prepare cold atom payloads for evaluation by the ESA science committees on scientific merit alone, without technical concerns about robustness for the space environment. One aim of the workshop in September 2021 was to engage the cold atom community in defining possible science payloads that might be used to establish a recognised pathway towards the use of cold atoms in the ESA science programme.

This community workshop brought together representatives of the cold atom, astrophysics, cosmology, fundamental physics, geodesy and earth observation communities to participate in shaping this development programme. It built upon one organised two years ago [3], which reviewed the landscape of present and prospective cold atom experiments in space. Subsequently, several White Papers were submitted [4–12] in response to the Voyage 2050 call ,[1] which outlined possible ultimate goals and reviewed experiments and technical developments underway that could help pave a way towards these goals.

One of the key objectives of this Workshop was to enhance the strength and coherence of a community embracing Cold Atom technology experts as well as prospective Users. In total, 331 people from 36 countries registered as participants in the Workshop and/or to follow the community roadmap process. As seen in the two upper panels of Fig. 1, the registered participants comprise essentially equal numbers of declared cold atom experts

[1] See, in particular, the Special Issue (Vol. 51) of *Experimental Astronomy* devoted to Voyage 2050 science themes for ESA's long-term plan for the science programme edited by K. O'Flaherty, L. Tacconi, C. Arridge, G. Hasinger and F. Favata.

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 4/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

and prospective Users. The upper right panel breaks the user community down by their self-declared primary research areas, which range from Earth Observation to cosmology, and the lower left panel provides another breakdown by their secondary research areas. As anticipated for a Europe-based event, about 80% of the registrations are from European countries, as seen in the lower right panel of Fig. 1, but there is also significant North American and Asian participation.

This community provides the backbone for long-term planning and the support needed to see the challenging cold atom missions foreseen in the road-map through to their successful completions. The participants display an excellent and diverse mix of expertise, building an outstanding basis for the success of the workshop and the development of the corresponding road-map, which defines possible interim and long-term scientific goals and outlines technological milestones. Section 2 is an introduction to our document, Sects. 3 to 5 discuss our main science themes, namely atomic clocks, Earth Observation and fundamental physics, Sect. 6 discusses the technology developments required before quantum sensors can be deployed in space, Sect. 7 summarises the discussions during the Workshop, and in Sect. 8 we outline the corresponding Community road-map.

2 Introduction

Quantum physics was developed primarily in Europe in the first half of the 20th century. In the second half, the first "quantum revolution" took place and was the engine of the main technological and societal transformations in recent decades considering, e.g., solid-state electronics and hence all information and computing technologies. It also enabled the space era thanks, e.g., to onboard semiconductor technologies (solar cells, avionics, communication systems radars, detectors, etc.). Similarly, the first half of the 21st century is being deeply impacted by the second "quantum revolution", with the possibility of exploiting and controlling quantum phenomena so far not applied outside the laboratory, e.g., macroscopic quantum coherence, superposition, entanglement.

Atomic quantum sensors are a newly-emerging technology of unparalleled accuracy and precision. Spaceborne quantum inertial sensors (e.g., accelerometers, gravimeters, gyroscopes) are today the most advanced sensing technologies that benefit from this revolution, exploiting matter-wave interferometry with Bose–Einstein condensates, using atom clouds cooled below nanoKelvin temperature. For example, whereas classical accelerometers suffer from high noise at low frequencies, cold atom interferometers (CAI) are highly accurate also at low frequencies and do not need any external calibration: see Fig. 5 below.

In the past twenty years, gravimetry missions have demonstrated a unique capability to monitor major climate-related changes of the Earth directly from space—quantifying the melt of large glaciers and ice sheets, global sea level rise, continental drought, major flooding events, and also the effects of large earthquakes and tsunamis. Adding to fundamental knowledge of the Earth, a quantum gravimetry mission for climate will provide essential climate variables (ECVs) of unprecedented quality for groundwater, mass balance of ice sheets and glaciers, heat and mass transport, building upon the successes of previous missions using classical technology, such as the Gravity field and steady-state Ocean Circulation Explorer (GOCE) [13], the Gravity Recovery and Climate Experiment mission (GRACE) [14] and its follow-on mission (GRACE-FO) [15]. A combination of classical sensors with CAI or, at a later stage, a full quantum sensor will bring the Quantum Mission for Climate to a sensitivity that will open many applications and satisfy user

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 5/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

needs [16] with respect to water management and hazard prevention. In this connection, we take special note of the adoption of Quantum Technology for Earth Observation by the European Commission, notably (but not exclusively) in the Horizon Europe programme, under the thrust of Commissioner T. Breton, and of the inclusion of Quantum Technology in the ESA Agenda 2025 [17].

Quantum Technology on Earth has revolutionised the measurement of time since the first atomic clocks in the 1950s, and these now provide the fundamental time frame across the globe. In space, atomic clocks have widespread applications such as satellite-based navigation systems (GPS, GALILEO). Terrestrial clocks based on atomic transitions are now reaching a relative accuracy on the order of 10^{-18}, a level at which a change of height in the Earth's gravitational field of 1 cm would be detectable as a gravitational redshift. This sensitivity brings both challenges and opportunities. The challenge for terrestrial clocks will be that changes in the local gravitational potential, either by human activity or by alterations in the local water table, will destroy the stability of the clock. This issue will certainly drive the siting of such clocks in space, with the implication that space qualification of the quantum technology will be essential for future development. The availability of such sensitive technology in space also offers significant opportunities to explore many aspects of fundamental physics.

Mounted on a space platform in a highly eccentric orbit, a sensitive atomic clock would provide an ideal laboratory to test the Einstein Equivalence Principle (EEP) of General Relativity beyond current precision as the spacecraft experiences varying gravitational potentials around the orbit. This is a test that is at the heart of General Relativity and all metric theories of gravitation and space-time. A fundamental aspect of the EEP of General Relativity is the Universality of Free Fall (UFF), which has been tested since the days of Galileo with ever-increasing accuracy. Quantum gravimetry using atom interferometers in space will allow pushing tests of UFF to new frontiers, with the potential of unveiling new physics beyond the Standard Model that modify our theory of gravity, e.g., via fifth forces that exhibit screening mechanisms. These experiments represent one of the best ways of exploring the unknown theoretical interface between quantum physics and our best-tested theory of gravity, General Relativity.

The deployment of cold atom technology in space will also enable many other sensitive experiments in fundamental physics, cosmology and astrophysics, such as searches for ultralight dark matter particles, measurements of gravitational waves from the mergers of massive black holes and phenomena in the early Universe, and ultrasensitive probes of quantum mechanics, complementing terrestrial laboratory experiments such as those at particle colliders.

The commonality of many subsystems between atomic clocks, gravimeters and fundamental physics experiments means that a well-planned programme of technical development should lead to the availability in space-borne missions of all these applications in fundamental science, Earth Observation, time keeping and navigation.

3 Atomic clocks review

3.1 Scientific and societal opportunities

3.1.1 Fundamental science

High-stability and -accuracy atomic clocks combined with state-of-the-art time and frequency links can be used to measure tiny variations in the space-time metric and test the validity of the Einstein's Equivalence Principle.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 6/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

As predicted by General Relativity, gravity influences the flow of time. When identical clocks experiencing a different gravitational potential are compared by exchanging timing signals, a relative frequency difference proportional to the difference of the gravitational potential at the location of the clocks can be measured. The effect, known as gravitational redshift, has been tested in 2018 with a precision of about 2×10^{-5} [18, 19] by using the clocks on-board the Galileo 5 and 6 satellites. The ACES (Atomic Clock Ensemble in Space) mission [20–22] will perform an absolute measurement of the redshift effect between the PHARAO clock on-board the International Space Station (ISS) and clocks on Earth, improving this limit by an order of magnitude. Optical clock missions on highly elliptical orbits around the Earth or cruising towards the Sun are expected to improve redshift tests by several orders of magnitude and to measure higher-order relativistic effects to high precision.

Local Lorentz Invariance (LLI) postulates the independence of any local test experiment from the velocity of the freely-falling apparatus. Optical clocks can be used to provide very stringent test of Lorentz symmetry and the Lorentz-Violating Standard Model Extension (SME) [23]. Distant Sr optical lattice clocks compared through optical fibre links have been used to constrain to 1×10^{-8} the Robertson–Mansouri-Sexl parameter used to test theories of special relativity (see [24] for a review) by searching for daily variations of the relative frequency difference [25]. In [26], two Yb^+ clocks confined in two traps with quantization axis aligned along non-parallel directions are compared while the Earth orbits around the Sun. The absence of frequency modulations at the level of 1×10^{-19} made possible an improvement in the limits on the Lorentz symmetry violation parameter for electrons.

Local Position Invariance (LPI) can also be tested by comparing clocks based on different atomic transitions. According to LPI, the outcome of any local test experiment is independent of where and when it is performed in the Universe. Transition frequencies depend differently on three fundamental constants: the fine structure constant α, the electron mass m_e/Λ_{QCD} (where Λ_{QCD} is the QCD scale parameter), and the quark mass m_q/Λ_{QCD}. Therefore, comparing atomic clocks based on different transitions can be used to constrain the time variation of fundamental constants and their couplings to gravity. As an example, the comparison of two $^{171}Yb^+$ clocks based on the electric quadrupole and electric octupole transitions and two Cs clocks repeated over several years has recently improved the limits on the time variation of the fine structure constant and of the electron-to-proton mass ratio [27]. At the same time, using the annual variation of the Sun's gravitational potential, it was possible to constrain the coupling of both constants to gravity.

Atomic clock networks can also be used to place bounds on Topological Dark Matter (TDM) models. TDM can be expressed as a scalar field that couples to fundamental constants such as the fine-structure constant and the electron mass, thus producing variations in the transition frequencies of atomic clocks at its passage. Cross-comparisons between atomic clocks connected in a network over large distances can be used to place bounds on the time variation of three fundamental constants and determine exclusion regions for the effective energy scale (inverse of the coupling strength) of the dark matter field as a function of its Compton wavelength [28–31]. Clock networks providing redundant measurements are a powerful tool to control systematic effects and confirm any detection above the noise threshold.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 7/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

The dark matter distribution in the Solar system is critical for the reach of dark matter direct detection experiments. It is possible that some local over-density of ultralight dark matter could be bound to the Sun. A clock-comparison satellite mission with two clocks onboard sent to the inner reaches of the solar system was proposed in [32] to search for the dark matter halo bound to the Sun, probe natural relaxion[2] parameter space, and look for the spatial variation of the fundamental constants associated with a change in the gravitation potential.

Optical clocks have also been proposed for gravitational wave detection [4, 5, 34]. A pair of clocks in drag-free satellites separated by a long-distance baseline share the interrogation laser via an optical link. The clocks act as narrowband detectors of the Doppler shift on the laser frequency due to the relative velocity between the satellites induced by the incoming gravitational wave. The atom interrogation sequence on the clock transition can be controlled, enabling precise tuning of the detection window over a wide frequency interval without loss of sensitivity. A frequency range between about 10 mHz and 10 Hz can be covered, thus bridging the gap between space-based and terrestrial optical interferometers, as discussed in more detail in Sect. 5.

3.1.2 Metrology

The basic 'second' in the International System of Units (SI) is the quantity that is fixed with by far the lowest uncertainty of all units. This is done by primary frequency standards (laser-cooled Cs fountain clocks) operated at National Metrology Institutes. Global time scales rely on the comparison of such high-performance atomic clocks connected in a global network. The Bureau International des Poids et Mesures (BIPM) generates International Atomic Time (TAI), which is based on the cross-comparison of the best primary frequency standards and, more recently, also optical clocks worldwide. Its rate is defined to be close to proper time on the geoid. Coordinated Universal Time (UTC) is produced by the BIPM and differs from TAI only by an integral number of seconds as published by the BIPM: it is the only recommended time scale for international reference and the basis of civil time in most countries.

Since optical clocks already outperform the primary frequency standards that operate in the microwave domain (see Sect. 3.2), the international metrology community represented by the Comité International des Poids et Mesures (CIPM) and its committees have devised a road-map for the redefinition of the second. This documents the high priority and strong commitment of a large community to the development and operation of optical clocks, with high relevance for society. Such a redefinition will enable a more accurate and stable international timescale [35, 36], which is key for precise navigation services via the global navigation satellite system (GNSS) network, the synchronization of worldwide exchanges and markets, communication networks, and national defence and security.

The coordination of time requires the permanent comparison and synchronisation of national timescales and clocks. With the increasing performance of optical clocks, the demands on the link quality are also increasing. Today's microwave links achieve neither the necessary stability nor the accuracy required by the new optical clocks [37]. Locally, fibre-optical links can be an alternative [38], but a global network is not within reach. Long-distance time and frequency links enabling frequency comparisons at the level of 1×10^{-18}

[2] The relaxion is a light spin-zero field that dynamically relaxes the Higgs mass with respect to its natural large value [33].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 8/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

are urgently needed. Such local networks may even be combined by space clocks, as in the ACES [21] or the proposed Space Optical Clock (SOC) [39] missions. Potentially, a space clock can overcome the limitations on the realization of the SI second and of timescales set by the knowledge of the gravity potential on the ground because the relativistic frequency shift, needed to transform the proper time of the clock to TAI, can be computed more accurately for a space clock than for a ground clock. Presently, this correction can only be determined with a fractional uncertainty of about 3×10^{-18}, equivalent to 3 cm height [40], which is already larger than the uncertainty of today's optical clocks (see Sect. 3.2.1).

3.1.3 Earth observation & geodesy

In view of climate change and its consequences for society, Earth observation and geodesy are of increasing importance. There exist highly accurate geometric reference frames based on the GNSS, Very Long Baseline Interferometry (VLBI), Satellite Laser Ranging (SLR) and Doppler Orbitography and Radiopositioning Integrated by Satellite (DORIS) networks [41, 42]. Physical height reference systems related to the geoid, which are sensitive to the flow of water, are much less accurate and fall behind the requirements set by the UN resolution for sustainable development [43] by more than an order of magnitude.

Presently, physical heights are locally derived by spirit levelling tied to reference points such as tide gauges or global observations from satellite missions like CHAMP [44], GOCE [13], GRACE [45], and GRACE-FO [15]. Although these missions were and are very successful, they lack spatial resolution and require considerable data processing, because the sensors are only sensitive to derivatives of the gravity potential. With clocks at a fractional uncertainty level of 10^{-18}, we now have sensors at hand that are directly sensitive to the gravity potential via the relativistic redshift [46, 47]. Therefore, we have an opportunity to establish a novel technique to realize a height reference system using a network of optical clocks from which the physical height differences at the respective locations can be derived.

Present clock performance already provides a height resolution better than the current geodetic state-of-the-art [40], and is likely to reach the millimetre level within the next decade. It is essential to establish links for cross-comparisons of optical clocks that are flexibly accessible and span the globe. Satellite-based approaches fulfil these requirements in an ideal fashion. While satellite-mediated ground-to-ground links with improved performance compared to, e.g., ACES, will enable a fast development of this field of application. Clocks operated in space are expected to improve the products from GRACE-like missions, in terms of estimates of potential coefficients, particularly in the low-frequency range of the spherical harmonics. If a suitable satellite formation is exploited, an improvement in the estimation of time variations in the potential could also be achieved. Space clocks may ultimately provide an independent, long-term stable and reproducible height reference for decades, even centuries of Earth monitoring.

3.2 Clocks: state of the art

3.2.1 Laboratory-based clocks

Figure 2 shows the historical progress of state-of-the-art laboratory atomic clocks of different types, distinguished by coloured symbols of different shapes. Cesium (Cs) microwave atomic clocks have been the primary standard for the SI second since 1967, which has helped to motivate the development of several generations of Cs clocks with reduced fractional frequency errors. However, in recent years, optical atomic clock technology has

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 9/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 2 Progress in the relative accuracy of atomic clocks [26, 27, 48, 51–79]. Cs microwave clocks have steadily improved since the emergence of laser-cooled fountain clocks in the early 1990s [80–82], but two distinct types of optical clock currently compete at a fractional frequency uncertainty of approximately 1×10^{-18}: trapped-ion clocks (Yb^+ [26], Al^+ [48], Hg^+ [68], Sr^+ [70], Ca^+ [83]) and optical lattice clocks (Sr [78], Yb [77], Hg [76], Cd [84])

matured significantly: the best optical atomic clocks now surpass Cs in relative accuracy by a factor of more than 100. The field of optical clocks encompasses a diverse range of trapped-ion clocks and optical lattice clocks, each with distinct merits. Trapped ion clocks naturally offer exquisite environmental isolation and quantum control, leading to small systematic shifts and high accuracy [26, 48]. However, optical lattice clocks have the key advantage of using many atoms in parallel, resulting in greater frequency stability and therefore allowing high-precision measurements within a significantly shorter averaging time [49, 50].

In order to verify the 10^{-18} precision of the best optical clocks and pursue the scientific opportunities discussed in Sect. 3.1, it is important to compare different clocks of comparable precision with each other. Local measurements of optical clocks within a single laboratory have been used to measure general relativistic redshifts corresponding to vertical displacements at the millimetre scale [85] and to search for possible variations in local physics induced by dark matter [64, 86, 87]. Distant comparisons between clocks in separate laboratories allow large numbers of independent clocks to be included, but must be mediated by a complex frequency link infrastructure. The longest distances are spanned by satellite links, which allow comparison at 10^{-16} fractional frequency uncertainty [37]. However, much higher precision at the 10^{-18} level can be carried out over shorter distances using terrestrial links, either with free-space lasers [88] or telecoms-wavelength lasers sent through optical fibres [38, 89, 90]. The most extensive optical fibre network is operated between European metrology institutes [38], across which several clocks have been compared to search for physics beyond the Standard Model (see Sect. 3.1.1) [25, 91].

The development of laboratory atomic clocks is fuelled by a broad community that spans universities, industry and several National Metrology Institutes. Optical lattice clocks are now particularly widespread, with more than a dozen strontium (Sr) [63, 75, 89, 92–102]

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 10/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

and some ytterbium (Yb) [102–106] clock laboratories in operation worldwide.[3] The commitment of the metrology community to continue investing in optical clocks is highlighted by the CIPM road-map for an optical redefinition of the SI second (see Sect. 3.1.2). The road-map mandates a research programme likely to span at least the next decade, in which several optical clocks will be developed at 1×10^{-18} relative accuracy and validated through clock-clock measurements. To carry out such measurements, the priority of the optical clock community will be to develop cold atom technology with higher technology readiness levels (TRLs), capable of combining state-of-the-art accuracy with robust, long-term operation—an investment which should have close synergies with a future programme for cold atoms in space.

3.2.2 *Transportable clocks*

Early in the development of optical frequency standards it was recognized that mobile devices (see, e.g., [107]) enable applications (see Sect. 3.1) of clocks that are impractical if the availability of clocks is restricted to only a few laboratories. The required engineering to develop delicate laboratory systems into robust mobile devices also opens the door to commercialization and space applications of clocks. While there have been several impressive demonstrations of compact optical frequency standards with high performance [108, 109], we focus here on activities that target a clock performance similar to the state-of-the-art of laboratory setups (see Sect. 3.2.1).

To maintain the outstanding frequency stability of optical clocks, ultra-stable interrogation lasers are required. For reasons of seismic and thermal insulation, these are typically neither robust nor compact. Therefore, the further development of these devices was identified by the community as an important challenge [110–112] and supported, e.g., by ESA activities [113–115] and is—with demonstrated fractional frequency instabilities significantly below 10^{-15}—on a good path. As ultra-stable laser systems have numerous applications beyond optical frequency standards, e.g., in atom interferometry, ultra-stable microwave generation, or optical telecommunication, the continued support of these activities is of high importance.

The realization of a fully transportable optical clock requires more lasers and a complex technical infrastructure (see Fig. 3), and thus poses a larger challenge. Nevertheless, several such systems working with neutral atoms [116–118] or single ions [83] have been realized, which already outperform the most accurate microwave standards. These setups are developed for space applications [39], and have been used in a geodetic context [119, 120] or to test fundamental aspects of physics [79]. We therefore conclude that the construction and reliable operation of optical clocks with fractional uncertainties of 1×10^{-17} and below and compact dimensions with volumes $\lesssim 1\ \mathrm{m}^3$ is already possible today [121, 122].

3.2.3 *Free space-time and frequency links*

Connecting (optical) atomic clocks worldwide lays the basis for applications such as the creation of TAI or a Positioning, Navigation and Timing (PNT) standard, and would also open the route to testing theories of fundamental physics (see Sects. 3.1 and 3.3).

[3] Importantly, these optical lattice clocks use the same Sr and Yb technology as proposed for atom-interferometer science missions such as AEDGE [5].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 11/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

Currently, primary microwave clocks are connected via satellites [37] in the microwave domain by the existing GNSS infrastructure [123] or dedicated two-way time and frequency transfer (TWTFT) links [124–126]. Demonstrated frequency transfer uncertainties of existing microwave links (MWLs) reach into the 10^{-16} range after averaging times of days [123, 127] and into the 10^{-17} range after averaging times of weeks with integer ambiguity resolution [123], and demonstrated time transfer uncertainties lie in the nanosecond region [126]. A new generation of MWL equipment is under development [21, 128], which reaches in laboratory tests timing instabilities of < 100 fs for averaging times τ = 10 s to 2000 s [21], which is equivalent to fractional frequency transfer uncertainties of $< 5 \times 10^{-17}$ at 2000 s. Similar performances are achieved in the optical domain by Time Transfer by Laser Link (T2L2) [129] and the European Laser Timing (ELT) experiment [130] employing time-of-arrival measurements of laser pulses.

A significantly improved uncertainty is achieved by techniques exploiting the optical carrier. Optical frequency dissemination using continuous wave laser signals [131] reaches fractional frequency transfer instabilities $< 5 \times 10^{-19}$ already after 100 s of averaging time in path-length stabilized operation [132]. A team at NIST has developed an optical TWTFT (OTWTFT) technique [88] combining carrier and time-of-flight information, allowing phase-coherent averaging over the signal dropouts that occur inevitably due to atmospheric turbulence [133]. Using this technique, the NIST team has demonstrated sub-10^{-18} frequency transfer uncertainty and sub-1 fs timing uncertainty at an averaging time of 1000 s in a 3-node network of two concatenated 14 km links [134]. Furthermore, they demonstrated OTWTFT to a flying drone with similar performance [135]. Despite the proven performance, however, the remaining steps to achieve ground-to-satellite world-wide coverage remain challenging: demonstrate techniques for higher relative speeds between sender and receiver, assess the impact of atmospheric turbulence, signal loss, potential loss of reciprocity on such ground-to-satellite links, and consider inclusion of relativistic effects. Recently, a first study addressed this scaling to ground-to-satellite connections [136] and came to a positive conclusion regarding the feasibility. Nevertheless, further experimental evidence gradually approaching the long-term ground-to-satellite goal is required. Synergies can be expected with the proposed combination of microwave and optical links in the context of new GNSS constellations [137].

3.3 International space activities

Space is the ideal laboratory to test general relativity and alternative theories of gravitation with atomic clocks. The large velocities and velocity variations, the access to large variations in the gravitational potential, and the possibility to establish a global network able to compare ground clocks across continents from space provide new opportunities both for fundamental physics research and for applications in other areas of research, such as clock synchronization and timescale distribution, geodesy, Earth observation, navigation, etc., as discussed elsewhere in this report.

ACES (Atomic Clock Ensemble in Space) [21] is an ESA mission designed to operate on the International Space Station. The two on-board clocks rely on atomic transitions in the microwave domain. The PHARAO clock, a primary frequency standard based on laser cooled Cs atoms, provides the ACES clock signal with a long-term stability and accuracy of 1×10^{-16} in fractional frequency; the active H-maser SHM is the on-board flywheel oscillator that will be used for the characterization of the PHARAO accuracy. The ACES

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 12/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 3 (Left) Flight model of the ACES payload during the assembly phase. SHM (vertical cylinder) and PHARAO (in the background) are installed on the bottom panel. The ACES computer, the PHARAO computer, and the on-board phase comparator are visible on the left panel. MWL electronics (not installed yet) and the antennae are accommodated on the top panel. (Right) The ACES payload installed on the Columbus External Payload Adapter (CEPA) for interface tests

clock signal is distributed to ground clocks by using two time and frequency links: MWL is a link in the microwave domain; ELT is an optical link using short laser pulses to exchange timing signals. A distributed network of MWL ground terminals will connect the clocks operated in the best research institutes worldwide (SYRTE, PTB, NPL and Wettzell in Europe, NIST and JPL in the US, NICT in Japan) to the ACES clock signal. Satellite laser ranging stations will also be connected to the clock network by using the ELT optical link. The space-to-ground clock de-synchronization measurement produced by MWL and ELT will be used to perform an absolute measurement of the gravitational redshift in the field of the Earth to < 2 ppm, to probe time variations of fundamental constants, and to perform tests of the Lorentz-Violating Standard Model Extension (SME) [23]. The possibility of searching for topological dark matter with the ACES network is also being investigated. ACES is expected to fly to the ISS in the 2025 time frame. The flight model of the ACES payload is shown in Fig. 3.

Significant advances in the development of microwave cold-atom clocks have been achieved in China with the launch and on-orbit operation of the CACES (Cold Atom Clock Experiment in Space) clock based on laser-cooled Rb atoms [138]. The experiment was successfully operated on the Chinese space laboratory Tiangong-2. From an analysis of the Ramsey fringes, an estimated short-term stability at the level of $3 \times 10^{-13}/\sqrt{s}$ was attained under free fall conditions. Unfortunately, a full characterization of the clock in space was not possible due to the absence of a stable frequency reference and of a space-to-ground link on board Tiangong-2. The clock performance is still to be optimised, but this experiment clearly demonstrates the robustness of the cold-atom clock technology for space.

In the US, the NASA's Jet Propulsion Lab has successfully demonstrated mercury trapped-ion clock technology in space [139]. The Deep Space Atomic Clock (DSAC) payload, consisting of a Hg^+ microwave clock and a dedicated GPS receiver, was launched

UNCORRECTED PROOF

AUTHOR'S PROOF

into a 720-km orbit around the Earth in June 2019. The space clock was compared to the clocks from the US Naval Observatory, and demonstrated a fractional frequency stability between 3×10^{-15} and 5×10^{-15} at 1 day, subsequently stabilizing at 3×10^{-15} as measured after 23 days [139]. The short-term stability of the clocks, which is below the GPS measurement system noise, could be estimated as $7 \times 10^{-13}/\tau^{1/2}$, where τ is the integration time. This technology can be used for navigation, planetary science, and fundamental physics.

As discussed above, optical clocks can provide an improvement in stability and accuracy of 2 orders of magnitude with respect to microwave clocks. Following the impressive progress of atomic clocks based on optical transitions, several initiatives are currently ongoing to advance the required technology to flight readiness.

Europe is developing key optical clock technology for space, e.g., cooling lasers, the clock laser, a high-finesse reference cavity, a clock control unit to stabilize the laser frequency on the atomic transition, and the lattice laser. A design study for a Sr clock physics package has been completed. Compact and transportable ground-based prototypes for a Sr optical lattice clock [116] and a Sr ion clock [140] are being characterized. Free-space coherent optical links reaching a fractional frequency uncertainty of 1×10^{-19} in a few days of measurement time are under development [132]. In parallel, the I-SOC Pathfinder platform has been proposed as the ACES follow-on mission. I-SOC Pathfinder is pushing further the microwave and optical link technology [141, 142] developed for ACES to continue operating a worldwide network of optical clocks on the ground to test fundamental laws of physics, to develop applications in geodesy and time & frequency transfer, and to demonstrate key time and frequency link technologies that are essential for all future atomic clock missions in space. The I-SOC Pathfinder mission could be operated on the ISS, but other platforms as geostationary satellites could be even preferable to enhance common view time slots between selected ground stations.

In the US, the FOCOS (Fundamental physics with an Optical Clock Orbiting in Space) mission concept has been proposed [122]. FOCOS relies on a Yb optical lattice clock with 1×10^{-18} stability and accuracy on a highly elliptical orbit around the Earth. A coherent optical link is used to compare the space clock to ground clocks for general relativity tests and timing applications. FOCOS will also serve as a pathfinder for future atom interferometry missions to test the Equivalence Principle, clock constellations in space to hunt for dark matter [28, 29], and gravitational wave observatories [34].

In parallel, CACES follow-on experiments based on optical clock technology are under development in China. It is expected that the Chinese space program will continue experiments including clocks over the coming five years, on the Tiangong space station, the Chang'e series of lunar probes, and the Tianwen-1 Mars probe [143].

Finally, major efforts and resources are being invested worldwide to improve the atomic clocks of the Global Navigation Satellite System (GNSS). Currently, available technology relies on the passive H-maser, the Rb atomic frequency standard, and the Cs beam frequency standard [144, 145]. Clocks for navigation satellites have reduced stability (in the 10^{-15} – 10^{-14} range for fractional frequency), but offer a more compact design with low mass (3 to 20 kg), low power consumption (30 to 70 W), and long lifetime on orbit. Alternative technologies are under study for the next generation of atomic clocks for navigation. Among them, it is worth mentioning the mercury ion clock technology [139], the pulsed optically pumped Rb clock [146], the Rb optical atomic clock [147], and the iodine

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 14/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

frequency reference [109, 148] that will soon be tested in the COMPASSO experiment on the ISS platform Bartolomeo [149]. These developments are maturing key technology for space (see Sect. 6) making it possible not only to deploy compact atomic clocks for global positioning and navigation, but also high-performance atomic clocks for fundamental physics and geodesy (see Sect. 4).

3.4 Recommendation: road-map to space clocks

As discussed above and in Sect. 5, there are many possible configurations of space-clock missions. Here we envisage the possibility of three missions, undertaken in stages:

- Complete ACES and launch it to the ISS with utmost urgency.
- Implement I-SOC Pathfinder (or an equivalent) on the ISS as an ACES follow-on mission. The payload, containing an active H-maser, a microwave link, and a laser-pulse optical link, is designed for comparison of ground-based optical clocks to 10^{-18} fractional frequency precision in 1 day. This would have applications in fundamental physics discovery, proof-of-concept optical timescales, and geodesy.
- Launch a dedicated satellite in a highly elliptical orbit containing a strontium optical lattice clock with a 1×10^{-18} systematic uncertainty and $1 \times 10^{-16}/\sqrt{\tau}$ instability, with a coherent optical link to ground. The goals of such a mission are similar to the ones proposed in FOCOS [122]; strong cooperation between the proponents of these missions would be very beneficial. A mission of this type will enable more precise comparisons across a wider network of ground clocks, and direct searches for new physics, including stringent tests of general relativity.

Within this road-map, we highlight an urgent need to develop coherent free-space optical links capable of clock comparisons at the 10^{-18} level in less than 1 day. Once optical links are in place, a critical element of the road-map is then the qualification of a strontium optical lattice clock for operation in space.[4] To achieve this, a technology development programme must be undertaken for several components of the clock: optical resonators to stabilize lasers to a noise floor of 1×10^{-16} in fractional frequency; laser sources at six different wavelengths to cool, optically confine and interrogate strontium atoms; compact physics packages with a controlled black-body radiation environment; and compact frequency combs. Further details of the required technologies are outlined in Sect. 6.

The possibility of establishing collaborations between Europe, US, and China on major future scientific missions should be investigated.[5] Further, synergies should be exploited between the atomic clock missions discussed here and the requirements for the AEDGE mission [5] discussed in Sect. 5, for which the same cold-strontium technology is envisaged.

4 Quantum gravimetry for Earth observation review

4.1 The observation of mass change and Earth observation requirements

4.1.1 Earth sciences and gravity field observations

The Earth sciences need gravity-field observations from satellites for understanding the Earth's structure and monitoring its changes related to geodynamics and climate change,

[4] Ytterbium could be used in place of strontium, as proposed in FOCOS [122]. The atoms share similar complexity and capability: either would be suitable for an atomic clock mission, or for an atom-interferometer science mission of the type proposed in AEDGE [5].

[5] For example, in the context of the US National Academies Decadal Survey on Biological and Physical Sciences Research in Space 2023–2032 [150].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 15/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

as well as many other needs of society related to its glacio- and hydrosphere. Satellite earth observations (EO) enable the observation and monitoring of the Earth's density distribution and mass variations, and provide a significant contribution to the determination of many Essential Climate Variables (ECVs) as defined by the Global Climate Observing System (GCOS) [151]. ECVs monitor phenomena that are changing the world we live in, such as climate change, changing water resources, flooding, melting of ice masses, global sea level rise and atmospheric changes. Better knowledge of such phenomena is bound to lead to significant societal benefits via, e.g., the operational prediction of floods and droughts, monitoring and prediction of sea level rise, and in applications regarding water management.

For these reasons, several initiatives and studies have been launched in the past years at the international level to foster continued observation and monitoring of mass and mass transport phenomena. In 2015 the International Union of Geodesy and Geophysics (IUGG) issued a resolution on "Future Satellite Gravity and Magnetic Mission Constellations" [152] and launched an international multidisciplinary study on science and user needs for the observation of mass transport to understand global change and to benefit society. In the resulting report it was stated that "… a satellite gravity infrastructure is needed with increased space-time sampling capability, higher accuracy and sustained observations" [153], see also [154]. Moreover, in 2019 the International Association of Geodesy (IAG) started the "Novel Sensors and Quantum Technology for Geodesy" (QuGe) initiative; in its framework, three working groups have been defined, one specifically devoted to "Quantum gravimetry in space and on ground" [155] and a second one targeting "Relativistic geodesy with clocks" [156] (See Sect. 3.1).

4.1.2 Past, present and planned gravimetry missions

In the past twenty years, satellite gravity missions have helped form a well-organized user community tracking the Earth mass movements and to study environmental changes on a global scale using data from satellite observations (see Fig. 4 and Table 1).

The general principle of gravity missions is based on precise tracking of satellite position (in free fall) and inter-satellite ranging, combined with the determination of non-gravitational accelerations by means of accelerometers on board the satellites. The technology exploited so far for gravimetry missions is represented by electrostatic accelerometers (EA), as in the CHAllenging Minisatellite Payload (CHAMP) mission that flew from 2000 to 2010 [44], where the accelerometer provides observations that represent the surface forces acting on the satellite, i.e., all non-gravitational accelerations (drag, solar and Earth radiation pressure), so that the Earth gravity field can be obtained from purely gravitational orbit perturbations (observed by satellite tracking). Based on the same EA technology but measuring orbit perturbations by satellite-to-satellite tracking, the NASA/DLR (Deutsches Zentrum für Luft- und Raumfahrt) GRACE mission that flew from 2002 to 2017 [45] and the GRACE-FO mission launched in 2018 [15] have provided and are providing routine measurements of the spatial variations of the Earth gravity field in space and in time on a monthly basis. These missions are based on the concept of flying a pair of low-Earth-orbiting satellites, precisely tracked using the Global Navigation Satellite System (GNSS) and an inter-satellite microwave ranging system, with the accelerometers enabling the measurement of non-gravitational forces. This allows for long-term monitoring of the gravity field and its time variations.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 16/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 4 The CHAMP [44], GRACE [45] and GOCE [13] satellites and mission concepts

Table 1 Accuracies in the determination of the gravity field by "classical" measurements (including the planned ESA-NASA NGGM/MAGIC mission [159])

Measurement type	Monitoring gravity field time variations			Static gravity field
	CHAMP 2000–2010	GRACE/GRACE-FO 2002—ongoing	NGGM Launch scheduled 2028	GOCE 2009–2013
EA accuracy Geoid undulations	$\sim 10^{-10}$ m/s^2 $\sim$ 10 cm @350*km*	$\sim 10^{-11}$ m/s^2 $\sim$ 10 cm @175*km*	$\sim 10^{-11}$ m/s^2 $\sim$ 1 mm @ 500 km every 3 days $\sim$ 1 mm @ 150 km every 10 days	$\sim 10^{-12}$ m/s^2 $\sim$ 1 cm @100*km*
Gravity anomalies	$\sim$ 0.02 mGal @1000*km*	$\sim$ 1 mGal @175*km*		$\sim$ 1 mGal @100*km*

However, the 20-year-long time series of observations needs to be prolonged: this is why several space agencies are working on follow-up missions. ESA plans a Next-Generation Gravity Mission—NGGM [157], scheduled for launch in 2028 and also known as MAGIC (Mass-change and Geosciences International Constellation), in cooperation with the corresponding NASA MCDO (Mass Change Designated Observable) initiative [158]. The MAGIC mission configuration will be based on two pairs of twin satellites: one pair will fly on a quasi-polar orbit, the other one on an orbit with inclination of about 67° [159], with laser ranging between each pair. This will allow for a reduction in the revisit time, providing higher temporal and spatial resolution.

Based on a different measurement concept, gravimetry can also be performed by acquiring observations from a gradiometer, measuring gravity gradients inside the satellite. This was done in the very successful GOCE mission flying from 2009 to 2013 [13], which exploited gradiometry for a unique mapping of the static gravity field, providing models with unprecedented accuracy for a range of geophysical and oceanographic applications (e.g., sea-level currents, a reference system for global height systems, and background data for geophysics and understanding the Earth interior).

Table 1 summarizes the status of the gravity missions based on classical technology, the accuracies attained so far and the prospects for the near future.

Although the classical gravity field missions have been highly successful, they have not satisfied all the user needs of science and society. These have been summarized in various international reports, in the form of tables such as those provided in [160], see Table 2.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 17/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

Table 2 Consolidated science and users' requirements for earth observation, as reported in [160]

Spatial resolution	Equivalent water height		Geoid	
	Monthly field	Long-term trend	Monthly field	Long-term trend
Threshold requirements				
400 km	5 mm	0.5 mm/yr	50 μm	5 μm/yr
200 km	10 cm	1 cm/yr	0.5 mm	0.05 mm/yr
150 km	50 cm	5 cm/yr	1 mm	0.1 mm/yr
100 km	5 m	0.5 m/yr	10 mm	1 mm/yr
Target objectives				
400 km	0.5 mm	0.05 mm/yr	5 μm	0.5 μm/yr
200 km	1 cm	0.1 cm/yr	0.05 mm	5 μm/yr
150 km	5 cm	0.5 cm/yr	0.1 mm	0.01 mm/yr
100 km	0.5 m	0.05 m/yr	1 mm	0.1 mm/yr

4.1.3 Earth observation requirements

The numbers in Table 2 point towards mission requirements that deliver measurements with higher sensitivity, greater accuracy, and more long-term stability. In summary, the needs are:

- Higher spatial resolution (implying lower orbits, 300–350 km) for detection of gravity changes due to movements of mass in the Earth system. However, it must be remarked that no space mission will be able to map the higher-frequency details of the gravity field, due to the atmospheric limitations of the orbit height. Therefore, a full detailed mapping of the spatial gravity field variations down to a few km resolution must be supplemented by airborne and ground gravity measurements;
- Shorter revisit times, which require flying a satellite constellation, e.g., double pairs such as in NGGM/MAGIC. The improved temporal resolution would be crucial for operational service applications such as near real-time flood tracking. A shorter revisit time will also aid in better determination of tidal effects, which must currently be modelled, and represent a limiting factor in the accuracy of current missions;
- Greater accuracy in the measurements, by exploiting new technologies such as laser interferometry and cold atom accelerometers, with accelerometers accurate to better than $\sim 10^{-10} - 10^{-11}$ m/s^2 (measurement range of $\pm 10^{-4}$ m/s^2) and gradiometers accurate to $\sim 10^{-12}$ m^2/s^2/$\sqrt{\text{Hz}}$ for a GOCE-like mission, over a larger spectral measurement band, due to the spectral behaviour of the quantum accelerometers at low frequencies compared with the spectral behaviour of electrostatic accelerometers, as illustrated in Fig. 5, which is taken from [161]. These improvements in instrument performance and low-frequency stability will become important for satellite constellations;
- Extension of the observation time series, which is essential for long-term monitoring of mass transport and variations, and especially for understanding the separation of natural and anthropogenic forcing.

4.2 Quantum sensors in the context of Earth observation

4.2.1 Classical and quantum sensors for gravimetry

Quantum sensors based on atom interferometry are extraordinarily sensitive to external forces [163], reaching an accuracy of 40 nm/s^2 for gravimeters [164], 8×10^{-8} 1/s^2 for gradiometers [165, 166] and 70 nrad/s for gyroscopes [167, 168]. In Table 3 some of the most relevant features [162, 169] of quantum accelerometers and the classical electrostatic accelerometers used so far for space accelerometry are compared. On the ground,

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 18/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 5 Comparison between the noise spectra in GOCE gradiometers and in a prospective cold atom gradiometer, illustrating the latter's potentially reduced level at low frequencies. Figure taken from [161], see [162] for a recent discussion

Table 3 Comparison of classical and quantum sensors

	Atomic accelerometer	Electrostatic accelerometer
Sensitivity	4×10^{-8} m/s^2/Hz$^{1/2}$ on ground (projection for space at 10^{-12} m/s^2/Hz$^{1/2}$ for interrogations of more than 20 s)	3×10^{-12} m/s^2/Hz$^{1/2}$ (demonstrated)
Measurement bandwidth	≤ 0.1 Hz	[0.005–0.1] Hz
Scale factor	Absolute	Calibration required
Stability	No drift	Drift
Measurement capability	Single axis	Three axes
Proof mass motion	Residual velocities → Coriolis acceleration	
SWaP	High	Low
TRL	Intermediate	High

the best quantum accelerometers are operating at sensitivities of about $5 \cdot 10^{-8}$ m/s^2/Hz$^{1/2}$ [164] for an interrogation time of about 100–200 ms. A space quantum accelerometer is expected to reach sensitivities in the low 10^{-12} m/s^2/Hz$^{1/2}$ when stretching the interrogation times to 20 s, similar to the very best electrostatic accelerometers, such as used for the GOCE [13] mission, but in a wider measurement band extending down to lower frequencies as illustrated in Fig. 5.[6] Absence of drifts is a consequence of the absolute character of quantum sensors, with stable scale factors determined by the wavelength of the laser beam-splitters and the duration of the measurement, and the possibility of evaluating accurately systematic effects. On the other hand, they are so far limited to single axis measurements, and have a much higher Size, Weight and Power (SWaP) budget. However, whilst the technology is currently less mature, it is being demonstrated in a number of national and international projects as outlined in Sect. 6.

[6] For the capabilities of hybrid electrostatic-quantum interferometers, see [170–172].

UNCORRECTED PROOF

AUTHOR'S PROOF



Figure 6 Spectra of gravity field recovery in equivalent water height obtainable with an atom interferometer and an electrostatic accelerometer, shown as black and red lines, respectively, in the left panel and colour-coded in the lower and upper maps in the right panels. Figure taken from [174]

4.2.2 Potential gain in Earth observation by quantum gravimeters

The promise of atomic accelerometers for providing better long-term stability, i.e., smaller measurement noise at the lowest frequencies, below 10 mHz, will enable the reconstruction of the Earth gravity field to be improved [162, 169, 173] over many spherical harmonic degrees, as seen in Fig. 6 [174]. This shows the uncertainties in the gravity-field recovery as a function of spherical harmonic degree, evaluated in equivalent water height, considering quantum (black) or drifting (red) accelerometers in the left panel, and colour-coded in the right panels. The computations were carried out without any empirical periodic parameter adjustment in the gravity-field reconstruction.

The potential gains in Earth observation obtainable using quantum sensors are illustrated in Fig. 7. The ellipses represent the required measurement resolutions for the indicated scientific objectives, with the spatial resolution on the horizontal axis and the temporal resolution on the vertical axis. Also shown are the sensitivity curves of the "classical" CHAMP [44], GRACE [45] and GOCE [13] missions and the prospective sensitivity of a possible quantum gravimetry mission employing atom interferometry.

4.2.3 Potential gain in Earth observation by chronometric levelling

As already mentioned in Sect. 3.1, a combination of ground and space clocks together with high performance time and frequency dissemination capabilities, in particular via satellites can benefit the stabilization and long-term validation of physical height networks. Though this approach is today less developed than quantum gravimeters, it is highly appealing [46, 47] because clocks offer access to a new observable in Earth observation, namely to gravity potential differences, in addition to the established determination of its derivatives. Time and frequency dissemination via satellite supported by space clocks can thus be a valuable ingredient [175] in the establishment of an improved height reference system as required by the United Nations initiative on the global geodetic reference frame for sustainable development [43].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 20/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 7 The spatial and temporal resolutions required for the indicated scientific objectives, compared with the sensitivity curves of the "classical" CHAMP [44], GRACE [45] and GOCE [13] missions and the prospective sensitivity of a possible quantum gravimetry mission. Figure adapted from [178]

4.3 Quantum space gravimetry pathfinder mission

4.3.1 Concepts for a quantum gravimetry pathfinder mission

For the European Union, deploying a quantum space gravimetry (QSG) pathfinder mission within this decade is a strategic priority to ensure technological non-dependence and leadership in this field and to pave the way towards an EU/ESA QSG mission within the next decade [176].[7] In this regard, the pathfinder will represent a fundamental technological step towards the feasibility of such a mission. As such, the pathfinder mission will be important in showing the fitness of cold atom interferometry for the purpose of gravity sensing, even if this first mission will not provide observations allowing for an improvement in the gravity field recovery. We note that the challenges inherent to the launch and operation of a dedicated quantum pathfinder mission have at an early stage led ESA to consider the possibility of embarking the pathfinder mission on one of the MAGIC satellite pairs, to be launched around 2028–30. This mission is at the moment at the stage of a phase A study, but a pathfinder add-on is currently deemed unrealistic from a risk/payload point of view.

[7] To this end, the European Commission has issued a call to enhance the Technical Readiness Levels (TRLs) of components for quantum gravimeters in space, e.g., components for cold atom interferometry (including Bose–Einstein Condensates) [177].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 21/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

4.3.2 Profile for a quantum gravimetry pathfinder mission

The main goal of the pathfinder mission should be to demonstrate the maturity of the cold atom technology to operate in space. It should also go beyond the present-day performance of ground atom accelerometers (few 10^{-12} m/s$^2/\sqrt{\text{Hz}}$) by one to two orders of magnitude thanks to the long interrogation times (several seconds) in microgravity. This interrogation time is limited by the expansion of the atomic ensembles, which challenges the detection and results in a number of systematic effects such as wave-front aberrations and the Coriolis effect [162, 179]. The pathfinder mission will also help to demonstrate the technical maturity of key components of cold atom sensors in space, such as the long operation times or the rotation compensation (see Sect. 6 for further details). Furthermore, the pathfinder mission will have a strategic importance also for geodesists who are looking forward to analyse observations and obtain meaningful geodetic results from the next (fully-fledged) quantum gravimetry mission. The pathfinder will in any case provide interesting observations and results useful for the recovery of the gravity field, even though a clear improvement will be available to end-users in geodesy and geophysics only from the following quantum gravimetry mission. In fact, this pathfinder mission will allow preparing for missions with larger interrogation times (more than 10 s) at resolutions of $\sim 10^{-11} - 10^{-12}$ m/s$^2/\sqrt{\text{Hz}}$ suitable for the wide user community.

The payload is expected to have a mass of a few hundred kg and would require a few hundred W to operate. In order for the quantum sensor to perform optimally, the platform needs to be designed to meet clear constraints (centre of gravity, rotation compensation, etc.). To operate with optimal performance, the orbit, altitude, flight modes and position within the platform should be chosen to ensure a successful operation of the quantum sensor.

4.4 Recommendation: road-map to quantum Earth observation in space

As has been outlined in Sect. 4.3, several possible scenarios can be envisaged for future Earth Observation missions based on quantum technology. Here we summarize the main possibilities, their drawbacks and strengths::

- In parallel with the conventional planned MAGIC gravimetry mission [159], it is necessary to update the quantum instrument specifications and requirements. However, embarking the quantum sensor as a passenger on an SST geodesy mission poses tremendous technical challenges and carries significant technological and programmatic risks for both aspects of the mission (classical and quantum sensors).
- Hence the prevailing outcome of the discussions among scientific experts during the workshop is a recommendation to launch a pathfinder mission within this decade with a performance of up to 10^{-10} m/s$^2/\sqrt{\text{Hz}}$ on a dedicated platform. Such a mission would balance the need to have a test of the quantum technology in space and the level of expectation from the quantum gravimetry pathfinder mission. It would also be a clear milestone for other communities, such as the fundamental physics one.
- The success of MAGIC and the Pathfinder mission will then enable the implementation of a full-fledged quantum space gravimetry mission to be launched to follow MAGIC. The definition of the mission scenario and the instrument baseline will be based on lessons learned from MAGIC and the Pathfinder mission.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 22/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

5 Atomic sensors for fundamental science review

5.1 Scientific opportunities

The promise of Cold Atom technologies for making precise experimental probes of topics in fundamental science such as general relativity, cosmology, quantum mechanics and the search for new physics beyond the Standard Model has been recognised in many terrestrial and space projects. To cite just a few examples: in the US the Matter-wave Atomic Gradiometer Interferometric Sensor (MAGIS) 100 m atom interferometer is under construction at Fermilab [180] and NASA has operated the Cold Atom Lab (CAL) Bose–Einstein condensate (BEC) experiment successfully for several years on the International Space Station (ISS) [181, 182]; in Europe the European Laboratory for Gravitation and Atom-interferometric Research (ELGAR) project [183] has been proposed; initial funding has been provided for a suite of experiments applying Quantum Technology for fundamental physics in the UK, including the terrestrial Atom Interferometer Observatory and Network (AION) experiment [184];[8] in France the Matter wave-laser based Interferometer Gravitation Antenna (MIGA) experiment is under construction [186]; in Germany there is the Very Long Baseline Atom Interferometry (VLBAI) programme [187] and a series of BEC experiments in microgravity using MAIUS sounding rockets [188, 189]; the Space-Time Explorer and Quantum Equivalence Principle Space Test (STE-QUEST) experiment has been proposed [190, 191], following the success of the MICROSatellite pour l'Observation du Principe d'Equivalence (MICROSCOPE) experiment [192, 193] that made a pioneering test of the Equivalence Principle in space; and in China the terrestrial Zhaoshan long-baseline Atom Interferometer Gravitation Antenna (ZAIGA) atom interferometer [194] is under construction.[9]

The deployment of cold atom technologies in space offers unique research opportunities in the fields of fundamental physics, cosmology and astrophysics, as represented in several White Papers submitted to the ESA Voyage 2050 call for mission concepts [4, 6–11]. We focus in the following on two of these mission concepts.

One is based on the previous STE-QUEST proposal [191], and proposes a double atom interferometer with rubidium and potassium "test masses" in quantum superposition to test the universality of free fall (UFF). It assumes a single satellite in a 700 km circular orbit, and applies recent developments on gravity gradient control by offsetting laser frequencies, which enables the atom positioning requirements to be relaxed by a factor > 100 [6, 196]. This offers the possibility of probing the UFF, i.e., the equivalence principle, with an unparalleled precision $\mathcal{O}(10^{-17})$ after 18 months in orbit [196], see Fig. 8.

The other is the Atomic Experiment for Dark matter and Gravity Exploration (AEDGE) concept for a satellite atom interferometer using strontium [5], illustrated in Fig. 9, which is similar in principle to the previous Space Atomic Gravity Explorer (SAGE) concept for a satellite atom interferometer experiment [199]. In the baseline AEDGE configuration [5] the atom clouds were assumed to be located inside the spacecraft and have sizes ~ 1 m. Future concepts with atomic ensembles outside the satellite could allow much higher number of momentum transfers and increase sensitivity if they turn out to be possible, a concept called AEDGE+.[10] AEDGE can search for waves of ultralight dark matter (ULDM) parti-

[8] For a recent review of atom interferometer detectors for gravitational waves, see [185].

[9] We also note that quantum correlation in the form of optical entanglement has been verified by the Micius satellite experiment [195] over distances exceeding a thousand km.g

[10] Technical issues for such a configuration are discussed in [200]. See also the Workshop talk by Nan Yu [201].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 23/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 8 Averaging of systematic uncertainties due to gravity gradients in a UFF test with Rb and K quantum sensors. Gravity Gradient Compensation (GGC) significantly reduces the systematic contributions [197, 198], such that the residual differential acceleration may be attenuated to an unprecedented degree through signal demodulation (orange curve). This not only allows for requirements on the source preparation that are greatly reduced compared to other mission proposals as STE-QUEST [191], but also paves the way for more ambitious mission scenarios targeting $\delta\eta \leq 10^{-17}$ in shot-noise limited operation (red curve). In contrast, even though the systematics are integrated down thanks to demodulation, the measurement would be limited by systematics without GGC (blue curve). The figure is taken from [196]



Figure 9 Possible scheme for a satellite atom interferometer experiment [5]. It has two master laser beams M1 and M2 (dotted and solid lines) and two reference beams (R1 and R2) there are two local oscillator lasers LO1 and LO2 (dashed lines) phase-locked with R2 and R1, respectively. Photodetectors (PD1 and PD2) measure the heterodyne beatnote between the reference beams R2 and R1 and the corresponding local lasers LO1 and LO2, respectively, providing feedback for the laser link. Non-polarizing beam splitters are denoted by BS, and tip-tilt mirrors used for controlling the directions of the laser beams are denoted by TTM. Small offsets between overlapping laser beams have been introduced for clarity. The figure is taken from [210]

cles with masses between $\mathcal{O}(10^{-21})$ and $\mathcal{O}(10^{-15})$ eV and measure gravitational waves in a frequency range between $\mathcal{O}(1)$ and $\mathcal{O}(10^{-2})$ Hz [5], intermediate between the ranges of frequency where the sensitivities of the terrestrial laser interferometers Laser Interferometer Gravitational-Wave Observatory (LIGO) [202], Virgo [203], Kamioka Gravitational Wave Detector (KAGRA) [204], Einstein Telescope (ET) [205] and Cosmic Explorer (CE) [206] are maximal ($\mathcal{O}(1)$ to $\mathcal{O}(10^2)$ Hz) and the optimal frequencies of the Laser Interferometer Space Antenna (LISA) [207], TianQin [208] and Taiji [209] ($\mathcal{O}(10^{-2})$ Hz).

In the following we present some representative examples of the capabilities of these cold atom concepts for probing fundamental physics, cosmology and astrophysics.

5.1.1 Tests of the universality of free fall

The Einstein Equivalence Principle (EEP) is the foundation of all theories of gravitation that describe it as a geometrical phenomenon, i.e., a curvature of space-time. Indeed, the universal coupling to all mass-energy that is implicit in the EEP is necessary for all metric

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 24/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

theories of gravitation, including general relativity among many others. As such, the EEP is one of the most foundational building blocks of modern physics. Nonetheless, many modified theories of gravity that go beyond the Standard Model and general relativity and/or account for dark matter/energy entail some violation of the EEP. Examples include scalar-tensor theories of gravity with screening mechanisms, such as chameleon and symmetron models, in which additional scalar degrees of freedom can give rise to long-range fifth forces that cause deviations from general relativity that depend on the density of extended objects (and not just their mass), as well as the local environment and the configuration of the experimental test.[11] Atom interferometry experiments already provide stringent constraints on certain screened fifth-force models (see, e.g., [212–216]), and further searches for fifth forces have been identified as a priority (see, e.g., [6] and Sect. 5.1.2 for more examples).

The best known aspect of EEP is the universality of free fall (UFF), i.e., the weak equivalence principle (WEP).[12] The history of experimental tests of UFF/WEP goes back as far as the Renaissance, and probably beyond. A simple phenomenological figure of merit for all UFF/WEP tests is the Eötvös ratio η_{AB} for two test objects A and B and a specified source mass of the gravitational field:

$$\eta_{AB} = 2\frac{a_A - a_B}{a_A + a_B}, \tag{5.1}$$

where a_i ($i = A, B$) is the gravitational acceleration of the object i with respect to the source mass. Note that for a given experiment the data can be interpreted with respect to different source masses (see, e.g., Ref. [218]) with correspondingly different results for η_{AB}.

Whilst η_{AB} is a useful tool for comparing different experiments, it cannot account for the diversity of possible underlying theories, e.g., different types of couplings depending on the source and test objects, or couplings to space-time varying background fields other than local gravity, e.g., [219, 220]. Thus, not only best performance in terms of the Eötvös ratio is required, but also a large diversity of test objects and source masses.

Table 4 presents the state of the art in UFF/WEP tests, separated into different classes as a function of the type of test-masses employed. In particular, we distinguish between tests using macroscopic test masses and atom-interferometry (AI) tests[13] that use matter waves in a quantum superposition, possibly condensed to quantum degenerate states (Bose–Einstein Condensates) with coherence lengths $\geq \mu$m. The "game-changing" results of the MICROSCOPE mission demonstrate the potential of going into a quiet and well-controlled space environment, with potentially "infinite" free fall times, as exemplified by the prospective sensitivity of the STE-QUEST-like space-borne cold atom interferometer mission displayed in Table 4.

5.1.2 Ultralight dark matter detection

Figure 10 shows examples of the present and prospective sensitivities of searches for linear couplings of ultralight scalar dark matter to photons (left panel) and electrons (right

[11] See [211] for a recent review of screening mechanisms and experimental tests.

[12] The EEP corresponds to the combination of the UFF/WEP with local position invariance and Lorentz invariance. See [217] for a test of Lorentz invariance using MICROSCOPE data.

[13] See also [221].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 25/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

Table 4 Compilation of UFF/WEP tests. Numbers in brackets are results expected in the near future, and we also show the performance of an STE-QUEST-like atom-interferometry mission in the context of this road-map. Further information about this road-map milestone are provided in Sect. 8. In addition, lunar laser ranging probes the UFF at a level $\sim 2 \times 10^{-14}$ for the Earth and Moon [222]

Class	Elements	η	Year [ref]	Comments
Classical	Be–Ti	2×10^{-13}	2008 [218]	Torsion balance
	Pt–Ti	1×10^{-14}	2017 [192]	MICROSCOPE first results
	Pt–Ti	(10^{-15})	2019+	MICROSCOPE full data
Hybrid	^{133}Cs–CC	7×10^{-9}	2001 [223]	Atom Interferometry
	^{87}Rb–CC	7×10^{-9}	2010 [224]	and macroscopic corner cube
Quantum	^{39}K–^{87}Rb	5×10^{-7}	2014 [225]	different elements
	^{87}Sr–^{88}Sr	2×10^{-7}	2014 [226]	same element, fermion vs. boson
	^{85}Rb–^{87}Rb	3×10^{-8}	2015 [227]	same element, different isotopes
	^{85}Rb–^{87}Rb	3.8×10^{-12}	2020 [228]	$\geq$ 10 m towers
	^{85}Rb–^{87}Rb	(10^{-13})	2020+ [198]	
	^{170}Yb–^{87}Rb	(10^{-13})	2020+ [229]	
	^{41}K–^{87}Rb	10^{-17}	2035+	STE-QUEST-like mission
Antimatter	$\overline{\text{H}}$–H	(10^{-2})	2020+ [230]	under construction at CERN

panel).[14] Such dark matter, because of its non-universal coupling to the fields of the standard model, is also one example of a violation of the Einstein equivalence principle and the universality of free fall, UFF. The shaded regions are excluded by current experiments including MICROSCOPE [193] and atomic clocks [86].[15] We also show the prospective sensitivities of rubidium-based terrestrial interferometers (MIGA [186] and ELGAR [183]) and strontium-based terrestrial and space-borne atom interferometers (AION [184] and AEDGE [5], respectively). MIGA, ELGAR and the 100 m and km versions of AION offer significantly greater sensitivity than current experiments (torsion balances, atomic clocks and MICROSCOPE [193]) to couplings of scalar ULDM with masses $\lesssim 10^{-12}$ eV to both photons and electrons. A sensitivity of $\sim 10^{-7}$ to the ULDM-photon coupling in this mass range could be obtained with a prospective cold-atom interferometer UFF probe with a precision of 10^{-17} as discussed in the context of this road-map [6], using Rb and K quantum probes (see also Table 4). As seen in Fig. 10, the AEDGE strontium-based space-borne atom interferometer concept [5] could offer the greatest sensitivity to both the ULDM-photon and -electron couplings for masses between 10^{-15} and 10^{-21} eV, with a maximum sensitivity $\sim 10^{-14}$ for masses between 10^{-17} and 10^{-18} eV.

5.1.3 Probes of general relativity

The measurements of gravitational waves using an atom interferometer in space offer unique prospects for probing modifications of general relativity. For example, AEDGE measurements of the inspiral of merging black holes with a combined mass of 10^4 solar masses at a redshift $z \sim 1$ would be sensitive to the possible appearance of a graviton mass $\lesssim 10^{-26}$ eV, over three orders of magnitude below the current upper limit from LIGO and Virgo [236]. These measurements could also be used to search for possible violations

[14] One may also consider quadratic couplings, which would be screened in terrestrial experiments if the quadratic couplings are positive, reducing the experimental sensitivity [220] (see also [231]). A space-based experiment such as AEDGE is less affected by this screening, and hence maintains sensitivity at larger masses, as discussed in [5, 184]. See [232, 233] for other suggestions how atom interferometers, gravimeters and accelerometers could be used to search for alternative dark matter candidates.

[15] See also the results from the Weekend Relaxion-Search Laboratory (WReSL) experiment on the couplings of ultralight scalar dark matter to photons and electrons oscillating at higher frequencies [234], and recent experimental bounds on the couplings to the up, down, and strange quarks and to gluons obtained using molecular spectroscopy [235].

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 26/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 10 Sensitivities to the possible couplings of scalar ULDM to photons (left panel) and to electrons (right panel) of the terrestrial AION [184], MIGA [186] and ELGAR [183] experiments, and of the AEDGE space-borne concept [5]. Also shown is a combination of the current constraints from MICROSCOPE [193] and terrestrial torsion balance and atomic clock experiments and (in the left panel) the sensitivity of a prospective measurement of the universality of free fall (UFF) with a precision of 10^{-17} [6]

of Lorentz invariance in the propagation of gravitational waves, with a sensitivity complementary to the searches by LIGO/Virgo and for gravitational Čerenkov radiation [236].

More classical GR tests using modern optical clocks are measurements of the gravitational Shapiro delay down to 10^{-8} [237] or tests of the gravitational red-shift at 10^{-9} as in the Fundamental physics with an Optical Clock Orbiting in Space (FOCOS) mission [122], proposed recently in the context of the NASA decadal survey. In both cases, these would provide 3–4 orders of magnitude improvements on best current knowledge.

There is also a proposal to probe models of dark energy by deploying a smart constellation of four satellites in an elliptic orbit around the Sun and making orientation-independent measurements of the differential accelerations between each pair of satellites, using as test masses atomic clouds far away from the spacecraft in open-space vacuum (see the Workshop presentation by Nan Yu [201]). It has also been suggested to deploy an atomic clock at a distance $\mathcal{O}(150)$ AU to probe the low-acceleration frontier of gravity and the local distribution of dark matter [9]. Another suggestion is to detect the gravito-magnetic field of the galactic dark halo by locating atomic clocks at Sun-Earth Lagrange points and measuring the time-of-flight asymmetries between electromagnetic signals travelling in opposite directions, which would be generated partly by the angular momentum of the Sun and partly by the angular momentum of the dark halo [8].

5.1.4 Quantum mechanics

It has been proposed to test quantum correlations over astronomical distances [7, 10], e.g., between the Earth and the Moon or Mars, or between LISA spacecraft.[16] The Micius measurements [195] already demonstrate that quantum correlations extend over 1200 km and that the apparent effective correlation speed exceeds 10^7 c, and Earth-Moon experiments could improve these sensitivities by factors $\sim 2 \times 10^4$ [10].

[16] See also [238] for a description of the proposed Deep Space Quantum Link (DSQL) mission, whose goals include long-range teleportation, tests of gravitational coupling to quantum states, and advanced tests of quantum nonlocality.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 27/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

It has also been proposed to test wavefunction collapse and models predicting the violation of the quantum superposition principle [239, 240] by monitoring the expansion of a cloud of cold atoms [241]. Current results already impose relevant constraints on the correlation length and rate parameters of the Continuous Spontaneous Localisation (CSL) model: see [241] for a detailed discussion.

In addition, atom interferometry can be exploited to perform nontrivial tests of the EEP in the quantum regime [242]. These include UFF tests with atoms in quantum superpositions of internal states [243] and with entangled atomic species [244], as well as gravitational redshift measurements with quantum clocks in a superposition of different heights [245].

As well as testing quantum mechanics and deploying it in applications and sensing opportunitites, cold atom ensembles can be investigated in their own rights. A freely falling experimental setup removes any need for measures to compensate for gravitational effects that could distort the overall trap potential. Consequently, reduced velocities and densities of the condensed atom ensemble can be achieved and near-monochromatic matter waves established. This enables more detailed studies of various phenomena in atom optics, such as Feshbach resonances [246], matter wave reflections [247] and bubble formation [248], improving on ground-based results. Other research areas include the study of quantum gas mixtures which, in ground based experiments, are separated due to differences in the gravitational sag and applied magnetic field. This allows more detailed studies of superfluidiy, vortex formation, and quantum droplets [249].

Carrying out these experiments requires cold atom laboratories in microgravity, such as Cold Atom Laboratory (CAL) and Bose Einstein Condensate and Cold Atom Laboratory (BECCAL). In view of the resources required for such space-based research, a next-generation quantum laboratory on a microgravity platform should enable a broad participation of researchers and their ideas.

5.1.5 Cosmology

AEDGE measurements also offer new opportunities in cosmology, such as unparalleled sensitivity to possible emissions from collapsing loops of cosmic strings in a network with tension $G\mu \gtrsim \mathcal{O}(10^{-18})$, more than 3 orders of magnitude below the current limit from the third Advanced LIGO–Virgo observing run [250] and an order of magnitude beyond the reach of LISA, as seen in the left panel of Fig. 11 [5]. Such measurements could also be sensitive to effects in the early Universe that cause it to deviate from the conventional expectation of adiabatic expansion, such as a period of matter dominance or kination, as seen in the right panel of Fig. 11 [251].[17] There are also some models that yield primordial gravitational waves that could be detected by AEDGE, see, e.g., Fig. 12 in [258].

Another cosmological opportunity is the search for a stochastic background of gravitational waves generated by a first-order phase transition in the early Universe, either in the electroweak sector of the Standard Model or in some extension that includes a new interaction generated by a massive boson beyond the reach of present and proposed collider experiments. The left panel of Fig. 12 compares the sensitivities in the (T_*, α) plane (where T_* denotes the critical temperature and α the strength of the transition) of the indicated

[17] The string tension $G\mu = 10^{-11}$ used in this plot is the maximum consistent [252, 253] with current data from pulsar timing arrays [254–257], which report strong evidence for a spectrally-similar low-frequency stochastic process.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 28/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 11 Left panel: Sensitivities to the GWs from a network of cosmic strings with tension $G\mu$ of AION-100 and -km, AEDGE [5] and AEDGE+ (a version of AEDGE that would use an ∼ 100m atomic cloud outside the spacecraft [251]), LIGO, ET and LISA. Right panel: Possible effects on the spectrum of GWs from cosmic strings with tension $G\mu = 10^{-11}$ of an epoch of matter domination (MD) or of kination [251]



Figure 12 Left panel: Sensitivities of AION-100 and -km, AEDGE, AEDGE+ and LISA in the (T_*, α) plane to GWs from generic first-order transitions with a transition rate $\beta/H = 100$ [259]. Right panel: The expected signal-to-noise ratio (SNR) with AEDGE for observing the stochastic GW background from a $U(1)_{B-L}$-breaking phase transition. In the red regions SNR > 1000 and the solid gray contour corresponds to SNR = 10 [259]

experiments to GWs from a generic phase transition with a transition rate $\beta/H = 10^2$, and the right panel shows the signal-to-noise ratio (SNR) in the $(m_{Z'}, g_{B-L})$ plane expected from AEDGE measurements of the stochastic GW background from a $U(1)_{B-L}$-breaking phase transition [259]. In the red regions SNR > 1000 and the solid gray contour corresponds to SNR = 10 [259].

5.1.6 Astrophysics

Opportunities in astrophysics are also opened up by these gravitational-wave measurements, including the observation of mergers of intermediate-mass black holes (IMBHs) that could reveal how the super-massive black holes at the centres of many galaxies were assembled [5], as illustrated in Fig. 13.[18] As also seen in Fig. 13, there are good prospects of synergies obtained by networking detectors working in different frequency ranges, e.g., LISA might measure the initial inspiral stage of IMBH mergers whose final stages would

[18] We note also that timing measurements using atomic clocks located on asteroids have been proposed as a way to measure gravitational waves in the μHz range [260], in the gap between LISA and pulsar timing arrays.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 29/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 13 Strain sensitivities of AION-10, -100 and -km, AEDGE and AEDGE+, compared with those of LIGO, LISA and ET and the signals expected from mergers of equal-mass binaries whose masses are 60, 10^4 and 10^7 solar masses. The assumed redshifts are $z = 0.1, 1$ and 10, as indicated. Also shown are the remaining times during inspiral before the final mergers [5, 251]



Figure 14 Left panel: The strain sensitivities of AEDGE and other experiments compared with the neutrino gravitational memory signal expected from a core-collapse supernova calculated using model Ac3G in [261]. Right panel: Sensitivities to the amplitude A of a delta function peak in the curvature power spectrum at scale k_* of AION-km, AEDGE and AEDGE+, LIGO, ET and LISA. The gray region is excluded by constrains on primordial black holes. The top axis indicates the horizon mass that corresponds approximately to the mass of the formed primordial black holes [251]

be measured by AEDGE, and AEDGE measurements of the initial inspiral stages of mergers of lower-mass black holes could be used to predict the direction and timing of their final stages, providing advance warning for multimessenger observations. Also, as shown in the left panel of Fig. 14, AEDGE could observe [251] the gravitational memory effect due to neutrinos emitted from a collapsing supernova in our galaxy [261] and, as shown in the right panel of Fig. 14, AEDGE would also be uniquely sensitive to possible features in the spectrum of cosmological density fluctuations that could lead to a population of primordial black holes with a density orders of magnitude below the current astrophysical limits [251].[19]

[19] We comment in passing that radio astronomy would gain greatly from extending the Event Horizon Telescope [262] concept to a space mission with a much longer baseline, which would benefit from using atomic clocks for synchronization.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 30/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

5.2 Connection to technology development section

5.2.1 Optical clocks

As outlined in Sect. 3.2, several current activities exist to advance the capabilities for optical clocks in space. Depending on the type, environment, and targeted timescale of these clocks, the achievable stability varies. Similar to the capabilities, the requirements for the above discussed experiments differ. LISA, similar to navigational applications and earth observation, require high short-term stability [207, 263], with the stability of the reference directly impacting the quality of the intended measurement. On the other hand, space-borne tests of special relativity, as proposed in [264] for instance, or made using the signals of misaligned Galileo satellites [19], require a high stability at orbit time.

As described in Sect. 3.2.1, the stabilization concept has to be chosen for a specific purpose. Optical frequency references based on *optical resonators* usually show the best performance on short time scales, where fluctuations in the resonator length can be dominated by thermal noise and external perturbations can be sufficiently suppressed [265]. If integrating over longer times, drifts of the resonator length caused by thermal or mechanical stress or material aging impact the achievable stability. To increase the long-term stability, efforts have been undertaken to reduce the impact of outside effects, such as by the choice of spacer material [265] and length of spacer, enclosure in thermal shields [114], or cryogenic environments [266]. The latter is especially interesting in ground-based experimental systems, such as for instance in [267].

Absolute frequency references based on the spectroscopy of atoms or molecules, on the other hand, benefit from prolonged integration times. In general, the width of the probed line determines the achievable stability. For such references, usually a trade-off has to be made between the required stability and the available SWaP budget. As described in Sect. 3.3, different concepts have been considered for space operation. We note in particular that space operation, especially outside the International Space Station, requires full automation and high reliability of the system.

As outlined above, fundamental experiments require quiet environments to measure the desired effects. While this does not always necessitate operation in orbit, space-qualified optical frequency references are a key technology to enable the research. Section 3.3 details current and planned missions for operation in orbit, demonstrating operation at a stability below 10^{-13} at orbit time. To improve on the current best measurements performed on the ground, the frequency stability of any space-borne frequency reference needs to be below $1 \cdot 10^{-15}$ at orbit time. Based on the current developments and demonstrated optical references in space, this appears to be feasible for both short-term (in the range of seconds to minutes) and long-term operation (in the range of hours).

5.2.2 Optical links

Optical links are required to fulfill the scientific goals of various planned and executed missions. Those missions can be divided in such examples with inter-satellite links, such as LISA [207] and AEDGE [5], and those with ground-to-satellite links, such as Micius [195]. As outlined in Sect. 3.2.3, currently mainly microwave links have been established, which do not satisfy the requirements for fundamental science missions.

As an obvious example, entangled optical photons, as required for Micius, can only be transferred by optical ground to satellite links. The success of Micius was a key step towards space-based fundamental quantum entanglement experiments. In future quantum

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 31/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

entanglement experiments, such as the quantum link to the Lunar Gateway in the proposed DSQL mission [238], one could envisage using two space-borne platforms to eliminate the atmospheric impact on the optical link.

Bidirectional ground-to-satellite optical links are a necessity for many applications, such as the proposed Kepler constellation [137]. Additionally, they could play a part in comparisons of optical frequency references operating in different gravitational potentials, i.e., on the ground and in space.

Precision in optical inter-satellite links is crucial to the scientific goals of missions employing long-range laser ranging, such as LISA [207], where picometre level changes over the separation of the satellites of the order of $5 \cdot 10^9$ metre will be detectable. In this case, the frequency of the deployed laser enables more precise measurements than achievable with a microwave link.

Finally, gravitational wave detection measurements with cold atom sensors require a pair of atom interferometers on two satellites irradiated by the same laser beams to achieve the necessary coupling. Current proposals foresee a linkage over $4 \cdot 10^7$ metre to reach the strain sensitivity displayed in Fig. 13. The achievable precision described in Sect. 3.2.3, underline the feasibility of missions such as LISA and AEDGE.

Outside of space-based experiments, also ground-based projects, such as ELGAR [183], require coherent long-distance free-space laser beams. Since the necessary length is short compared to the optical links discussed above and the accessibility of the system, the involved technology is currently at hand.

5.2.3 Atom interferometry

In the sections above, laser stabilization and information transfer over optical links has been discussed. The execution of future spaced-based experiments for fundamental science exploitation, such as STE-QUEST [191] and AEDGE [5], requires high-precision atom interferometers, with different interferometer schemes. Whereas, for a test of the weak equivalence principle two different masses (here atomic species) need to be observed in the same place, the detection of gravitational waves and other astrophysical phenomena require elaborate schemes with a single atomic species and optical links between the atom interferometers, interrogated by the same laser.

As described in [191], weak equivalence principle experiments are made using two different masses, whose free-fall behaviours are observed and compared. Improvements on the current experimental sensitivity are possible, in principle, using any pair of different atomic species. The visibility of any deviation from the equivalence principle prediction increases with the interrogation time. As an example, according to current estimates, experiments with ^{85}Rb and ^{87}Rb require interrogation times of 10 s or more in interleaved operation. This enables measurements of differential accelerations better than 10^{-13} m/ s^{-2}, necessary for the targeted precision in the Eötvös parameter, see Table 4. These types of experiments require two species interferometers on a single satellite, which also allows for the increased free-fall time.

The detection of gravitational waves, on the other hand, strongly profits from large distances between two atom interferometers in a gradiometric configuration. Such experiments require optical links over long periods with two connected satellites in Earth orbit. The specific interferometer concept requires an atomic species with a clock transition at optical frequencies, e.g., strontium. The free-fall times of the atoms anticipated for these

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 32/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

experiments are substantially longer than those discussed above. For example, to enable gravitational wave detection as outlined in Sect. 5.1, free-fall times in the order of 600 s are required to enable characteristic strain measurements down to 10^{-23} at about 80 mHz.

Atom interferometry in microgravity has been performed for several years. Experiments have been performed in a drop tower (QUANTUS, PRIMUS) [268, 269], on parabolic flights (Interférométrie Cohérente pour l'Espace, I.C.E.) [270], on sounding rockets (MAIUS) [188], and on the ISS (CAL) [181]. The planned BECCAL mission [271] is the next-generation ultra-cold atom laboratory, including high precision atom interferometry, and is being prepared for deployment on the ISS. Important challenges in miniaturization and automation have been addressed in the development of CAL and BECCAL. We note in addition that pathfinders for the key underlying optical technologies such as FOKUS [272], KALEXUS [273] and JOKARUS [274] have already demonstrated reliable operation. The next step with the proven technology is leaving the International Space Station and integrating an atom interferometer into a satellite. With that development, space-borne tests of the weak equivalence principle, as proposed in [191], would come within reach.

AEDGE represents the following step in advancing quantum technologies for deployment in space. It requires two optically-linked atom interferometers, increasing the distance and thereby sensitivity of gravitational wave detectors with respect to ground-based measurements. The measurement is enabled by employing strontium as opposed to rubidium or potassium. This leads to an additional complexity in the system, such as availability of miniaturized laser systems, electronics, and optics.

Based on the present technology in space-based rubidium and potassium systems [181, 188, 271] and ground-based strontium systems [275, 276], the developments required for the next steps appear feasible.

In this connection, we note that the experimental landscape of atom interferometry projects for fundamental science exploitation has expanded significantly in recent years, with several terrestrial experiments, based on different cold atom technologies, currently under construction, planned or proposed.

As mentioned above, four large-scale prototype projects are funded and currently under construction, i.e., MAGIS [180] in the US, MIGA [186] in France, ZAIGA [194] in China and AION [184] in the UK. These will demonstrate the feasibility of atom interferometry at macroscopic scales, paving the way for terrestrial km-scale experiments as the next steps. There are proposals to build one or several km-scale detectors, including AION-km at the Science and Technology Facilities Council (STFC) Boulby facility in the UK, MAGIA-advanced and ELGAR [183] in Europe, MAGIS-km at the Sanford Underground Research facility (SURF) in the US, and advanced ZAIGA in China. It is foreseen that by about 2035 one or more km-scale detectors will have entered operation. These km-scale experiments would not only be able to explore sensitively ultralight dark matter and the mid-frequency band of gravitational waves, but would also serve as the ultimate technology readiness demonstrators for space-based missions like STE-QUEST [191] and AEDGE [5] that would reach the ultimate sensitivity for exploring the fundamental physics goals outlined in this section.

The perspectives for large-scale atom interferometer projects are very rich today, with a central focus on establishing the readiness of cold atom technology for use in experiments

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 33/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

to explore fundamental science. These terrestrial pathfinders are advancing further the relevant technologies, closing gaps and addressing fundamental physics questions.

However, in order to deploy effectively strontium or ytterbium on any microgravity platform, in addition to these terrestrial developments a space-borne pathfinder mission or technology demonstrator is key. The development of critical components benefits from synergies between all systems. As such, laser modules developed for clock deployment could be used as the basis for interferometric missions. Similar synergies are present for vacuum generation, frequency stabilization, and low noise electronics between the various systems. With a technology demonstrator for a space-bourne optical lattice system, many applied and fundamental missions could be supported and further developments triggered.

5.3 Recommendation: road-map for fundamental physics in space

To summarize Sect. 5.2, which details the requirements to enable the scientific opportunities from Sect. 5.1, the following recommendations for developments can be made:

- Build upon the ongoing large-scale terrestrial atom interferometer projects for fundamental science exploitation, such as MAGIS [180] in the US, MIGA [186] in France, ZAIGA [194] in China and AION [184] in the UK to construct one or more of the proposed km-scale successors, which will enable technology development for space-based missions like STE-QUEST [191] and AEDGE [5]. The ELGAR experiment [183] would also supplement existing and future, possibly satellite-based, scientific measurements by targeting relevant gravitational wave frequencies.
- Perform fundamental tests in ground based microgravity facilities, such as the drop tower, the Einstein Elevator, parabolic flights, and sounding rockets, to develop technology and support scientific findings.
- Prepare a satellite mission as the next step in cold and condensed atom technology in space with a two-species interferometer based on the available rubidium and potassium sources.
- Prepare optical frequency references for operation in space to test General Relativity and probe modified gravity theories, as well as support space missions such as LISA. This is in accordance with the recommendations in Sect. 3.
- Develop components for deployment in space, with a focus on those with synergies for different missions, for instance optical preparation, laser modules, vacuum generation, and magnetic field control.
- Advance the development of optical lattice systems on the ground, in ground-based microgravity platforms, and in space-based pathfinder missions to enable future gravitational wave detection missions. This includes miniaturization of subsystems and proof-of-principle missions or studies for individual components.

6 Technology development, space qualification and pathfinders

6.1 Requirements for cold atoms in space/specifications for space borne quantum sensors

6.1.1 Atomic clock mission

We advocate a single dedicated satellite in a highly elliptical orbit, containing a strontium optical lattice clock with a 1×10^{-18} systematic uncertainty and $1 \times 10^{-16}/\sqrt{\tau}$ instability, and including a coherent optical link for comparisons to ground-based clocks. Such

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 34/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

a mission would enable a wide network of ground-based clocks to be compared to each other with 10^{-18} precision in 1 day, which would have applications in fundamental physics discovery, proof-of-concept optical timescales, and geodesy. Furthermore, the clock in elliptical orbit would enable direct searches for new physics, including stringent tests of general relativity (see Sects. 3 and 5). To support such a mission, improved optical links must be developed for clock comparisons at 1×10^{-18}. Efforts to realize such links are currently underway in the Atomic Clock Ensemble in Space (ACES) and ISS Space Optical Clock (I-SOC) Pathfinder ESA programmes. However, the bulk of the challenge of a space-clock mission will be to develop a space-qualified strontium optical lattice clock. Present strontium clock technology occupies at least a few m^3 and consists of several complex, delicate components (see Sect. 6.3) which must all be brought up through the TRL scale.

6.1.2 Earth observation mission

As outlined in Sect. 4, quantum sensors offer the perspective of enhancing missions for Earth Observation by embarking them either in a gradiometric configuration on a single satellite for a GOCE-like mission concept, or as accelerometers combined with laser links between satellite pairs for a GRACE-like mission concept in an earth orbit with an altitude of few hundred kilometers and nadir pointing mode. Similar to previous missions in Earth Observation, the targeted mission duration is several years. Sensitivities of ^{87}Rb atom interferometers to accelerations or differential accelerations would be in the range of 10^{-10} to 10^{-12} m/ s^{-2} $Hz^{-1/2}$ at low frequencies, complementing current sensor technology. The SWaP budget of a current quantum sensor may include a few hundred kilograms of payload and a few hundred Watts of power, which need to be reduced. Moreover, the payload could be designed to link multiple quantum sensors on a single satellite by interrogating them with the same beam splitter laser for a GOCE-like mission, or they could be implemented as accelerometers for drag correction in a GRACE-like mission.

6.1.3 Fundamental physics

As discussed in Sect. 5.2, key technologies in three areas are required for fundamental physics tests: optical clocks, optical links, and atom interferometers. The requirements for optical frequency references are described in the paragraph above. In case of optical links, the requirements depend on the specific mission, and range from single photon transmission in case of entanglement missions to long distance coherent laser light transmission in case of gravitational wave detection. For atom interferometers, the requirements are naturally more restrictive the higher the desired precision, which has implications for the atomic flux, preparation of the atoms, coherent manipulation, and interrogation times. The test of the universality of free fall outlined in Sect. 5.2 targets a measurement of the Eötvös ratio at the level of $\leq 10^{-17}$, and requires atom interferometers using two different kinds of atomic species simultaneously (e.g., ^{85}Rb and ^{87}Rb or ^{41}K and ^{87}Rb) with interrogation times of over 10 s to reach a projected sensitivity to differential accelerations of 10^{-13} m/ s^{-2} at a cycle time of 10 s in an interleaved operation. For the case of the proposed space-borne gravitational detector, the scheme is based on strontium atoms, requiring an increased atomic flux, large momentum transfers, and interrogation times of 600 s to enable characteristic strain measurements down to 10^{-23} at about 80 mHz. These parameters place requirements, e.g., on the atomic source as well as the dimensions of the central elements, such as the vacuum chamber, affecting the overall design of the sensor head.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 35/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

Fundamental physics experiments require a variety of different orbital scenarios. For example, entanglement experiments require large distances to close additional loopholes and test the validity of quantum mechanics, hence a geostationary or lunar satellite would be of interest. Clock redshift and local position invariance tests benefit from orbits with large variations in the gravitational potential, whereas local Lorentz invariance tests benefit from short orbital times with high orbital velocities and velocity variations. A prime candidate for such a mission is a satellite in low Earth orbit, while nevertheless avoiding vibrations caused by drag in the atmosphere. Orbital heights of 600 km with orbital times in the order of 90 min appear appropriate. A similar orbit may be chosen for EEP tests, motivated by the effect of the gravitational field and the absence of vibrations due to drag. The satellite should have an optical link to an optical frequency reference on the ground. The proposed gravitational-wave experiment described here would require two satellites in medium Earth orbit with a longer-baseline optical link in a calm environment to reduce gravitational noise.

In all cases, the proposed mission duration spans multiple years.

6.2 Technology development path and milestones

As indicated in the previous Section, the requirements for Cold Atoms in space basically call for three types of instrument to be developed: Atomic Clocks, Atom Interferometers and Optical Links.

In order to have those instruments introduced and accepted into a space mission, a solid development and qualification approach should be established. This is expected to be based on existing and well-proven approaches currently applied in space projects, that need to be tailored to the specific technologies and trends to reach a suitable balance between risk and mission objectives. Depending on the technologies and objectives, the approach could include in-orbit demonstrators or pathfinder missions. As a guideline, and irrespective of the type of mission (in-orbit demonstrator, pathfinder, …) a generic development approach for such instruments would typically include the following steps.

First, the scientific/mission objectives are defined, e.g., a test of the universality of free fall or a strain measurement to a certain level, together with a high-level baseline mission scenario including, e.g., the orbit, in order to derive the expected preliminary mission lifetime and environment (mechanical, thermal, magnetic field, radiation, …). In parallel, the technical requirements for the instrument(s) should be defined (functional, performance, operational, volume/mass/power, interfaces, …). This is based on both a flow-down of mission requirements (top-down) and the review of existing ground instruments and/or experiments (bottom-up). The outcome of this first step would be the issue of the consolidated mission definition and technical requirements, e.g., a certain sensitivity to differential accelerations or phase shifts induced by a gravitational wave.

Once these requirements have been reviewed and agreed by the community, each instrument can follow its own development path. This includes first the definition of the instrument architecture and its external interfaces with the spacecraft and with other instruments. Secondly, the instrument architecture definition is further refined into subsystems, modules or units, e.g., the vacuum system including peripheral optics, the laser system, or the control electronics. For each of these elements, interfaces with upper levels are defined, together with technical requirements based on a flow-down from the upper level. The granularity of the instrument architecture definition depends on the type and

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 36/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

complexity of the instrument. The outcome of this step is the issue of the instrument architecture definition and technical requirements for the subsystems.

The next step is the development of the subsystems, modules or units, whose approach is tailored to the specific element and the maturity of its underlying technology. It is usual practice to start the development at Breadboard level to demonstrate the basic functionalities and performance, and to develop further to an Elegant Breadboard and/or Engineering Model (EM). An Engineering Model is fully representative of the Flight Model in terms of form, fit and function, but does not require the use of qualified high-reliability parts.

Once all subsystems have demonstrated compliance with their technical requirements, they are integrated into the instrument, which is in turn validated and verified according to an agreed method. It is likely that instruments based on Cold Atom technologies will not be able to reach their full performance on-ground, and therefore appropriate verification methods will have to be defined (e.g., based on a combination of test and analysis/extrapolation through modelling, and/or microgravity test facilities such as a drop tower, Einstein Elevator, or zero-g Airbus flight). In the event of successful validation and verification at instrument level according to EM standards, the instrument will have reached TRL6.

From there, the instrument will follow a qualification phase. For complex instruments like the ones we are considering here, this will most probably involve a Qualification Model (QM) that is fully representative of the Flight Model in terms of build standard (using qualified high-reliability parts), and that will be subject to a qualification test campaign according to agreed qualification levels and duration. This qualification step also includes the qualification of all lower-level elements, including their materials, parts and processes, in accordance with the requirements applicable to the mission (e.g., due to launch loads, the operational environment. etc.). After successful completion of the qualification phase, the instrument will have reached TRL7, and the manufacturing of the Flight Model is released.

6.3 Technology readiness level

6.3.1 Atomic clock mission

Once optical links are in place, several technologies must be developed with improved TRL in order to launch a strontium optical lattice clock into space: optical resonators to stabilize lasers to a noise floor of 1×10^{-16} in fractional frequency, laser sources at six different wavelengths to cool, optically confine and interrogate strontium atoms, compact physics packages with a controlled black-body radiation environment, and compact frequency combs.

6.3.2 Earth observation

Key concepts have been demonstrated, including gravimeters [277] with sensitivity $4.2 \cdot 10^{-8}$ m/s^2/Hz$^{1/2}$ and gradiometers [165, 166] with sensitivity $3 \cdot 10^{-8}$ s^{-2} Hz$^{-1/2}$, both operating with ^{87}Rb atoms, as well as matter-wave collimation of BECs and BEC interferometers in a drop tower and onboard of a sounding rocket, utilising atom-chip technology for fast and robust production of ^{87}Rb BECs. The latter relied on adaptation and developments of the physics package, the laser system, and the control electronics to realise compact and robust setups. Further developments are required for operation in the specific conditions imposed in satellite missions on a component level, which is partially ongoing, but also

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 37/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

for demonstrating the desired performance which relies on the extended free fall times in a microgravity environment.

6.3.3 Fundamental physics

As outlined above, different scenarios for testing the limits of quantum mechanics with cold and condensed atoms exist. As such, the TRL differs widely for the involved components. At this time, payload for experiments on cold Rb atoms and BECs as well as quantum entanglement have been operated on satellites or the ISS. To achieve the targeted sensitivities for the discussed missions, further developments for components, such as laser systems, vacuum technology, radiation-hard electronics and autonomous operation are necessary to accommodate, for instance, high-precision Rb/K or Sr interferometers. Specific payloads, such as a setup for BEC experiments and interferometry, or components, such as frequency combs, have been successfully operated on sounding rockets and therefore feature a higher TRL than other parts. Moreover, several components require developments to cope with vibrational loads during launch, SWaP budget limitations, and other environmental conditions.

6.4 Technology evolution

Operating quantum sensors based on cold atoms on a satellite implies a completely new technology going to space. Between various mission scenarios utilising atomic clocks or atom interferometers for earth observation or fundamental physics several general building blocks are shared. They consist of a physics package with a vacuum system surrounded by optics, coils and other peripherals, a laser system with laser sources and optical benches for light distribution and switching, an electronics system with various controllers, e.g., laser drivers, and a computer for executing sequences, collecting, storing and evaluating data. This leads to potential synergies in technology developments. Within these subsystems, it is crucial to identify critical components and to start their development without delay for a mission within this decade. While different missions may require modified components imposing a new verification, this does not imply a completely new development, as the concepts, their capabilities and the approach for verification are known.

Mission-specific technology developments that may focus on performance, miniaturisation, robustness, lifetime or other relevant topics, as required, will typically follow a stepwise approach. On a conceptual level, ground-based facilities, although incompatible with deployment in space, can serve to test and verify experimental procedures, sequences, and concepts for a future mission. Initially, the system needs to be defined based on the necessary functionalities. Top-level examples are, e.g., an atom interferometer based on Rb or Sr, a Sr optical lattice clock, optical frequency dissemination based on fibres, free space, and ground-space-ground communication with mission-specific performance requirements. The next step is to identify suitable subsystems (e.g., a laser system or vacuum system/science chamber) with respect to performance, to define components (e.g., a laser head or an atom chip) and develop either as required. This may include reliability and partial environmental tests on a subsystem level, depending on the estimated criticality. The subsystems have then to be integrated, and subjected to end-to-end verification and performance tests. Subsequently, the full ground system is to be implemented and tested, including reliability and partial environmental tests. Facilities such as the microgravity simulator in Bordeaux, the drop tower in Bremen, or the Einstein Elevator in Hannover

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 38/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

offer the possibility for operating a payload or parts thereof in up to a few seconds of microgravity. Additional options for such tests are early flight tests, as enabled by a zero-g Airbus flight or a sounding rocket. Gravity, the available microgravity time of the aforementioned facilities, or special (e.g., environmental) constraints of a mission may lead to the recommendation of a pathfinder mission with opportunities sponsored by ESA, the EU or national agencies. Finally, after the development and verification steps, the definition and planning of a full mission concludes the technology evolution.

6.5 Development milestones

Preceding or in parallel to the technology developments, several scientifically-justifiable milestones can be defined to assess the maturity on a conceptual level. These are related to the demonstration of basic functionalities and concepts, including feasibility studies. A starting point is the definition of the mission concept and the scientific motivation, be it for earth observation, fundamental physics or other purposes. Critical concepts can then be tested in ground-based setups (e.g., trapping and cooling atoms with novel atom's chips/gratings), in certain cases with reduced performance (e.g., due to reduced free fall time), but still demonstrating their basic feasibility. This may include a special scheme for an atom interferometer up to a large-scale device for gravitational wave detection as in MIGA, MAGIS, ELGAR or AION. Depending on the possibility for extrapolation, the confidence in the modelling and understanding (e.g., due to different behaviour in the absence of gravitational sag, non-moving atoms), deploying a compact test setup in a microgravity facility such as the drop tower can demonstrate source performance, interrogation procedures, and detection of the atoms, accompanied by refined modelling, without the need for components qualified for a satellite mission. On the side of the mission concept, the orbit needs to be defined and evaluated accordingly, including the estimated implications on the payload and performance. The latter requires a simulator for the atom interferometer, clock or other relevant payload element to be developed. It can subsequently provide the modelling of the measurement output, including dependencies on internal and external disturbances, compared to the desired signal. Within a dedicated pathfinder mission on a satellite, demonstrating, e.g., a high-contrast atom interferometer with extended free-fall times and long-term operation, and performing statistical and systematic studies, provide further milestones for assessing the maturity of sensors.

6.5.1 In-orbit validation

To date, following the successful operation of quantum sensors in laboratories, key technologies and concepts have been demonstrated, e.g., in dedicated microgravity experiments on BEC generation and interferometry with rubidium atoms. Limited microgravity time on the available platforms prevented long-term operation, extensive statistics and a detailed systematic analysis in this special environment.

A dedicated satellite platform for a pathfinder avoids conflicts of programmatic or technical constraints due to interface or other requirements of either part of the payload in joint mission concepts. Other currently existing or planned payloads have to provide multiple functionalities and are consequently neither dedicated nor optimised for quantum sensors. Despite considerable efforts for miniaturisation and robustness for accommodation in the available microgravity platforms, a satellite mission will impose additional constraints and requirements on the payload and the operation of the quantum sensor. This

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 39/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

step towards a full-blown mission utilising quantum sensors motivates a timely pathfinder mission.

Several examples of pathfinder missions have been mentioned in earlier sections. Here we discuss as an example some aspects of a prospective mission deploying a quantum sensor with rubidium BECs on a satellite. Such a system offers the opportunity to achieve multiple goals. It enables the technology demonstration of a BEC source, beam splitters, detection, remote control, and autonomous execution of sequences on a satellite up to the uninterrupted operation of an atom interferometer over weeks to months, validating the maturity of the individual key components and functionalities. Furthermore, it can serve for testing and validating onboard data evaluation and autonomous optimisation of parameters for the source, interferometer, and other manipulations of the atoms, reducing the need for user intervention. Depending on the satellite bus and orbit, the quantum sensor is subjected to specific disturbances, necessitating mitigation strategies such as, e.g., rotation compensation with a higher dynamic range than for stationary setups on Earth. Testing such techniques for free fall times of several seconds is crucial for future high-performance sensors. Finally, only a pathfinder mission can provide a detailed performance evaluation for a satellite-based quantum sensor due to the persistent microgravity, enabling the accumulation of extensive statistics in the satellite-specific environment.

The higher level of maturity shown by the technology demonstration in such a pathfinder mission would have a direct impact on proposals for an Earth Observation mission based on Rb BEC interferometers or a fundamental physics mission testing the universality of free fall with Rb/K dual-species BEC interferometers. Future missions with optical lattice clocks would also benefit from a pathfinder mission with a Rb BEC interferometer for validating vacuum technology, parts of the control electronics, the experiment control computer, onboard data evaluation, and autonomous operation. Introducing other atomic species for atom interferometry such as Sr implies more similarities with the outlined pathfinder than a clock-type mission due to additional shared concepts, but will in addition require dedicated development activities for the atomic source and laser systems.

7 Workshop summary

As discussed in Sect. 2, the discovery of quantum mechanics in the first part of the 20th century made possible many of the technological innovations developed in the second half of the century. We are now witnessing the breakout from the laboratory of many quantum phenomena not yet applied, which offer sensor technologies of unparalleled accuracy for timing, accelerometry, gravimeters, etc.. The potential of this second quantum revolution has been recognized by the Senior Science Committee advising ESA on its Voyage 2050 programme, which has recommended an intensive programme of R&D to prepare quantum sensor technologies for deployment in space. Many of the cutting-edge developments of quantum technologies currently taking place in laboratories around Europe and elsewhere were discussed in this Workshop. Realizing their full potential in space-borne applications of immediate value to society as well as to fundamental science will require a community effort to outline high-level objectives and a road-map for achieving them that optimizes the synergies between different mission concepts.

Atomic clocks have already attained an accuracy of 10^{-18}, and the ACES mission is in an advanced state of preparation for deployment on the ISS. Atomic accelerometers have already exhibited a precision of 4×10^{-8} m/s$^2/\sqrt{\text{Hz}}$ on the ground, and offer drift-free stability extending to frequencies below 10 mHz. There has been a series of Bose–Einstein

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 40/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

Condensate (BEC) experiments in microgravity using MAIUS sounding rockets, the CAL BEC experiment has operated successfully for several years on the ISS, the MICROSCOPE experiment has tested the Weak Equivalence Principle (WEP) in space, and the MAGIS, MIGA, ZAIGA and AION atom interferometer experiments to look for ultralight dark matter and pave the way for future measurements of gravitational waves are under construction.

We present below a first draft for this community road-map, based on the cold atom achievements so far, the ongoing research, and the high-level objectives for the future. The draft road-map is centred around three topics: atomic clocks, quantum accelerometry, and atom interferometry, and is oriented towards the following objectives: next-generation standards for time standards and navigation, next-generation Earth observation and its potential for monitoring mass and climate change, and fundamental science including tests of relativity, searches for dark matter and novel measurements of gravitational waves. We stress the existence of an ongoing technology development programme including terrestrial and space-borne pathfinder projects, and the need for follow-on pathfinder experiments on Earth and in space.

Milestones in the road-map towards space clocks discussed in Sect. 3 include the completion of ACES [20] and its deployment on the ISS, to be followed by a follow-on mission such as I-SOC Pathfinder [141, 142]. Its objectives would include a comparison of ground-based optical clocks with a precision of 10^{-18} over a day (see Fig. 2), which would have applications in fundamental physics experiments as establishing a proof-of-concept for setting timescales, and in geodesy. This should be followed by a dedicated satellite in a highly elliptical orbit containing a strontium optical lattice clock with similar precision and a coherent optical link to ground, with goals similar to those of FOCOS [122]. Such a mission would enable more precise comparisons across a wider network of ground clocks, and stringent tests of general relativity. The development of coherent free-space optical links will be key, to be accompanied by the qualification of a strontium optical lattice clock for operation in space, which will require a technology development programme for several clock components. There are multiple synergies between these atomic clock missions and the requirements for a fundamental science mission such as AEDGE [5] based on strontium atom interferometry.

As discussed in Sect. 4, quantum accelerometry has exciting potential (see Figs. 6 and 7), and we envisage two atom accelerometer missions aimed at realizing it. For the technical reasons discussed in Sect. 4, deploying a quantum sensor as a passenger on a conventional geodesy mission such as MAGIC [159] would pose severe technical challenges, implying significant technological and programmatic risks for both the classical and quantum aspects of such a joint mission. For this reason, the discussions among scientific experts during the workshop led to a recommendation to launch a separate quantum pathfinder mission within this decade on a dedicated platform, with a target performance of 10^{-10} m/s$^2/\sqrt{\text{Hz}}$. Such a mission would combine the need for a test of the quantum technology in space with optimizing the results to be expected from a subsequent quantum gravimetry pathfinder mission. It would also serve as a milestone for other communities, such as that interested in applications of cold atoms to probes of fundamental physics. The success of MAGIC and the quantum pathfinder mission would pave the way for a full-fledged quantum space gravimetry mission to follow on from MAGIC, whose definition would be based on the experience gained with MAGIC and the pathfinder mission.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 41/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

We outlined in Sect. 5 the requirements for enabling the opportunities for exploring fundamental science, some of which are illustrated in Figs. 10 and 13. The first step is to construct and operate the ongoing large-scale terrestrial atom interferometer projects for fundamental science, such as MAGIS [180] in the US, MIGA [186] in France, ZAIGA [194] in China and AION [184], to be followed by one or more of the proposed km-scale experiments such as ELGAR [183] and the successors to MAGIS, ZAIGA and AION, which will serve as ultimate conceptual technology readiness demonstrators for a space-based mission such as AEDGE [5]. In parallel, there should be a satellite mission demonstrating cold and condensed atom technology in space, building on the experience with CAL [181] and MAIUS [188] and using a two-species interferometer (see [191]), based on the available rubidium and potassium sources. It will also be necessary to prepare optical frequency references for operation in space (see also Sect. 3), which will also support space missions such as LISA [207]. This will require advancing strontium development on the ground, in ground-based microgravity platforms, and in space-based pathfinder missions, including individual components, the miniaturization of subsystems and proof-of-principle missions.

8 Community proposal for an ESA road-map for cold atoms in space

We present in this Section a draft for a possible road-map for the development and deployment of cold atoms in space, formulated following the discussions during the Community Workshop that are summarised in the previous Sections of this document, which we propose for discussion with ESA, European national space agencies and interested potential partners. The proposed road-map is mainly driven by high-level scientific applications and includes synergetically the areas of Earth Observation and Fundamental Physics, as well as applications of space-borne Atomic Clocks. The status and prospects in these three areas are reviewed in Sects. 4, 5, and 3 of the workshop summary, respectively, requirements for technology development, space qualification and pathfinders are reviewed in Sect. 6, and the main points of these Sections are summarised in Sect. 7.

Figure 15 presents a high-level timeline of the main milestones of the road-map, where dates indicate the start of an activity or the launch of a space mission. A detailed discussion of the individual milestones and their interdependences is provided in Sect. 8.1 to 8.9. Although the road-map focuses on possible ESA missions and their milestones, we also mention potential synergies with other missions and options to integrate the ESA programme into a broader international context. We note that the timeline is conservative, in the sense that it could be shortened in the event of substantial international collaboration, and that uncertainties are larger for later milestones in the road-map.

8.1 Terrestrial pathfinders: underway in 2021

Scientific experiments in terrestrial environments and additional technology development, including relevant environmental testing, such as vibration or radiation hardness, are crucial to the definition, planning, and successful deployment of the space-borne missions discussed in the following road-map. Several of these developments are currently ongoing in different institutes. While some institutes focus on individual subsystems, such as more compact and robust laser systems, other efforts are underway to operate large-scale experiments, such as MAGIS-100 (Sr) in the US, MIGA (Rb) in France, AION-10 (Sr) in the UK, VLBAI (Rb and Yt) in Germany, and ZAIGA (Rb) in China, which are funded

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 42/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF



Figure 15 A first draft of the community road-map proposed for discussion with ESA. The figure shows the main milestones of the road-map, where the colour code indicates the relevance for the three main areas of Earth Observation (purple), Atomic Clocks (black) and Fundamental Physics (red). The main cold atom species targeted by the milestones are indicated with circles coloured blue for strontium and green for rubidium. These do not necessarily indicate that a milestone is based on this Cold Atom species, but may also indicate that this milestone will contribute to its space readiness. The years indicated in each milestone represent the starting date of the milestone activity or the launch date of a mission, assuming preparatory work took place before. The further the timeline extends into the future, the more uncertain the milestone starting date becomes. Further details about each of the milestones and their interdependences are given in Sect. 8

and currently under construction. These are prototypes for planned apparatuses spanning several km, such as ELGAR in Europe, SURF in the USA, and expansions of AION in the UK and ZAIGA in China. These systems could explore systematically mid-frequency range gravitational waves and would complement existing and future gravitational wave antennae, both in space and on the ground. In addition, these large-scale projects could perform research in other areas, e.g., to search for oltralight bosonic dark matter that is weakly coupled to Standard Model particles, and to test the Equivalence Principle. These terrestrial Cold Atom pathfinder projects will complement the other pathfinders listed in our road-map and space-borne cold atom experiments such as CACES, MAIUS, CAL and BECCAL, as well as the successful pathfinders for key underlying optical technologies such as FOKUS, KALEXUS and JOKARUS.

8.2 Development programme launch: 2022

In parallel with the construction, commissioning and operation of these terrestrial pathfinder experiments, it will be necessary to start without delay a programme to raise the TRLs of the associated technologies, demonstrating their robustness and reliability, and reducing their size, weight and power requirements, so that they become realistic candidates for deployment in space-borne experiments. In addition to the sensor technologies used directly in the experimental apparatus, such as atomic clocks, accelerometers and interferometers, it will be necessary to raise the TRLs of ancillary technologies such as optical links. This technology development programme will be essential for the deployment in space of atomic clocks, applications to Earth Observation and experiments in

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 43/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

fundamental physics. Technologies using rubidium are currently more mature than those using strontium, but both will require substantial development effort.

In addition to a development programme building systematically on the ongoing terrestrial pathfinders outlined in Sect. 8.1, a pathway that has proven to be successful in increasing TRLs for space operation is provided by terrestrial microgravity environments. These include, for instance, the Einstein Elevator in Hanover, the drop tower in Bremen, parabolic flights operated by Novespace, and sounding rocket missions from Kiruna. Operation within these environments requires the experimental setups to ensure miniaturization of the setup and perform environmental tests. This enables aspects of fundamental research to be performed and technology necessary for space operation to be developed. The scientific output supplements and focuses planned space-based missions and can bridge gaps until space-based missions become a reality. Prominent examples of such experiments are the I.C.E experiment in parabolic flights, the QUANTUS and PRIMUS experiments in the drop tower, and the MAIUS and JOKARUS campaigns in sounding rocket flights. Other important stepping-stones for fundamental research and technology development for further space-based experiments are the cold atom experiments on the ISS, namely CAL and BECCAL. With CAL being currently mounted to the ISS and BECCAL being prepared for deployment within this decade, they pave the way towards more complex and specialized experiments and increase the TRLs of individual components as well as complete setups. Establishing this Development Programme will be essential for the success of the following Road-Map milestones outlined in Sects. 8.3–8.9.

8.3 ACES: 2025

The ACES experiment is currently being prepared for launch by 2025. It will measure the gravitational redshift between the Project of atomic clock using in orbit atom cooling (PHARAO) clock on-board the ISS and clocks on Earth, improving on current measurements of the redshift effect by an order of magnitude. In addition to providing the first in-orbit demonstration of the operation of a cold atom clock using cesium, as well as the operation of a hydrogen maser, it will also pioneer the deployment of improved optical links for atomic clocks in space. ACES will serve as a pathfinder for future projects deploying atomic clocks in space (Sect. 8.8) as well as providing important new tests of general relativity, paving the way for future cold atom experiments on fundamental physics in space (Sects. 8.7 and 8.9). The experience gained with ACES will be particularly relevant for subsequent experiments using strontium clocks in space.

8.4 Earth observation pathfinder mission: 2029

As discussed in the main text, gravimeters and accelerometers based on cold atom technology show great promise for future Earth Observation missions, in view of the high precision, stability and low-frequency performance they offer. However, we judge that it would be premature to plan a standalone full Earth Observation mission using cold atoms, and that a prior pathfinder mission will be required. For the reasons discussed in the text, we consider that it not be advantageous to combine this quantum pathfinder mission with an Earth Observation mission using classical technology. While relevant primarily for a subsequent Earth Observation mission (Sect. 8.6), this pathfinder mission will also pave the way for subsequent fundamental science missions using atomic clocks. It would

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 44/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

use rubidium atoms, whose terrestrial development is relatively mature, and its operational experience would be relevant to an STE-QUEST-like fundamental physics mission (Sect. 8.7), which would also use rubidium.

8.5 I-SOC pathfinder: 2030

I-SOC Pathfinder would push further the microwave and optical link technologies being developed for ACES (Sect. 8.3), with a view to continue the operation of a worldwide network of optical clocks on the ground to test fundamental laws of physics, to develop applications in geodesy and time & frequency transfer, and to demonstrate key technologies for future atomic clock missions in space. The main objective of I-SOC Pathfinder will be to increase the versatility of atomic clocks in space, acting as a pathfinder for a subsequent mission to exploit the full capabilities of atomic clocks, also for applications in fundamental physics based on strontium. Its operation will support other technology developments by operating optical links from the space station to ground.

8.6 Earth observation mission: 2036

This would be a standalone Earth Observation mission to deliver the prospective improvements in spatial and temporal resolution over classical Earth Observation missions such as GRACE and GOCE that are illustrated in Fig. 7. The full definition of the mission will be informed by the lessons learned from the next-generation classical MAGIC mission and the Pathfinder mission outlined in Sect. 8.4. Although the primary purpose of this mission would be Earth Observation, the technical developments it requires will also benefit the STE-QUEST-like fundamental physics mission outlined in Sect. 8.7, which will also use rubidium.

8.7 STE-QUEST-like M-class mission: 2037

Building on experience with the successful MICROSCOPE mission and development work undertaken for the previous STE-QUEST proposal, this mission would deploy a double atom interferometer with rubidium and potassium "test masses" in quantum superposition to test the Einstein equivalence Principle (universality of free fall, UFF) and search for ultralight dark matter. It would use a single satellite in a 700 km circular orbit, and offers the possibility of probing the Einstein Equivalence Principle, with a precision $\mathcal{O}(10^{-17})$, as seen in Fig. 10. This is essentially a fundamental physics mission that would, however, take advantage of the synergy with the Rb-based Earth Observation missions described in Sects. 8.4 and 8.6 as well as experiments performed on the ISS in the CAL and BECCAL facilities and those performed in terrestrial microgravity facilities outlined in Sect. 8.1, e.g., QUANTUS, PRIMUS, MAIUS and I.C.E.

Programmatically, this mission profile, supported by the comprehsive and ambitious development programme outlined in Sect. 8.2, would fit well in the scope of an ESA M-class mission. The time-frame of the call announced recently by ESA's Science Directorate [278] would be fully in line with the timeline proposed here of a STE-QUEST-like mission launching in 2037, and there is widespread support in the cold atom community for the proposal of such a mission profile in response to this ESA call.

8.8 Atomic clock mission: 2038

This mission would translate the high precision of the most accurate atomic clocks shown in Fig. 2 into a global time standard that would take metrology to the next level, with cor-

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 45/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

responding advantages for navigation, geodesy and fundamental physics. It would use a dedicated satellite in a highly elliptical orbit containing a strontium optical lattice clock with a 10^{-18} systematic uncertainty and $10^{-16}/\sqrt{\tau}$ stability, and a coherent optical link to ground. In addition to enable more precise comparisons across a wider network of ground clocks, it would provide direct searches for new physics including stringent tests of general relativity than possible with previous missions. The technology development for a Sr optical lattice clock is ongoing, and is expected to continue and bring to flight maturity the critical elements of the instrument, e.g., lasers, reference cavity and atomic source. The optical clock mission will build upon the operational experience previously gained with ACES (Sect. 8.3) and I-SOC Pathfinder (Sect. 8.5) and on a continuously improving network of ground-based optical clocks connected to the space clock via a coherent optical link. We remark that the time frame envisaged in Fig. 15 could be substantially shortened by cooperation with other space agencies that show a strong interest in the development of optical clocks for space [122], which might allow such a mission to be realised on a substantially shorter time-scale.

8.9 AEDGE-like L-class mission: 2045

This would be a fundamental science mission based on atom interferometry using strontium. It would provide the ultimate sensitivity to ultralight dark matter, as seen in Fig. 10, and gravitational waves in the deciHz band intermediate between the maximum sensitivities of LIGO/Virgo/KAGRA and other terrestrial laser interferometers and the spaceborne laser interferometer LISA, as seen in Fig. 13. Thus it would complement the capabilities of these other detectors and offer interesting synergies when operating as part of an international network. The configurations considered assume two satellites in medium Earth orbit separated by $\sim 40,000$ km using atom clouds that might be either inside or possibly outside the satellites. It would be based upon developments pioneered by many of the pathfinders described above including the terrestrial atom interferometers now under construction (Sect. 8.1), ACES (Sect. 8.3), I-SOC Pathfinder (Sect. 8.5) and the proposed dedicated atomic clock mission (Sect. 8.8). It would also have many elements in common with missions using rubidium atoms such as the dedicated Earth Observation and STE-QUEST-like missions (Sects. 8.4, 8.6 and 8.7), such as techniques to minimize the size, weight and power requirements for atomic clocks, as well as optical links.

8.10 Road-map summary

We have highlighted in the Section the important synergies and interdependences between fundamental science and Earth Observation missions based on cold atom technology, as well as the key roles played by terrestrial pathfinder experiments and the importance of a vigorous development programme. We note that are strong, diverse, and well-funded terrestrial pathfinder programmes for deploying cold atom technology already well underway in many countries. ESA leadership of a technical development programme for cold atoms in space, building on the advances made by the terrestrial pathfinders to raise the TRLs of the associated technologies for space readiness, would provide the backbone for future cold atom missions.

This development programme will complement ACES and the Earth Observation and I-SOC pathfinder missions, which will pave the way for ultimate Earth Observation and atomic clock missions targeting next-generation geodesy, time-keeping and fundamental

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 46/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

physics to launch around 2036 and 2038. The technical developments in cold atom technology and ancillary technologies such as optical links will also prepare the ground for an STE-QUEST-like M-Class mission around 2037 to probe general relativity via tests of the equivalence principle (universality of free fall) and search for ultralight dark matter. An STE-QUEST-like M-class mission around 2037 is being proposed by cold atom community members in response to the recent call by ESA's Science Directorate [278], and it enjoys widespread and strong support in cold atom community. The ultimate sensitivity to dark matter, along with unique capabilities for measuring gravitational waves, would be provided by an AEDGE-like L-class mission in the 2040 s.

We propose this road-map for discussion by the interested cold atom, Earth Observation, fundamental physics and other prospective scientific user communities, together with ESA and national space and research funding agencies.

Acknowledgements
We thank the CERN Quantum Technology Initiative for their support of the workshop that laid the basis for this paper.

Availability of data and materials
Not applicable. For all requests relating to the paper, please contact author Oliver Buchmueller.

Declarations

Ethics approval and consent to participate
Not applicable.

Consent for publication
Consents given by the authors at the time of submission

Competing interests
The authors declare no competing interests.

Author contributions
The author OB is the contact person for this paper. The following authors are Section Editors and/or Workshop Organisers: AB, KB, PB, OB, LC, OC, MLC, AMC, ADR, MD, JE, RF, NG, RH, SK, TL, CL, FM, EM, EMR, SS, CS, CS, GMT, WK, EW, PW, LW. All authors discussed the content of the paper and contributed to the writing of the manuscript. All authors read and approved the final manuscript.

Author details
[1]Higher Polytechnic School, University of the Balearic Islands, Valldemossa Road, Palma de Mallorca, 07122, Spain. [2]University of Washington, Physics-Astronomy Building, 15th Ave NE, Seattle, WA 98195-1560, USA. [3]University of South Carolina, Main Street, Columbia, SC 29208, USA. [4]Imperial College London, Prince Consort Road, London, SW7 2AZ, UK. [5]CERN, CH-1211, Geneva, 23, Switzerland. [6]Aarhus University, Ny Munkegade, 8000 Aarhus C, Denmark. [7]King's College London, Strand, London, WC2R 2LS, UK. [8]University of Belgrade, Pregrevica 118, 11080, Belgrade, Serbia. [9]Dipartimento di Fisica e Astronomia, Università di Firenze, Via Sansone 1, 50019 Sesto Fiorentino, Firenze, Italy. [10]European Laboratory for Non-Linear Spectroscopy (LENS), Università degli Studi di Firenze, Via Giovanni Sansone, 1, 50019 Sesto Fiorentino, Firenze, Italy. [11]LIGO, California Institute of Technology, 1201 E California, Pasadena, CA 91125, USA. [12]Institute of Nuclear and Particle Physics, NCSR Demokritos, Agia Paraskevi, 15310, Greece. [13]University of Pisa, Largo Lazzarino, Pisa, 56122, Italy. [14]Kitzbühel Centre for Physics, Kitzbühel, Austria. [15]University of Trieste, Strada Costiera 11, 34151, Trieste, Italy. [16]Istituto Nazionale di Fisica Nucleare, Trieste Section, Via Valerio 2, 34127, Trieste, Italy. [17]LP2N, Laboratoire Photonique, Numérique et Nanosciences, Université Bordeaux-IOGS-CNRS: UMR 5298, F-33400, Talence, France. [18]SYRTE, Observatoire de Paris, Université PSL, Sorbonne Université, LNE, 61 avenue de l'Observatoire, 75014, Paris, France. [19]DPHY, ONERA, Université Paris-Saclay, Chemin de la Hunière—BP80100, F-91123, Palaiseau, France. [20]Department of Theoretical Physics, University of Valencia, Joint Centre Univ. Valencia-CSIC, 46100, Burjassot-Valencia, Spain. [21]IFIC, University of Valencia, Joint Centre Univ. Valencia-CSIC, 46100, Burjassot-Valencia, Spain. [22]Rutherford Appleton Laboratory, Harwell Campus, Didcot, OX11 0QX, UK. [23]University of Strathclyde, Glasgow, G4 0NG, UK. [24]Grup de Física Teòrica, Departament de Física, Universitat Autònoma de Barcelona, 08193, Bellaterra, Spain. [25]Institut de Fisica d'Altes Energies, The Barcelona Institute of Science and Technology, Campus UAB, 08193, Bellaterra (Barcelona), Spain. [26]University of Birmingham, Birmingham, B15 2TT, UK. [27]German Aerospace Center (DRL), Institute of Quantum Technologies, Wilhelm-Runge-Straße 10, 89081, Ulm, Germany. [28]Institute of Microelectronics, Ulm University, Albert-Einstein-Allee 43, 89081, Ulm, Germany. [29]Department of Physics, Clarendon Laboratory, University of Oxford, South Parks Road, Oxford, OX1 3PU, UK. [30]Johannes Gutenberg-Universität Mainz, Helmholtz-Institut Mainz, GSI Helmholtzzentrum für Schwerionenforschung, 5128, Mainz, Germany. [31]Department of Physics, University of California, Berkeley, CA 94720, USA. [32]Instituto de Telecomunicações, Instituto Superior Técnico, Av. Rovisco Pais, Torre Norte, Lisboa, 1049-001, Portugal. [33]University of Liverpool, Liverpool, L69 7ZE, UK. [34]European Space Agency, Keplerlaan 1, PO Box 299, NL-2200, AG Noordwijk, The Netherlands.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 47/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

[35]Physikalisch-Technische Bundesanstalt, Bundesallee 100, Braunschweig, 38116, Germany. [36]University of Sussex, Brighton, BN1 9QH, UK. [37]Istituto Nazionale di Ricerca Metrologica, Strada delle Cacce 91, Turin, 10035, Italy. [38]Institute of Space Science, 409 Atomistilor Street, Bucharest, Magurele, Ilfov, 077125, Romania. [39]University of St Andrews, North Haugh, St Andrews, KY16 9SS, UK. [40]Leibniz Universität Hannover, Welfengarten 1, Hannover, 30167, Germany. [41]Fermilab, Batavia, Illinois, USA. [42]Stanford University, Stanford, CA 94305, USA. [43]Indian Institute of Technology, New Delhi, India. [44]Peking University, Beijing, 100871, China. [45]Hefei National Laboratory for Physical Sciences at the Microscale and Department of Modern Physics, University of Science and Technology of China, Hefei, 230026, China. [46]Shanghai Branch CAS Center for Excellence in Quantum Information and Quantum Physics, University of Science and Technology of China, Shanghai, 201315, China. [47]Shanghai Research Center for Quantum Sciences, Shanghai 201315, China. [48]Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, Largo Pontecorvo 3, I-56125, Pisa, Italy. [49]Department of Physics and Astronomy, University of California, Riverside, CA 92521, USA. [50]Antwerp University, B 2610, Wilrijk, Belgium. [51]Oak Ridge National Laboratory, 1 Bethel Valley Road, Oak Ridge, TN 37830, USA. [52]University of Nevada, Reno, Nevada 89557, USA. [53]Italian Space Agency, Via del Politecnico, Rome, 00133, Italy. [54]University of Nis, Visegradska 33, Nis, Serbia. [55]Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA. [56]Foundation for Research and Technology-Hellas, 100 N. Plastira Street, Vassilika Vouton, GR 70013, Heraklion, Crete, Greece. [57]The University of Tokyo Institutes for Advanced Study, The University of Tokyo, Kashiwa, Chiba, 277-8583, Japan. [58]Los Alamos National Laboratory, Los Alamos, New Mexico, USA. [59]Faculty of Science, Mohammed V University in Rabat, 4 Avenue Ibn Battouta, B.P. 1014 RP, Rabat, Morocco. [60]The Johns Hopkins University, 3400 N Charles St, Baltimore, MD 21218, USA. [61]University College London, Gower Street, London, WC1E 6BT, UK. [62]School of Physics, University of New South Wales, Sydney, 2052, Australia. [63]Technical University of Denmark, Kongens Lyngby, 2800, Denmark. [64]University of Nottingham, Nottingham, NG7 2RD, UK. [65]Northwestern University, Evanston, USA. [66]The Pennsylvania State University, University Park, PA 16802, USA. [67]University of Cambridge, J.J. Thomson Avenue, Cambridge, CB3 0HE, UK. [68]National Physical Laboratory, Hampton Road, Teddington, TW11 0LW, UK. [69]Department of Physics and Astronomy, Wayne State University, Detroit, MI 48202, USA. [70]Laboratoire Kastler Brossel, Sorbonne Université, CNRS, ENS-Université PSL, Collège de France, **???**, France. [71]Humboldt-Universität zu Berlin, Newtonstraße 15, Berlin 12489, Germany. [72]Okinawa Institute of Science and Technology, Okinawa, Japan. [73]Universität Bremen, Am Fallturm 2, 28359, Bremen, Germany. [74]University of Bergen, Allégaten 55, 5007, Bergen, Norway. [75]Jožef Stefan Institute, Jamova 39, 1000, Ljubljana, Slovenia. [76]University of Zurich, Winterthurerstrasse 190, 8057, Zurich, Switzerland. [77]Vilnius University, Saulėtekio 3, Vilnius, LT-10257, Lithuania. [78]University of Ljubljana, Jadranska ulica 19, 1000, Ljubljana, Slovenia. [79]Physics Division, National Technical University of Athens, Zografou, Athens, 15780, Greece. [80]University of Latvia, Riga, LV-1004, Latvia. [81]Teledyne e2v, 106 Waterhouse Lane, Chelmsford, CM1 2QU, UK. [82]University of Wisconsin, Madison, WI 53706, USA. [83]Faculty of Physics, Warsaw University of Technology, ul. Koszykowa 75, 00-662, Warsaw, Poland. [84]University of the Witwatersrand, Johannesburg, Wits, 2050, South Africa. [85]Indian Institute of Science Education and Research, Bhopal, 462066, India. [86] Department of Physics, Brown University, Providence, RI 02912, USA. [87]National Institute of Standards and Technology, 325 Broadway Street, Boulder, Colorado 80305, USA. [88]Centre National d'Etudes Spatiales, 18 Avenue Edouard Belin, 31400, Toulouse, France. [89]University of Warsaw, ul. Pasteura 5, 02-093, Warsaw, Poland. [90]Purdue University, West Lafayette, IN 47907, USA. [91]Autonomous University of Aguascalientes, Av. Universidad 940, Ciudad Universitaria, Aguascalientes, C. P. 20131, Mexico. [92]Potsdam Institute for Climate Impact Research, Member of the Leibniz Association, 14473, Potsdam, Germany. [93]Università degli Studi di Salerno, Via Giovanni Paolo II 132, I-84084, Fisciano (SA), Italy. [94]Bates College, Lewiston, ME 04240, USA. [95]Fayoum University, El-Fayoum, Egypt. [96]Politecnico di Milano, Piazza Leonardo da Vinci 32, 20133, Milano, Italy. [97]University of Warwick, Coventry, CV4 7AL, UK. [98]Department of Physics, Koç University, İstanbul, Sarıyer, 34450, Turkey. [99]University of Glasgow, Glasgow, G12 8QQ, UK. [100]ColdQuanta, Boulder, Colorado, USA. [101] AMOLF, Science Park 1098 XG, Amsterdam, The Netherlands. [102]University of Crete, P.O. Box 2208, 71003, Heraklion, Greece. [103]Queen's University, Belfast, BT7 1NN, UK. [104]Center for Theoretical Physics PAS, Aleja Lotnikow 32/46, 02-668, Warsaw, Poland. [105]Tyndall National Institute, University College Cork, Cork, Ireland. [106]Stevens Institute of Technology, Castle Point on the Hudson, Hoboken, NJ 07030, USA. [107]Stockholm University, AlbaNova, SE-10691, Stockholm, Sweden. [108]Department of Physics and Astronomy, University of Manchester, Manchester, M13 9PL, UK. [109]Hoursec, Energy Efficient AI On-Chip, Uetlibergstrasse 111B, Zurich, 8045, Switzerland. [110]University of Bologna, Viale Berti-Pichat 6/2, Bologna, I-40126, Italy. [111]Department of Physics, The University of Arizona, Tucson, AZ 85721-0081, USA. [112]Scuola Internazionale Superiore di Studi Avanzati, Via Bonomea 265, I-34136, Trieste, Italy. [113]ETH Zurich, Robert-Gnehm-Weg 15, CH-8093, Zurich, Switzerland. [114]University of Delaware, Newark, Delaware 19716, USA. [115]Institute of Physics of the Czech Academy of Sciences, Na Slovance 2, 182 21, Praha 8, Czechia. [116]Istituto Nazionale di Fisica Nucleare, Sezione di Firenze, via Sansone 1, Sesto Fiorentino, Firenze, Italy. [117]Van der Waals-Zeeman Institute, University of Amsterdam, Science Park 904, 1098XH, **???**, The Netherlands. [118]Heinrich-Heine-Universität Düsseldorf, Germany. [119]Laboratoire Univers et Théorie, Observatoire de Paris-PSL, 92190, Meudon, France. [120]University of Vienna, Boltzmanngasse 5, A-1090, Vienna, Austria. [121]Polish Academy of Sciences, PL 31-342, Krakow, Poland. [122]CEICO–FZU, Institute of Physics of the Czech Academy of Sciences, Na Slovance 1999/2, 182 21, Prague, Czechia. [123]INO-CNR, Largo Enrico Fermi 2, 50125, Firenze, Italy. [124]Institut de Ciències de l'Espai (ICE, CSIC), Campus UAB, Carrer de Can Magrans s/n, 08193, Cerdanyola del Vallès, Spain. [125]Institut d'Estudis Espacials de Catalunya (IEEC), Edifici Nexus, Carrer del Gran Capità 2–4, despatx 201, 08034, Barcelona, Spain. [126]Istituto Nazionale di Fisica Nucleare, Sezione di Genova, via Dodecaneso 33, 16146, Genova, Italy. [127]National and Kapodistrian University of Athens, Athens, Greece. [128]The University of Sydney, **???**, NSW 2006, Australia. [129]High Energy Physics Group, 513 Keen Building, Dept. of Physics, Florida State University, 77 Chieflan Way, Tallahassee, FL 32306, USA. [130]Department of Physics, University of Basel, **???**, Switzerland. [131]Department of Physics and Astronomy, University of California, Irvine, CA 92697-4575, USA. [132]Thales Alenia Space, 26 av. J.-F. Champollion, 31100, Toulouse, France. [133]Institut de Fisica d'Altes Energies, The Barcelona Institute of Science and Technology, Campus UAB, 08193, Bellaterra (Barcelona), Spain. [134]DiSPeA, Virgo Group, University of Urbino, via S. Chiara, Urbino, I61029, Italy. [135]Institute for Applied Physics, Technical University of Darmstadt, Hochschulstrasse 4A, D-64289, Darmstadt, Germany. [136]EQOQI, QUANTUM, Institute of Physics, Johannes Gutenberg-University Mainz, 55122, Mainz, Germany. [137]Photonics Laboratory, Engineering Physics, Münster University of Applied Sciences, Stegerwaldstr. 39, 48565, Steinfurt, Münster, Germany. [138]University of Malta, Msida, MSD 2080, Malta. [139]Faculty of Engineering and Natural Sciences, International University of Sarajevo, Hrasnička cesta 15, 71210, Ilidža, Sarajevo, Bosnia-Herzegovina. [140]Department of Physics, National Taiwan University, No. 1 Sect. 4 Roosevelt Road, Taipei, 10617, Taiwan. [141]Wuhan Institute of Physics and Mathematics, Innovation Academy for Precision

UNCORRECTED PROOF

AUTHOR'S PROOF

Measurement Science and Technology, Chinese Academy of Sciences, Wuhan, 430071, China. [142]University of Cincinnati, Cincinnati, Ohio 45221, USA.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 12 April 2022 Accepted: 8 October 2022

References

1. CERN. Community Workshop on Cold Atoms in Space. https://indico.cern.ch/event/1064855/.
2. Voyage 2050: final recommendations from the Voyage 2050 Senior Committee. https://www.cosmos.esa.int/documents/1866264/1866292/Voyage2050-Senior-Committee-report-public.pdf/e2b2631e-5348-5d2d-60c1-437225981b6b?t=1623427287109.
3. CERN. Workshop on Atomic Experiments for Dark Matter and Gravity Exploration. https://indico.cern.ch/event/830432/.
4. Bertoldi A, Bongs K, Bouyer P, Buchmueller O, Canuel B, Caramete L-I et al. AEDGE: atomic experiment for dark matter and gravity exploration in space. Exp Astron. 2021;51:1417–26.
5. El-Neaj YA, Alpigiani C, Amairi-Pyka S, Araújo H, Balaž A, Bassi A et al. AEDGE: atomic experiment for dark matter and gravity exploration in space. EPJ Quantum Technol. 2020;7:6.
6. Battelier B, Bergé J, Bertoldi A, Blanchet L, Bongs K, Bouyer P et al. Exploring the foundations of the physical universe with space tests of the equivalence principle. Exp Astron. 2021;51:1695–736.
7. Kaltenbaek R, Acin A, Bacsardi L, Bianco P, Bouyer P, Diamanti E et al. Quantum technologies in space. Exp Astron. 2021;51:1677–94.
8. Tartaglia A, Bassan M, Casalino L, Crosta M, Lattanzi M, Lorenzini E et al. Detecting the gravito-magnetic field of the dark halo of the milky way—the ladahad mission concept. Exp Astron. 2021;51:1773–91.
9. Bergé J, Baudis L, Brax P, Chiow S-W, Christophe B, Doré O et al. The local dark sector. Exp Astron. 2021;51:1737–66.
10. Schneider J. Quantum correlations at Earth-moon distance. Exp Astron. 2021;51:1767–72.
11. Sedda MA, Berry CPL, Jani K, Amaro-Seoane P, Auclair P, Baird J et al. The missing link in gravitational-wave astronomy. Exp Astron. 2021;51:1427–40.
12. Sedda MA, Berry CPL, Jani K, Amaro-Seoane P, Auclair P, Baird J et al. The missing link in gravitational-wave astronomy: discoveries waiting in the decihertz range. Class Quantum Gravity. 2020;37:215011.
13. Floberghagen R, Fehringer M, Lamarre D, Muzi D, Frommknecht B, Steiger C et al. Mission design, operation and exploitation of the gravity field and steady-state ocean circulation explorer mission. J Geod. 2011;85:749–58.
14. Tapley BD, Watkins MM, Flechtner F, Reigber C, Bettadpur S, Rodell M et al. Contributions of GRACE to understanding climate change. Nat Clim Change. 2019;9:358–69.
15. Kornfeld RP, Arnold BW, Gross MA, Dahya NT, Klipstein WM, Gath PF et al. Grace-fo: the gravity recovery and climate experiment follow-on mission. J Spacecr Rockets. 2019;56:931–51. https://doi.org/10.2514/1.A34326.
16. Pail R, Bingham R, Braitenberg C, Eicker A, Horwath M, Ivins E, et al. Observing mass transport to understand global change and to benefit society: science and user needs-an international multi-disciplinary initiative for IUGG, Deutsche Geodätische Kommission: B (Angewandte Geodäsie) Heft Nr. 320: Reihe B, Angewandte Geodäsie, Elektronische Ressource. 2015. https://publikationen.badw.de/en/044436255.
17. ESA AGENDA 2025: make space for Europe. https://esamultimedia.esa.int/docs/ESA_Agenda_2025_final.pdf.
18. Delva P, Puchades N, Schönemann E, Dilssner F, Courde C, Bertone S et al. Gravitational redshift test using eccentric Galileo satellites. Phys Rev Lett. 2018;121:231101.
19. Herrmann S, Finke F, Lülf M, Kichakova O, Puetzfeld D, Knickmann D et al. Test of the gravitational redshift with Galileo satellites in an eccentric orbit. Phys Rev Lett. 2018;121:231102.
20. Cacciapuoti L, Salomon C. The ACES experiment. Eur Phys J Spec Top. 2009;172:57.
21. Cacciapuoti L, Armano M, Much R, Sy O, Helm A, Hess MP et al. Testing gravity with cold-atom clocks in space. Eur Phys J D. 2020;74:164.
22. Savalle E, Guerlin C, Delva P, Meynadier F, le Poncin-Lafitte C, Wolf P. Gravitational redshift test with the future ACES mission. Class Quantum Gravity. 2019;36:245004.
23. Colladay D, Kostelecky VA. Lorentz violating extension of the standard model. Phys Rev D. 1998;58:116002. arXiv:hep-ph/9809521.
24. Lämmerzahl C, Braxmaier C, Dittus H, Müller H, Peters A, Schiller S. Kinematical test theories for special relativity: a comparison. Int J Mod Phys D. 2002;11:1109–36.
25. Delva P, Lodewyck J, Bilicki S, Bookjans E, Vallet G, Targat RL et al. Test of special relativity using a fiber network of optical clocks. Phys Rev Lett. 2017;118:221102. arXiv:1703.04426.
26. Sanner C, Huntemann N, Lange R, Tamm C, Peik E, Safronova MS et al. Optical clock comparison for Lorentz symmetry testing. Nature. 2019;567:204–8.
27. Lange R, Huntemann N, Rahm JM, Sanner C, Shao H, Lipphardt B et al. Improved limits for violations of local position invariance from atomic clock comparisons. Phys Rev Lett. 2021;126:011102.
28. Roberts BM, Blewitt G, Dailey C, Murphy M, Pospelov M, Rollings A et al. Search for domain wall dark matter with atomic clocks on board global positioning system satellites. Nat Commun. 2017;8:1195. arXiv:1704.06844v3.
29. Roberts BM, Delva P, Al-Masoudi A, Amy-Klein A, Bærentsen C, Baynham CFA et al. Search for transient variations of the fine structure constant and dark matter using fiber-linked optical atomic clocks. New J Phys. 2020;22:093010.
30. Barontini G, Boyer V, Calmet X, Fitch N, Forgan E, Godun R et al. QSNET, a network of clock for measuring the stability of fundamental constants. In: Quantum technology: driving commercialisation of an enabling science II. vol. 11881. Bellingham: SPIE; 2021. p. 63–6.
31. Barontini G, et al. Measuring the stability of fundamental constants with a network of clocks. arXiv:2112.10618.
32. Tsai Y-D, Eby J, Safronova MS. arXiv:2112.076741.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 49/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

33. Banerjee A, Kim H, Matsedonskyi O, Perez G, Safronova MS. Probing the relaxed relaxion at the luminosity and precision frontiers. J High Energy Phys. 2020;07:153.
34. Kolkowitz S, Pikovski I, Langellier N, Lukin MD, Walsworth RL, Ye J. Gravitational wave detection with optical lattice atomic clocks. Phys Rev D. 2016;94:124043.
35. Grebing C, Al-Masoudi A, Dörscher S, Häfner S, Gerginov V, Weyers S et al. Realization of a timescale with an accurate optical lattice clock. Optica. 2016;3:563. arXiv:1511.03888.
36. Yao J, Sherman JA, Fortier T, Leopardi H, Parker T, McGrew W et al. Optical-clock-based time scale. Phys Rev Appl. 2019;12:044069.
37. Riedel F, Al-Masoudi A, Benkler E, Dörscher S, Gerginov V, Grebing C et al. Direct comparisons of European primary and secondary frequency standards via satellite techniques. Metrologia. 2020;57:045005. arXiv:1910.06736.
38. Lisdat C, Grosche G, Quintin N, Shi C, Raupach S, Grebing C et al. A clock network for geodesy and fundamental science. Nat Commun. 2016;7:12443.
39. Origlia S, Schiller S, Pramod MS, Smith L, Singh Y, He W et al. Development of a strontium optical lattice clock for the SOC mission on the ISS. In: Stuhler J, Shields AJ, editors. Comptes rendus physique. vol. 9900. 2016. p. 990003. http://proceedings.spiedigitallibrary.org/proceeding.aspx?doi=10.1117/12.2229473. arXiv:1503.08457v1.
40. Denker H, Timmen L, Voigt C, Weyers S, Peik E, Margolis HS, et al. Geodetic methods to determine the relativistic redshift at the level of 10^{-18} in the context of international timescales: a review and practical results. J Geod. 2017; 1–30.
41. IERS. The International Terrestrial Reference Frame (ITRF). https://www.iers.org/IERS/EN/DataProducts/ITRF/itrf.html.
42. IERS. ITRF2020 Call for participation. https://www.iers.org/IERS/EN/DataProducts/ITRF/itrf.html.
43. United Nations General Assembly. United Nations Resolution a/re/69/266: a global geodetic reference frame for sustainable development. https://ggim.un.org/documents/a_res_69_266_e.pdf.
44. Reigber C, Schwintzer P, Lühr H. The CHAMP geopotential mission. Boll Geofis Teor Appl. 1999;40:285.
45. Tapley B, Bettadpur S, Watkins M, Reigber C. The gravity recovery and climate experiment: mission overview and early results. Geophys Res Lett. 2004;31.
46. Wu H, Müller J, Lämmerzahl C. Clock networks for height system unification: a simulation study. Geophys J Int. 2019;216:1594–607.
47. Wu H, Müller J. Towards an international height reference frame using clock networks, IUGG2019 General Assembly, International Association of Geodesy Symposia. https://doi.org/10.1007/1345_2020_97.
48. Brewer SM, Chen J-S, Hankin AM, Clements ER, Chou CW, Wineland DJ et al. Al+ quantum-logic clock with a systematic uncertainty below 10–18. Phys Rev Lett. 2019;123:1–5. arXiv:1902.07694.
49. Schioppo M, Brown RC, McGrew WF, Hinkley N, Fasano RJ, Beloy K et al. Ultrastable optical clock with two cold-atom ensembles. Nat Photonics. 2017;11:48–52.
50. Schwarz R, Dörscher S, Al-Masoudi A, Benkler E, Legero T, Sterr U et al. Long term measurement of the ^{87}Sr clock frequency at the limit of primary Cs clocks. Phys Rev Res. 2020;2:033242.
51. Schnatz H, Lipphardt B, Helmcke J, Riehle F, Zinner G. First phase-coherent frequency measurement of visible radiation. Phys Rev Lett. 1996;76:18–21.
52. Udem T, Huber A, Gross B, Reichert J, Prevedelli M, Weitz M et al. Phase-coherent measurement of the hydrogen 1S-2S transition frequency with an optical frequency interval divider chain. Phys Rev Lett. 1997;79:2646–9.
53. Niering M, Holzwarth R, Reichert J, Pokasov P, Udem T, Weitz M et al. Measurement of the hydrogen 1S-2S transition frequency by phase coherent comparison with a microwave cesium fountain clock. Phys Rev Lett. 2000;84:5496–9.
54. Udem T, Diddams SA, Vogel KR, Oates CW, Curtis EA, Lee WD et al. Absolute frequency measurements of the Hg^+ and Ca optical clock transitions with a femtosecond laser. Phys Rev Lett. 2001;86:4996–9.
55. Margolis H, Barwood G, Huang G, Klein H, Lea S, Szymaniec K et al. Hertz-level measurement of the optical clock frequency in a single $^{88}Sr^+$ ion. Science. 2004;306:1355–8.
56. Oskay W, Diddams S, Donley E, Fortier T, Heavner T, Hollberg L, et al. Single-Atom Optical Clock with High Accuracy, Phys Rev Lett. 2006;97.
57. Boyd MM, Ludlow AD, Blatt S, Foreman SM, Ido T, Zelevinsky T et al. ^{87}Sr lattice clock with inaccuracy below 10^{-15}. Phys Rev Lett. 2007;98:083002.
58. Stalnaker JE, Diddams SA, Fortier TM, Kim K, Hollberg L, Bergquist JC et al. Optical-to-microwave frequency comparison with fractional uncertainty of 10^{-15}. Appl Phys B, Lasers Opt. 2007;89:167–76.
59. Poli N, Barber ZW, Lemke ND, Oates CW, Ma LS, Stalnaker JE et al. Frequency evaluation of the doubly forbidden $^1S_0 \rightarrow {}^3P_0$ transition in bosonic ^{174}Yb. Phys Rev A. 2008;77:7–10. arXiv:0803.4503.
60. Chwalla M, Benhelm J, Kim K, Kirchmair G, Monz T, Riebe M et al. Absolute frequency measurement of the $^{40}Ca^+$ $4s^2S_{1/2} - 3d^2D_{5/2}$ clock transition. Phys Rev Lett. 2009;102:023002.
61. King SA, Godun RM, Webster SA, Margolis HS, Johnson LAM, Szymaniec K et al. Absolute frequency measurement of the $^2S_{1/2} \rightarrow {}^2F_{7/2}$ electric octupole transition in a single ion of $^{171}Yb^+$ with 10^{-15} fractional uncertainty. New J Phys. 2012;14:013045.
62. McFerran JJ, Yi L, Mejri S, Di Manno S, Zhang W, Guéna J et al. Neutral atom frequency reference in the deep ultraviolet with fractional uncertainty=5.7×10^{-15}. Phys Rev Lett. 2012;108:183004.
63. Targat RL, Lorini L, Coq YL, Zawada M, Guéna J, Abgrall M et al. Experimental realization of an optical second with strontium lattice clocks. Nat Commun. 2013;4:1–9.
64. Huntemann N, Lipphardt B, Tamm C, Gerginov V, Weyers S, Peik E. Improved limit on a temporal variation of m_p/m_e from comparisons of Yb^+ and Cs atomic clocks. Phys Rev Lett. 2014;113:1–5. arXiv:1407.4408.
65. McGrew WF, Zhang X, Leopardi H, Fasano RJ, Nicolodi D, Beloy K et al. Towards the optical second: verifying optical clocks at the SI limit. Optica. 2019;6:448–54.
66. Schwarz R, Dörscher S, Al-Masoudi A, Benkler E, Legero T, Sterr U et al. Long term measurement of the ^{87}Sr clock frequency at the limit of primary Cs clocks. Phys Rev Res. 2020;2:033242.
67. Weyers S, Gerginov V, Kazda M, Rahm J, Lipphardt B, Dobrev G et al. Advances in the accuracy, stability, and reliability of the PTB primary fountain clocks. Metrologia. 2018;55:789–805.
68. Rosenband T, Hume DBD, Schmidt PO, Chou CW, Brusch A, Lorini L et al. Frequency ratio of Al^+ and Hg^+ nn optical clocks; metrology at the 17th decimal place. Science. 2008;319:1808–12.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 50/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

69. Chou CW, Hume DB, Koelemeij JCJ, Wineland DJ, Rosenband T. Frequency comparison of two high-accuracy Al+ optical clocks. Phys Rev Lett. 2010;104:1.
70. Madej AA, Dubé P, Zhou Z, Bernard JE, Gertsvolf M. $^{88}Sr^{+}$ 445 THz single-ion reference at the 10^{-17} level via control and cancellation of systematic uncertainties and its measurement against the SI second. Phys Rev Lett. 2012;109:5.
71. Huntemann N, Okhapkin M, Lipphardt B, Weyers S, Tamm C, Peik E. High-accuracy optical clock based on the octupole transition in $^{171}Yb^{+}$. Phys Rev Lett. 2012;108:1.
72. Huntemann N. High-accuracy optical clock based on the octupole transition in Yb+171. Ph.D. thesis. 2014. arXiv:1111.2446. https://doi.org/10.1103/PhysRevLett.108.090801.
73. Barwood GP, Huang G, Klein HA, Johnson LAM, King SA, Margolis HS et al. Agreement between two $^{88}Sr^{+}$ optical clocks to 4 parts in 10^{17}. Phys Rev A. 2014;89:050501.
74. Nicholson T, Campbell S, Hutson R, Marti G, Bloom B, McNally R et al. Systematic evaluation of an atomic clock at 2×10^{-18} total uncertainty. Nat Commun. 2015;6:6896.
75. Ushijima I, Takamoto M, Das M, Ohkubo T, Katori H. Cryogenic optical lattice clocks. Nat Photonics. 2015;9:185–9.
76. Yamanaka K, Ohmae N, Ushijima I, Takamoto M, Katori H. Frequency ratio of ^{199}Hg and ^{87}Sr optical lattice clocks beyond the SI limit. Phys Rev Lett. 2015;114:1.
77. McGrew WF, Zhang X, Fasano RJ, Schäffer SA, Beloy K, Nicolodi D et al. Atomic clock performance enabling geodesy below the centimetre level. Nature. 2018;564:87–90.
78. Bothwell T, Kedar D, Oelker E, Robinson JM, Bromley SL, Tew WL et al. JILA SrI optical lattice clock with uncertainty of 2.0×10^{-18}. Metrologia. 2019;56:065004.
79. Takamoto M, Ushijima I, Ohmae N, Yahagi T, Kokado K, Shinkai H et al. Test of general relativity by a pair of transportable optical lattice clocks. Nat Photogr. 2020;14:411–5.
80. Kasevich M, Riis E, Chu S, DeVoe R. RF spectroscopy in an atomic fountain. Phys Rev Lett. 1989;63:612–5.
81. Clairon A, Salomon C, Guellati S, Phillips W. Ramsey resonance in a Zacharias fountain. Europhys Lett. 1991;16:165.
82. Jefferts S, Shirley J, Parker T, Heavner T, Meekhof D, Nelson C et al. Accuracy evaluation of NIST-F1. Metrologia. 2002;39:321.
83. Cao J, Zhang P, Shang J, Cui K, Yuan J, Chao S et al. A compact, transportable single-ion optical clock with 7.8×10^{-17} systematic uncertainty. Appl Phys B. 2017;123:112.
84. Yamaguchi A, Safronova M, Gibble K, Katori H. Narrow-line cooling and determination of the magic wavelength of Cd. Phys Rev Lett. 2019;123:113201.
85. Bothwell T, Kennedy CJ, Aeppli A, Kedar D, Robinson JM, Oelker E, et al. Resolving the gravitational redshift within a millimeter atomic sample. arXiv:2109.12238.
86. Kennedy CJ, Oelker E, Robinson JM, Bothwell T, Kedar D, Milner WR et al. Precision metrology meets cosmology: improved constraints on ultralight dark matter from atom-cavity frequency comparisons. Phys Rev Lett. 2020;125:201302.
87. Godun RM, Nisbet-Jones PBR, Jones JM, King SA, Johnson LAM, Margolis HS et al. Frequency ratio of two optical clock transitions in $^{171}Yb^{+}$ and constraints on the time variation of fundamental constants. Phys Rev Lett. 2014;113:210801. arXiv:1407.0164.
88. Giorgetta FR, Swann WC, Sinclair LC, Baumann E, Coddington I, Newbury NR. Optical two-way time and frequency transfer over free space. Nat Photonics. 2013;7:434–8.
89. Yamaguchi A, Fujieda M, Kumagai M, Hachisu H, Nagano S, Li Y et al. Direct comparison of distant optical lattice clocks at the 10^{-16} uncertainty. Appl Phys Express. 2011;4:082203.
90. Beloy K, Bodine MI, Bothwell T, Brewer SM, Bromley SL, Chen J-S et al. Frequency ratio measurements at 18-digit accuracy using an optical clock network. Nature. 2021;591:564–9.
91. Roberts BM, Delva P, Al-Masoudi A, Amy-Klein A, Bærentsen C, Baynham CFA et al. Search for transient variations of the fine structure constant and dark matter using fiber-linked optical atomic clocks. New J Phys. 2020;22:093010. arXiv:1907.02661.
92. Bober M, Morzyński P, Cygan A, Lisak D, Masłowski P, Prymaczek M et al. Strontium optical lattice clocks for practical realization of the metre and secondary representation of the second. Meas Sci Technol. 2015;26:075201.
93. Bongs K, Singh Y, Smith L, He W, Kock O, Świerad D et al. Development of a strontium optical lattice clock for the SOC mission on the ISS. C R Phys. 2015;16:553–64. arXiv:1503.08457v1.
94. Falke S, Lemke N, Grebing C, Lipphardt B, Weyers S, Gerginov V, et al. A strontium lattice clock with 3×10^{-17} inaccuracy and its frequency. New J Phys. 2014;16. arXiv:1312.3419.
95. Lin Y-G, Wang Q, Li Y, Meng F, Lin B-K, Zang E-J et al. First evaluation and frequency measurement of the strontium optical lattice clock at NIM. Chin Phys Lett. 2015;32:90601.
96. Ludlow A, Boyd M, Zelevinsky T, Foreman S, Blatt S, Notcutt M et al. Systematic study of the ^{87}Sr clock transition in an optical lattice. Phys Rev Lett. 2006;96:033003.
97. Poli N, Schioppo M, Vogt S, Falke S, Sterr U, Lisdat C et al. A transportable strontium optical lattice clock. Appl Phys B. 2014;117:1107–16.
98. Strelkin SA, Galyshev AA, Berdasov OI, Gribov AY, Sutyrin DV, Khabarova KY et al. Narrow line cooling of 88Sr atoms in the magneto-optical trap for precision frequency standard. Phys Proc. 2015;72:184–8.
99. Takamoto M, Hong F-L, Higashi R, Katori H. An optical lattice clock. Nature. 2005;435:321–4.
100. Hobson R, Bowden W, Vianello A, Silva A, Baynham CFA, Margolis HS et al. A strontium optical lattice clock with 1×10^{-17} uncertainty and measurement of its absolute frequency. Metrologia. 2020;57:065026.
101. Zheng X, Dolde J, Lochab V, Merriman BN, Li H, Kolkowitz S. High precision differential clock comparisons with a multiplexed optical lattice clock. arXiv:2109.12237.
102. Akamatsu D, Yasuda M, Inaba H, Hosaka K, Tanabe T, Onae A et al. Frequency ratio measurement of ^{171}Yb and ^{87}Sr optical lattice clocks. Opt Express. 2014;22:7898–905.
103. Barber ZW, Hoyt CW, Oates CW, Hollberg L, Taichenachev AV, Yudin VI. Direct excitation of the forbidden clock transition in neutral ^{174}Yb atoms confined to an optical lattice. Phys Rev Lett. 2006;96:083002.
104. Pedrozo-Peñafiel E, Colombo S, Shu C, Adiyatullin AF, Li Z, Mendez E et al. Entanglement on an optical atomic-clock transition. Nature. 2020;588:414–8.
105. Gao Q, Zhou M, Han C, Li S, Zhang S, Yao Y et al. Systematic evaluation of a ^{171}Yb optical clock by synchronous comparison between two lattice systems. Sci Rep. 2018;8:8022.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 51/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

106. Kim H, Heo M-S, Park CY, Yu D-H, Lee W-K. Absolute frequency measurement of the ^{171}Yb optical lattice clock at KRISS using TAI for over a year. Metrologia. 2021;58:055007.
107. Kersten P, Mensing F, Sterr U, Riehle F. A transportable optical calcium frequency standard. Appl Phys B. 1999;68:27–38.
108. Olson J, Fox RW, Fortier TM, Sheerin TF, Brown RC, Leopardi H et al. Ramsey–Bordé matter-wave interferometry for laser frequency stabilization at 10^{-16} frequency instability and below. Phys Rev Lett. 2019;123:073202.
109. Döringshoff K, Gutsch FB, Schkolnik V, Kürbis C, Oswald M, Pröbster B et al. Iodine frequency reference on a sounding rocket. Phys Rev Appl. 2019;11:054068.
110. Argence B, Prevost E, Lévèque T, Goff RL, Bize S, Lemonde P et al. Prototype of an ultra-stable optical cavity for space applications. Opt Express. 2012;20:25409–20.
111. Chen Q-F, Nevsky A, Cardace M, Schiller S, Legero T, Häfner S et al. A compact, robust, and transportable ultra-stable laser with a fractional frequency instability of 1×10^{-15}. Rev Sci Instrum. 2014;85:113107.
112. Häfner S, Herbers S, Vogt S, Lisdat C, Sterr U. Transportable interrogation laser system with an instability of $\mathbf{mod}\,\sigma_y = 3 \times 10^{-16}$. Opt Express. 2020;28:16407–16.
113. Webster S, Gill P. Force-insensitive optical cavity. Opt Lett. 2011;36:3572. arXiv:1108.4819.
114. Sanjuan J, Abich K, Gohlke M, Resch A, Schuldt T, Wegehaupt T et al. Long-term stable optical cavity for special relativity tests in space. Opt Express. 2019;27:36206–20.
115. Hill IR, Hendricks RJ, Donnellan S, Gaynor P, Allen B, Barwood GP et al. Dual-axis cubic cavity for drift-compensated multi-wavelength laser stabilisation. Opt Express. 2021;29:36758–68.
116. Origlia S, Pramod MS, Schiller S, Singh Y, Bongs K, Schwarz R, et al. A high-performance optical lattice clock based on bosonic atoms. arXiv:1803.03157.
117. Koller SB, Grotti J, Vogt S, Al-Masoudi A, Dörscher S, Häfner S et al. Transportable optical lattice clock with 7×10^{-17} uncertainty. Phys Rev Lett. 2017;118:073601.
118. Ohmae N, Takamoto M, Takahashi Y, Kokubun M, Araki K, Hinton A, et al. Transportable strontium optical lattice clocks operated outside laboratory at the level of 10^{-18} uncertainty. Adv Quantum Technol. 2021;2100015.
119. Grotti J, Koller S, Vogt S, Häfner S, Sterr U, Lisdat C et al. Geodesy and metrology with a transportable optical clock. Nat Phys. 2018;14:437–41.
120. Huang Y, Zhang H, Zhang B, Hao Y, Guan H, Zeng M et al. Geopotential measurement with a robust, transportable Ca^+ optical clock. Phys Rev A. 2020;102:050802.
121. Bongs K, Singh Y, Smith L, He W, Kock O, Swierad D et al. Development of a strontium optical lattice clock for the SOC mission on the ISS. CRP. 2015;16:553–64. arXiv:1503.08457 [physics.atom-ph].
122. Derevianko A, Gibble K, Hollberg L, Newbury NR, Oates C, Safronova MS, et al. Fundamental Physics with a State-of-the-Art Optical Clock in Space. arXiv:2112.10817.
123. Petit G. Sub-10–16 accuracy GNSS frequency transfer with IPPP. GPS Solut. 2021;25:1–9. https://doi.org/10.1007/s10291-020-01062-2.
124. Kirchner D, Ressler H, Grudler P, Baumont F, Veillet C, Lewandowski W et al. Comparison of GPS common-view and two-way satellite time transfer over a baseline of 800 km. Metrologia. 1993;30:183.
125. ITU technical recommendation: the operational use of two-way satellite time and frequency transfer employing pseudorandom noise codes, ITU-R TF. 1153-4 (2015).
126. Huang Y-J, Fujieda M, Takiguchi H, Tseng W-H, Tsao H-W. Stability improvement of an operational two-way satellite time and frequency transfer system. Metrologia. 2016;53:881.
127. Droste S, Grebing C, Leute J, Raupach SMF, Matveev A, Hänsch TW et al. Characterization of a 450 km baseline GPS carrier-phase link using an optical fiber link. New J Phys. 2015;17:083044.
128. Schäfer W, Feldmann T. Perspectives of time and frequency transfer via satellite. J Phys Conf Ser. 2016;723:012038.
129. Samain E, Vrancken P, Guillemot P, Fridelance P, Exertier P. Time transfer by laser link (T2L2): characterization and calibration of the flight instrument. Metrologia. 2014;51:503–15.
130. Schreiber K, Prochazka I, Lauber P, Hugentobler U, Schäfer W, Cacciapuoti L et al. Ground-based demonstration of the European Laser Timing (ELT) experiment. IEEE Trans Ultrason Ferroelectr Freq Control. 2010;57:728–37.
131. Djerroud K, Acef O, Clairon A, Lemonde P, Man CN, Samain E et al. Coherent optical link through the turbulent atmosphere. Opt Lett. 2010;35:1479–81.
132. Dix-Matthews BP, Schediwy SW, Gozzard DR, Savalle E, Esnault F-X, Lévèque T et al. Point-to-point stabilized optical frequency transfer with active optics. Nat Commun. 2021;12:515.
133. Sinclair LC, Bergeron H, Swann WC, Baumann E, Deschênes J-D, Newbury NR. Comparing optical oscillators across the air to milliradians in phase and 10^{-17} in frequency. Phys Rev Lett. 2018;120:050801.
134. Bodine MI, Ellis JL, Swann WC, Stevenson SA, Deschênes J-D, Hannah ED et al. Optical time-frequency transfer across a free-space, three-node network. APL Photon. 2020;5:076113. https://doi.org/10.1063/5.0010704.
135. Bergeron H, Sinclair LC, Swann WC, Khader I, Cossel KC, Cermak M et al. Femtosecond synchronization of optical clocks off of a flying quadcopter. Nat Commun. 2019;10:1819.
136. Shen Q, Guan J-Y, Zeng T, Lu Q-M, Huang L, Cao Y et al. Experimental simulation of time and frequency transfer via an optical satellite–ground link at 10–18 instability. Optica. 2021;8:471–6.
137. Glaser S, et al. Reference system origin and scale realization within the future GNSS constellation "Kepler". J Geod. 2020;94.
138. Liu L, Lü D-S, Chen W-B, Li T, Qu Q-Z, Wang B et al. In-orbit operation of an atomic clock based on laser-cooled ^{87}Rb atoms. Nat Commun. 2018;9:2760.
139. Burt EA, Prestage JD, Tjoelker RL, Enzer DG, Kuang D, Murphy DW et al. Demonstration of a trapped-ion atomic clock in space. Nature. 2021;595:43–7.
140. Barwood GP, Gill P, Klein HA, Hosaka K, Huang G, Lea SN et al. Trapped strontium ion optical clock. In: Armandillo E, Costeraste J, Karafolas N, editors. International conference on space optics—ICSO 2006. vol. 10567. International Society for Optics and Photonics, SPIE; 2017. p. 681–7. https://doi.org/10.1117/12.2308177.
141. Prochazka I, Blazej J, Kodet J. Note: space qualified solid state photon counting detector with reduced detection delay temperature drift. Rev Sci Instrum. 2018;89:056106.
142. Prochazka I, Blazej J, Flekova T, Kodet J. Silicon based photon counting detector providing femtosecond detection delay stability. IEEE J Sel Top Quantum Electron. 2020;26:1–5.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 52/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

143. The State Council Information Office of the People's Republic of China. "China's Space Program: a 2021 Perspective." http://www.cnsa.gov.cn/english/n6465645/n6465648/c6813088/content.html (2021).
144. Steigenberger P, Montenbruck O. Galileo status: orbits, clocks, and positioning. GPS Solut. 2017;21:319–31.
145. Vannicola F, Beard R, White J, Senior K, Largay M, Buisson J. GPS block IIF atomic frequency standard analysis. In: Proceedings of the 42nd annual precise time and time interval systems and applications meeting. Reston, Virginia, November 2010. 2010. p. 181–96.
146. Shen Q, Lin H, Deng J, Wang Y. Pulsed optically pumped atomic clock with a medium- to long-term frequency stability of 10^{-15}. Rev Sci Instrum. 2020;91:045114.
147. Phelps G, Lemke N, Erickson C, Burke J, Martin K. Compact optical clock with 5×10^{-13} instability at 1 s. Navigation. 2018;65:49–54.
148. Schkolnik V, Döringshoff K, Gutsch FB, Oswald M, Schuldt T, Braxmaier C et al. Jokarus—design of a compact optical iodine frequency reference for a sounding rocket mission. EPJ Quantum Technol. 2017;4:9.
149. DLR. Compasso. https://www.dlr.de/gk/en/desktopdefault.aspx/tabid-14174/28130_read-59743.
150. US National Academies Decadal Survey on Biological and Physical Sciences Research in Space 2023–2032. https://www.nationalacademies.org/our-work/decadal-survey-on-life-and-physical-sciences-research-in-space-2023-2032.
151. WMO. The Status of the Global Climate Observing System 2021: The GCOS Status Report (GCOS-240). https://gcos.wmo.int/en/gcos-status-report-2021.
152. IUGG. Resolution on future satellite gravity and magnetic mission constellations. https://www.iugg.org/assemblies/2015prague/2015_Prague_Comptes_Rendus_Part1.pdf (2015).
153. Roland P, editor. Observing Mass Transport to Understand Global Change and to benefit Society: Science and User Needs—An international multi-disciplinary initiative for IUGG. https://dgk.badw.de/fileadmin/user_upload/Files/DGK/docs/b-320.pdf (2015).
154. Pail R, Bingham R, Braitenberg C, Dobslaw H, Eicker A, Güntner A et al. Science and user needs for observing global mass transport to understand global change and to benefit society. Surv Geophys. 2015;36:743–72.
155. dos Santos FP, et al. International Association of Geodesy Project—Novel Sensors and Quantum Technology for Geodesy (QuGe) WG Q.1: Quantum gravimetry in space and on ground. https://quge.iag-aig.org/, https://quge.iag-aig.org/quge-working-groups/230.
156. Petit G, et al. International Association of Geodesy Project—Novel Sensors and Quantum Technology for Geodesy (QuGe) WG Q.3: Relativistic geodesy with clocks. https://quge.iag-aig.org/, https://quge.iag-aig.org/quge-working-groups/232.
157. Haagmans R, Siemes C, Massotti L, Carraz O, Silvestrin P. ESA's next-generation gravity mission concepts. Rend Lincei, Sci Fis Nat. 2020;1–11.
158. NASA. Mass Change. https://science.nasa.gov/earth-science/decadal-mc.
159. Bender P, Wiese D, Nerem R. A possible dual-GRACE mission with 90 degree and 63 degree inclination orbits. In: Proceedings of the third international symposium on formation flying, missions and technologies. ESA/ESTEC Noordwijk. 2008. p. 1–6.
160. Pail R. Impact of new measurement technologies for the monitoring of mass transport processes in the Earth system; "Quantum Gravimetry in Space and on Ground" Workshop, 26–27 May 2021. https://quge.iag-aig.org/, https://quge.iag-aig.org/quge-meetings/workshop-wgq1.
161. Carraz O, Siemes C, Massotti L, Haagmans R, Silvestrin P. A spaceborne gravity gradiometer concept based on cold atom interferometers for measuring Earth's gravity field. Microgravity Sci Technol. 2014;26:139–45. arXiv:1406.0765.
162. Trimeche A, Battelier B, Becker D, Bertoldi A, Bouyer P, Braxmaier C et al. Concept study and preliminary design of a cold atom interferometer for space gravity gradiometry. Class Quantum Gravity. 2019;36:215004.
163. Geiger R, Landragin A, Merlet S, Pereira Dos Santos F. High-accuracy inertial measurements with cold-atom sensors. AVS Quantum Sci. 2020;2:024702. arXiv:2003.12516.
164. Karcher R, Imanaliev A, Merlet S, Santos FPD. Improving the accuracy of atom interferometers with ultracold sources. New J Phys. 2018;20:113041.
165. Chiow S-W, Williams J, Yu N. Noise reduction in differential phase extraction of dual atom interferometers using an active servo loop. Phys Rev A. 2016;93:013602.
166. Asenbaum P, Overstreet C, Kovachy T, Brown DD, Hogan JM, Kasevich MA. Phase shift in atom interferometry due to spacetime curvature. Phys Rev Lett. 2016;118:183602. arXiv:1610.03832.
167. Berg P, Abend S, Tackmann G, Schubert C, Giese E, Schleich WP et al. Composite-light-pulse technique for high-precision atom interferometry. Phys Rev Lett. 2015;114:063002.
168. Savoie D, Altorio M, Fang B, Sidorenkov LA, Geiger R, Landragin A. Interleaved atom interferometry for high-sensitivity inertial measurements. Sci Adv. 2018;4:eaau7948.
169. Douch K, Wu H, Schubert C, Müller J, dos Santos FP. Simulation-based evaluation of a cold atom interferometry gradiometer concept for gravity field recovery. Adv Space Res. 2018;61:1307–23.
170. Bidel Y, Carraz O, Charrière R, Cadoret M, Zahzam N, Bresson A. Compact cold atom gravimeter for field applications. Appl Phys Lett. 2013;102:144107. arXiv:1302.1518.
171. Bonnin A, Diboune C, Zahzam N, Bidel Y, Cadoret M, Bresson A. New concepts of inertial measurements with multi-species atom interferometry. Appl Phys B, Lasers Opt. 2018;124:181. arXiv:1710.06289.
172. Abrykosov P, Pail R, Gruber T, Zahzam N, Bresson A, Hardy E et al. Impact of a novel hybrid accelerometer on satellite gravimetry performance. Adv Space Res. 2019;63:3235–48.
173. Müller J, Wu H. Using quantum optical sensors for determining the Earth's gravity field from space. J Geod. 2020;94:1–14.
174. Lévèque T, Fallet C, Mandea M, Biancale R, Lemoine JM, Tardivel S et al. Gravity field mapping using laser-coupled quantum accelerometers in space. J Geod. 2021;95:15.
175. Müller J, Dirkx D, Kopeikin SM, Lion G, Panet I, Petit G et al. High performance clocks and gravity field determination. Space Sci Rev. 2018;214:1–31.
176. Domps F. Quantum Space Gravimetry at European Commission. https://indico.cern.ch/event/1064855/contributions/4524210/attachments/2315397/3941302/QSG%40EC|_CERN.pdf.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 53/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

177. Strategic autonomy in developing, deploying and using global space-based infrastructures, services, applications and data 2021 (HORIZON-CL4-2021-SPACE-01). https://ec.europa.eu/info/funding-tenders/opportunities/portal/screen/opportunities/topic-details/horizon-cl4-2021-space-01-62.
178. Visser P, Bettadpur S, Chambers D, Diament M, Gruber T, Hanna E, et al. Towards a sustained observing system for mass transport to understand global change and to benefit society. 2017. p. 32–33. http://www.swarm2017.org/.
179. Hensel T, Loriani S, Schubert C, Fitzek F, Abend S, Ahlers H et al. Inertial sensing with quantum gases: a comparative performance study of condensed versus thermal sources for atom interferometry. Eur Phys J D. 2021;75:108.
180. Abe M, Adamson P, Borcean M, Bortoletto D, Bridges K, Carman SP, et al. Matter-wave Atomic Gradiometer Interferometric Sensor (MAGIS-100). arXiv:2104.02835v1.
181. Elliott ER, Krutzik MC, Williams JR, Thompson RJ, Aveline DC. NASA's Cold Atom Lab (CAL): system development and ground test status. npj Microgravity. 2016;4:16.
182. Aveline DC, Williams JR, Elliott ER, Dutenhoffer C, Kellogg JR, Kohel JM et al. Observation of Bose–Einstein condensates in an Earth-orbiting research lab. Nature. 2020;582:193–7.
183. Canuel B et al. ELGAR—a European laboratory for gravitation and atom-interferometric research. Class Quantum Gravity. 2020;37:225017. arXiv:1911.03701.
184. Badurina L et al. AION: an atom interferometer observatory and network. JCAP. 2020;05:011. arXiv:1911.11755.
185. Bertoldi A, Bouyer P, Canuel B. Quantum sensors with matter waves for GW observation. arXiv:2010.14604.
186. Canuel B et al. Exploring gravity with the MIGA large scale atom interferometer. Sci Rep. 2018;8:14064. arXiv:1703.02490.
187. Schlippert D, Meiners C, Rengelink R, Schubert C, Tell D, Wodey É et al. Matter-wave interferometry for inertial sensing and tests of fundamental physics. In: CPT and Lorentz symmetry: proceedings of the eighth meeting on CPT and Lorentz symmetry. Singapore: World Scientific; 2020. p. 37–40.
188. Becker D, Lachmann MD, Seidel ST, Ahlers H, Dinkelaker AN, Grosse J et al. Space-borne Bose–Einstein condensation for precision interferometry. Nature. 2018;562:391–5.
189. Lachmann MD, Ahlers H, Becker D, Dinkelaker AN, Grosse J, Hellmig O et al. Ultracold atom interferometry in space. Nat Commun. 2021;12:1317.
190. Schubert C, Hartwig J, Ahlers H, Posso-Trujillo K, Gaaloul N, Velte U, et al. Differential atom interferometry with ^{87}Rb and ^{85}Rb for testing the UFF in STE-QUEST. arXiv:1312.5963.
191. Aguilera D et al. STE-QUEST–test of the universality of free fall using cold atom interferometry. Class Quantum Gravity. 2014;31:115010. arXiv:1312.5980.
192. Touboul P, Métris G, Rodrigues M, André Y, Baghi Q, Bergé J et al. MICROSCOPE. Phys Rev Lett. 2017;119:231101.
193. Berge J, Brax P, Metris G, Pernot-Borras M, Touboul P, Uzan J-P. MICROSCOPE mission: first constraints on the violation of the weak equivalence principle by a light scalar dilaton. Phys Rev Lett. 2018;120:141101. arXiv:1712.00483.
194. Zhan M-S et al. ZAIGA: Zhaoshan long-baseline atom interferometer gravitation antenna. Int J Mod Phys D. 2019;28:1940005. arXiv:1903.09288.
195. Yin J, Cao Y, Li Y-H, Liao S-K, Zhang L, Ren J-G et al. Satellite-based entanglement distribution over 1200 kilometers. Science. 2017;356:1140–4.
196. Loriani S, Schubert C, Schlippert D, Ertmer W, Pereira Dos Santos F, Rasel EM et al. Resolution of the colocation problem in satellite quantum tests of the universality of free fall. Phys Rev D. 2020;102:124043.
197. Roura A. Circumventing Heisenberg's uncertainty principle in atom interferometry tests of the equivalence principle. Phys Rev Lett. 2017;118:160401.
198. Overstreet C, Asenbaum P, Kovachy T, Notermans R, Hogan JM, Kasevich MA. Effective inertial frame in an atom interferometric test of the equivalence principle. Phys Rev Lett. 2018;120:183604.
199. Tino GM et al. SAGE: a proposal for a space atomic gravity explorer. Eur Phys J D. 2019;73:228. arXiv:1907.03867.
200. Dimopoulos S, Graham PW, Hogan JM, Kasevich MA, Rajendran S. An atomic gravitational wave interferometric sensor (AGIS). Phys Rev D. 2008;78:122002. arXiv:0806.2125.
201. Yu N. Opportunities of Cold Atom Quantum Sensors under Microgravity and in Space—With focus on direct detection of dark energy. https://indico.cern.ch/event/1064855/contributions/4524214/attachments/2321220/3952720/ECWCAP2021_nyu_UR.pdf.
202. LIGO Scientific collaboration, Abbott BP et al Exploring the sensitivity of next generation gravitational wave detectors. Class Quantum Gravity. 2017;34:044001. arXiv:1607.08697.
203. VIRGO collaboration, Acernese F et al Advanced virgo: a second-generation interferometric gravitational wave detector. Class Quantum Gravity. 2015;32:024001. arXiv:1408.3978.
204. KAGRA collaboration, Akutsu T, et al. Overview of KAGRA: KAGRA science. arXiv:2008.02921.
205. Punturo M et al. The Einstein telescope: a third-generation gravitational wave observatory. Class Quantum Gravity. 2010;27:194002.
206. Reitze D et al. Cosmic explorer: the U.S. contribution to gravitational-wave astronomy beyond LIGO. Bull Am Astron Soc. 2019;51:035. arXiv:1907.04833.
207. LISA collaboration, Audley H, et al. Laser Interferometer Space Antenna. arXiv:1702.00786.
208. Luo J, Chen L-S, Duan H-Z, Gong Y-G, Hu S, Ji J et al. TianQin: a space-borne gravitational wave detector. Class Quantum Gravity. 2016;33:035010.
209. Ruan W-H, Guo Z-K, Cai R-G, Zhang Y-Z. Taiji program: gravitational-wave sources. Int J Mod Phys A. 2020;35:2050075. arXiv:1807.09495.
210. MAGIS collaboration, Graham PW, Hogan JM, Kasevich MA, Rajendran S, Romani RW. Mid-band gravitational wave detection with precision atomic sensors. arXiv:1711.02225.
211. Brax P, Casas S, Desmond H, Elder B. Testing screened modified gravity. Universe. 2021;8:11. arXiv:2201.10817.
212. Hamilton P, Jaffe M, Haslinger P, Simmons Q, Müller H, Khoury J. Atom-interferometry constraints on dark energy. Science. 2015;349:849–51. arXiv:1502.03888.
213. Elder B, Khoury J, Haslinger P, Jaffe M, Müller H, Hamilton P. Chameleon dark energy and atom interferometry. Phys Rev D. 2016;94:044051. arXiv:1603.06587.

UNCORRECTED PROOF

AUTHOR'S PROOF

214. Burrage C, Sakstein J. A compendium of chameleon constraints. JCAP. 2016;11:045. arXiv:1609.01192.
215. Jaffe M, Haslinger P, Xu V, Hamilton P, Upadhye A, Elder B et al. Testing sub-gravitational forces on atoms from a miniature, in-vacuum source mass. Nat Phys. 2017;13:938 arXiv:1612.05171.
216. Sabulsky DO, Dutta I, Hinds EA, Elder B, Burrage C, Copeland EJ. Experiment to detect dark energy forces using atom interferometry. Phys Rev Lett. 2019;123:061102. arXiv:1812.08244.
217. Bars HP-L, Guerlin C, Hees A, Peaucelle R, Tasson JD, Bailey QG et al. New test of Lorentz invariance using the MICROSCOPE space mission. Phys Rev Lett. 2019;123:231102.
218. Schlamminger S, Choi KY, Wagner TA, Gundlach JH, Adelberger EG. Test of the equivalence principle using a rotating torsion balance. Phys Rev Lett. 2008;100:041101. arXiv:0712.0607.
219. Kostelecky VA, Tasson JD. Matter-gravity couplings and Lorentz violation. Phys Rev D. 2011;83:016013.
220. Hees A, Minazzoli O, Savalle E, Stadnik YV, Wolf P. Violation of the equivalence principle from light scalar dark matter. Phys Rev D. 2018;98:064051. arXiv:1807.04512.
221. Bonnin A, Zahzam N, Bidel Y, Bresson A. Characterization of a simultaneous dual-species atom interferometer for a quantum test of the weak equivalence principle. Phys Rev A. 2015;92:023626. arXiv:1506.06535.
222. Biskupek L, Müller J, Torre J-M. Benefit of new high-precision LLR data for the determination of relativistic parameters. Universe. 2021;7:34.
223. Peters A, Chung KY, Chu S. High-precision gravity measurements using atom interferometry. Metrologia. 2001;38:25.
224. Merlet S, Bodart Q, Malossi N, Landragin A, Santos FPD, Gitlein O et al. Comparison between two mobile absolute gravimeters: optical versus atomic interferometers. Metrologia. 2010;47:L9–11.
225. Schlippert D, Hartwig J, Albers H, Richardson L, Schubert LC, Roura A et al. Quantum test of the universality of free fall. Phys Rev Lett. 2014;112:203002.
226. Tarallo MG, Mazzoni T, Poli N, Sutyrin DV, Zhang X, Tino GM. Test of Einstein equivalence principle for 0-spin and half-integer-spin atoms: search for spin-gravity coupling effects. Phys Rev Lett. 2014;113:023005.
227. Zhou L, Long S, Tang B, Chen X, Gao F, Peng W et al. Test of equivalence principle at 10^{-8} level by a dual-species double-diffraction Raman atom interferometer. Phys Rev Lett. 2015;115:013004.
228. Asenbaum P, Overstreet C, Kim M, Curti J, Kasevich MA. Atom-interferometric test of the equivalence principle at the 10^{-12} level. Phys Rev Lett. 2020;125:191101.
229. Hartwig J, Abend S, Schubert C, Schlippert D, Ahlers H, Posso-Trujillo K et al. Testing the universality of free fall with rubidium and ytterbium in a very large baseline atom interferometer. New J Phys. 2015;17:035011.
230. Perez P, Sacquin Y. The GBAR experiment: gravitational behaviour of antihydrogen at rest. https://doi.org/10.1088/0264-9381/29/18/184008.
231. Stadnik YV. New bounds on macroscopic scalar-field topological defects from non-transient signatures due to environmental dependence and spatial variations of the fundamental constants. Phys Rev D. 2020;102:115016. arXiv:2006.00185.
232. Geraci AA, Derevianko A. Sensitivity of atom interferometry to ultralight scalar field dark matter. Phys Rev Lett. 2016;117:261301.
233. Figueroa NL, Budker D, Rasel EM. Dark matter searches using accelerometer-based networks. Quantum Sci Technol. 2021;6:034004.
234. Antypas D, Tretiak O, Zhang K, Garcon A, Perez G, Kozlov MG et al. Probing fast oscillating scalar dark matter with atoms and molecules. Quantum Sci Technol. 2021;6:034001.
235. Oswald R, et al. Search for oscillations of fundamental constants using molecular spectroscopy. arXiv:2111.06883.
236. Ellis J, Vaskonen V. Probes of gravitational waves with atom interferometers. Phys Rev D. 2020;101:124013. arXiv:2003.13480.
237. Ashby N, Bender PL, Hall JL, Ye J, Diddams SA, Jefferts SR et al. Measurement of gravitational time delay using drag-free spacecraft and an optical clock. In: Klioner SA, Seidelman PK, Soffel MH, editors. Relativity in fundamental astronomy proceedings IAU symposium. Vol. 261. 2010. p. 414.
238. Mohageg M, et al. The Deep Space Quantum Link: Prospective Fundamental Physics Experiments using Long-Baseline Quantum Optics. arXiv:2111.15591.
239. Bassi A, Ghirardi G. Dynamical reduction models. Phys Rep. 2003;379:257–426.
240. Bassi A, Lochan K, Satin S, Singh TP, Ulbricht H. Models of wave-function collapse, underlying theories, and experimental tests. Rev Mod Phys. 2013;85:471–527.
241. Bilardello M, Donadi S, Vinante A, Bassi A. Bounds on collapse models from cold-atom experiments. Phys A, Stat Mech Appl. 2016;462:764–82.
242. Zych M, Brukner Č. Quantum formulation of the Einstein equivalence principle. Nat Phys. 2018;14:1027–31.
243. Rosi G, D'Amico G, Cacciapuoti L, Sorrentino F, Prevedelli M, Zych M et al. Quantum test of the equivalence principle for atoms in coherent superposition of internal energy states. Nat Commun. 2017;8:15529.
244. Geiger R, Trupke M. Proposal for a quantum test of the weak equivalence principle with entangled atomic species. Phys Rev Lett. 2018;120:043602.
245. Roura A. Gravitational redshift in quantum-clock interferometry. Phys Rev X. 2020;10:021014.
246. Inouye S, Andrews M, Stenger J, Miesner H-J, Stamper-Kurn DM, Ketterle W. Observation of Feshbach resonances in a Bose–Einstein condensate. Nature. 1998;392:151–4.
247. Pasquini T, Saba M, Jo G-B, Shin Y, Ketterle W, Pritchard DE et al. Low velocity quantum reflection of Bose–Einstein condensates. Phys Rev Lett. 2006;97:093201.
248. Lundblad N, Carollo R, Lannert C, Gold M, Jiang X, Paseltiner D et al. Shell potentials for microgravity Bose–Einstein condensates. npj Microgravity. 2019;5:1–6.
249. Cabrera C, Tanzi L, Sanz J, Naylor B, Thomas P, Cheiney P et al. Quantum liquid droplets in a mixture of Bose–Einstein condensates. Science. 2018;359:301–4.
250. LIGO Scientific, Virgo, KAGRA collaboration, Abbott R et al Constraints on cosmic strings using data from the third advanced LIGO–virgo observing run. Phys Rev Lett. 2021;126:241102. arXiv:2101.12248.
251. Badurina L, Buchmueller O, Ellis J, Lewicki M, McCabe C, Vaskonen V. Prospective sensitivities of atom interferometers to gravitational waves and ultralight dark matter. Philos Trans R Soc Lond A. 2022;380:20210060. arXiv:2108.02468.
252. Ellis J, Lewicki M. Cosmic string interpretation of NANOGrav pulsar timing data. Phys Rev Lett. 2021;126:041304. arXiv:2009.06555.

UNCORRECTED PROOF

«EPJQT 40507 class: bmc-onecol-v1.cls 2022/10/25 v1.0.1 BMC onecolumn layout file: epjqt147.tex (Ausra) Prn:26/10/2022; 12:04 p. 55/55»
« doctopic: ReviewPaper numbering style: ContentOnly reference style: vancouver»

AUTHOR'S PROOF

253. Blasi S, Brdar V, Schmitz K. Has NANOGrav found first evidence for cosmic strings? Phys Rev Lett. 2021;126:041305. arXiv:2009.06607.
254. NANOGrav collaboration, Arzoumanian Z et al The NANOGrav 12.5 yr data set: search for an isotropic stochastic gravitational-wave background. Astrophys J Lett. 2020;905:L34. arXiv:2009.04496.
255. Goncharov B, et al. On the evidence for a common-spectrum process in the search for the nanohertz gravitational-wave background with the Parkes Pulsar Timing Array. arXiv:2107.12112.
256. Chen S et al. Common-red-signal analysis with 24-yr high-precision timing of the European pulsar timing array: inferences in the stochastic gravitational-wave background search. Mon Not R Astron Soc. 2021;508:4970–93. arXiv:2110.13184.
257. Antoniadis J, et al. The International Pulsar Timing Array second data release: Search for an isotropic Gravitational Wave Background. arXiv:2201.03980.
258. Campeti P, Komatsu E, Poletti D, Baccigalupi C. Measuring the spectrum of primordial gravitational waves with CMB, PTA and laser interferometers. JCAP. 2021;01:012. arXiv:2007.04241.
259. Ellis J, Lewicki M, Vaskonen V. Updated predictions for gravitational waves produced in a strongly supercooled phase transition. JCAP. 2020;11:020. arXiv:2007.15586.
260. Fedderke MA, Graham PW, Rajendran S. Asteroids for microhertz gravitational-wave detection. arXiv:2112.11431.
261. Mukhopadhyay M, Cardona C, Lunardini C. The neutrino gravitational memory from a core collapse supernova: phenomenology and physics potential. arXiv:2105.05862.
262. Event Horizon Telescope collaboration, Akiyama K et al First M87 event horizon telescope results. I. The shadow of the supermassive black hole. Astrophys J Lett. 2019;875:L1. arXiv:1906.11238.
263. Schuldt T, Döringshoff K, Kovalchuk EV, Keetman A, Pahl J, Peters A et al. Development of a compact optical absolute frequency reference for space with 10^{-15} instability. Appl Opt. 2017;56:1101–6.
264. Gürlebeck N, Wörner L, Schuldt T, Döringshoff K, Gaul K, Gerardi D et al. BOOST: a satellite mission to test Lorentz invariance using high-performance optical frequency references. Phys Rev D. 2018;97:124051.
265. Ludlow AD, Huang X, Notcutt M, Zanon-Willette T, Foreman SM, Boyd MM et al. Compact, thermal-noise-limited optical cavity for diode laser stabilization at 1×10^{-15}. Opt Lett. 2007;32:641–3.
266. Seel S, Storz R, Ruoso G, Mlynek J, Schiller S. Cryogenic optical resonators: a new tool for laser frequency stabilization at the 1 Hz level. Phys Rev Lett. 1997;78:4741–4.
267. Tobar ME, Wolf P, Bize S, Santarelli G, Flambaum V. Testing local Lorentz and position invariance and variation of fundamental constants by searching the derivative of the comparison frequency between a cryogenic sapphire oscillator and hydrogen maser. Phys Rev D. 2010;81:1.
268. van Zoest T, Gaaloul N, Singh Y, Ahlers H, Herr W, Seidel ST et al. Bose–Einstein condensation in microgravity. Science. 2010;328:1540–3. https://www.science.org/doi/pdf/10.1126/science.1189164.
269. Müntinga H, Ahlers H, Krutzik M, Wenzlawski A, Arnold S, Becker D et al. Interferometry with Bose–Einstein condensates in microgravity. Phys Rev Lett. 2013;110:093602.
270. Nyman R et al. I.C.E.: a transportable atomic inertial sensor for test in microgravity. Appl Phys B. 2006;84:673–81.
271. Frye K, Abend S, Bartosch W, Bawamia A, Becker D, Blume H et al. The Bose–Einstein condensate and cold atom laboratory. EPJ Quantum Technol. 2021;8:1.
272. Lezius M, Wilken T, Deutsch C, Giunta M, Mandel O, Thaller A et al. Space-borne frequency comb metrology. Optica. 2016;3:1381–7.
273. Dinkelaker AN, Schiemangk M, Schkolnik V, Kenyon A, Lampmann K, Wenzlawski A et al. Autonomous frequency stabilization of two extended-cavity diode lasers at the potassium wavelength on a sounding rocket. Appl Opt. 2017;56:1388–96.
274. Döringshoff K et al. Iodine frequency reference on a sounding rocket. Phys Rev Appl. 2019;11:054068.
275. Zhang X, del Aguila RP, Mazzoni T, Poli N, Tino GM. Trapped-atom interferometer with ultracold Sr atoms. Phys Rev A. 2016;94:043608.
276. Rudolph J, Wilkason T, Nantel M, Swan H, Holland CM, Jiang Y et al. Large momentum transfer clock atom interferometry on the 689 nm intercombination line of strontium. Phys Rev Lett. 2020;124:83604. arXiv:1910.05459.
277. Hu Z-K, Sun B-L, Duan X-C, Zhou M-K, Chen L-L, Zhan S et al. Demonstration of an ultrahigh-sensitivity atom-interferometry absolute gravimeter. Phys Rev A. 2013;88:043610.
278. ESA. Call for a medium-size and a fast mission opportunity in ESA's science programme. https://www.cosmos.esa.int/web/call-for-missions-2021.

UNCORRECTED PROOF

Submit your manuscript to a SpringerOpen journal and benefit from:

► Convenient online submission
► Rigorous peer review
► Open access: articles freely available online
► High visibility within the field
► Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com