



Review of CP-violation and spectroscopy measurements at LHCb



Neville Harnew
University of Oxford


On behalf of the LHCb Collaboration

**Corfu Summer
Institute
2 September 2021**


Outline

- General introduction
- An update of mixing and CP-violation measurements
 - New unitarity triangle measurements
 - Update on the angle γ
 - CP violation and mixing in charm
- New measurements in spectroscopy
- The upgraded LHCb detector and outlook
- Summary

Unitarity Triangle measurements




1995



2004

- Amazing progress in the last 26 years; the SM remains intact, but a whole lot still to learn


<http://ckmfitter.in2p3.fr>



Now (dominated by LHCb)

LHCb
Mixing and CP-
violation in
beauty and charm


$B_{(s)}$ mixing at LHCb




Previous Workshop

$$\frac{N(B^0 \rightarrow B^0) - N(B^0 \rightarrow \bar{B}^0)}{N(B^0 \rightarrow B^0) + N(B^0 \rightarrow \bar{B}^0)}$$

NEW




— $B_s^0 \rightarrow D_s^- \pi^+$ — $\bar{B}_s^0 \rightarrow B_s^0 \rightarrow D_s^- \pi^+$ — Untagged



$$\Delta m_d = (505.0 \pm 2.1 \pm 1.0) \text{ ns}^{-1}$$

Eur. Phys. J. C76 (2016) 412

Mixing measurements dominated by LHCb



LHCb-PAPER-2021-005, arXiv:2104.04421



$$\Delta m_s = 17.7683 \pm 0.0051 \pm 0.0032 \text{ ps}^{-1}$$

The angle γ (a key measurement)

- Loop processes are very sensitive to the presence of New Physics
- Constraints on the triangle apex largely come from **loop** decay measurements
- Large uncertainty on γ , the only angle accessible at **tree** level : forms a **SM benchmark***
- γ measurement theoretically very clean

JHEP 01 (2014) 051, PRD 92(3):033002 (2015)

* assuming no significant New Physics in tree decays



γ : indirect vs direct determinations


$$\gamma \equiv \arg \left[- \frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*} \right]$$

γ combination from all direct measurements from tree decays

$$\gamma = (72.1^{+5.4}_{-5.7})^\circ$$

(As of Summer 2019)

Reaching degree level precision from direct measurements is crucial



Determination from CKM fit excluding all direct measurements of γ


$$\gamma = (65.8^{+0.9}_{-1.3})^\circ$$

<http://ckmfitter.in2p3.fr>

The time-integrated mode: $B^- \rightarrow D^0 K^-$

$$\gamma \equiv \arg \left[-\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*} \right] \quad (\text{and charge conjugate mode } B^+ \rightarrow \bar{D}^0 K^+)$$

- Interference possible if D^0 and \bar{D}^0 decay to **same** final state
- Two possible decay paths to final state via D^0 and \bar{D}^0





- Branching fraction for favoured B decay only $\sim 10^{-4}$
 - Measurements require high statistics

New GLW & ADS γ measurements


GLW : where D^0 and \overline{D}^0 decay to CP eigenstates

ADS : where D^0 and \overline{D}^0 decay to flavour-specific states



$$A_K^{CP} = \frac{\Gamma(B^- \rightarrow [hh]_D^0 K^-) - \Gamma(B^+ \rightarrow [hh]_D^0 K^+)}{\Gamma(B^- \rightarrow [hh]_D^0 K^-) + \Gamma(B^+ \rightarrow [hh]_D^0 K^+)}$$

JHEP 04 (2021) 081



LHCb combination from different modes

LHCb-CONF-2021-001


- The most recent combination includes the following modes:

<i>B</i> decay	<i>D</i> decay	Ref.	Dataset	Lumi (fb ⁻¹)	Status since Ref. [21]	<i>D</i> decay	Ref.	Dataset	Lumi (fb ⁻¹)	Status since Ref. [21]
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+h^-$	[23]	Run 1&2	9	Updated	$D \rightarrow h^+h^-$	[35-37]	Run 1&2	9	New
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[24]	Run 1	3	As before	$D \rightarrow h^+h^-$	[38]	Run 1	3	New
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow h^+h^-\pi^0$	[25]	Run 1	3	As before	$D \rightarrow h^+h^-$	[39]	Run 1&2	9	New
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 h^+h^-$	[22]	Run 2	9	Updated	$D \rightarrow K^+\pi^-$	[40]	Run 1	3	New
$B^\pm \rightarrow Dh^\pm$	$D \rightarrow K_S^0 K^\pm \pi^\mp$	[26]	Run 1&2	9	Updated	$D \rightarrow K^+\pi^-$	[41]	Run 1&2	5	New
$B^\pm \rightarrow D^*h^\pm$	$D \rightarrow h^+h^-$	[23]	Run 1&2	5	Updated	$D \rightarrow K^\pm \pi^\mp \pi^+ \pi^-$	[42]	Run 1	3	New
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^+h^-$	[27]	Run 1&2	5	As before	$D \rightarrow K_S^0 \pi^+ \pi^-$	[43, 44]	Run 1&2	9	New
$B^\pm \rightarrow DK^{*\pm}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[27]	Run 1&2	5	As before	$D \rightarrow K_S^0 \pi^+ \pi^-$	[45]	Run 1	1	New
$B^\pm \rightarrow Dh^\pm \pi^+ \pi^-$	$D \rightarrow h^+h^-$	[28]	Run 1	3	As before					
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K^+\pi^-$	[29]	Run 1&2	5	Updated					
$B^0 \rightarrow DK^{*0}$	$D \rightarrow h^+\pi^-\pi^+\pi^-$	[29]	Run 1&2	5	New					
$B^0 \rightarrow DK^+\pi^-$	$D \rightarrow h^+h^-$	[30]	Run 1	3	Supersede					
$B^0 \rightarrow DK^{*0}$	$D \rightarrow K_S^0 \pi^+ \pi^-$	[31]	Run 1	3	As before					
$B^0 \rightarrow D^\mp \pi^\pm$	$D^+ \rightarrow K^- \pi^+ \pi^+$	[32]	Run 1	3	As before					
$B_s^0 \rightarrow D_s^\mp K^\pm$	$D_s^+ \rightarrow h^+h^- \pi^+$	[33]	Run 1	3	As before					
$B_s^0 \rightarrow D_s^\mp K^\pm \pi^+ \pi^-$	$D_s^+ \rightarrow h^+h^- \pi^+$	[34]	Run 1&2	9	New					

LHCb combination from different modes

New LHCb average

$$\gamma = (65.4^{+3.8}_{-4.2})^\circ$$



Previous measurement from LHCb

$$\gamma = (74.0^{+5.0}_{-5.8})^\circ \quad \text{LHCb-CONF-2018-002}$$

LHCb dominates world average

Reminder of indirect constraint

$$\gamma = (65.8^{+0.9}_{-1.3})^\circ$$

LHCb-CONF-2021-001

BaBar : $\gamma = (69^{+17}_{-16})^\circ$



PRD 87 (2013) 052015

Belle: $\gamma = (73^{+15}_{-14})^\circ$

arXiv:1301.2033

Breakdowns and evolution of γ results

LHCb-CONF-2021-001




$$\gamma = (65.4^{+3.8}_{-4.2})^\circ$$

Combination	Value	68.3% CL	95.4% CL
B^+	61.7	[57.1, 65.9]	[52.6, 69.8]
B^0	82.0	[73.7, 90.5]	[64.0, 98.0]
B_s^0	79.0	[59.0, 98.0]	[41.0, 106.0]

Beauty and Charm unitarity triangles

■ Beauty system



B system : angles $\alpha, \beta, \gamma \sim 1$

B_s system : angle $\beta_s \sim \lambda^2$


■ Charm system



Charm system : angle $\beta_c \sim \lambda^4$

Diagrams from Jolanta Brodzicka

B_s weak mixing phase ϕ_s in $B_s \rightarrow J/\psi \phi$




- “Golden mode” for this study is $B_s \rightarrow J/\psi \phi (\rightarrow K^+K^-)$
- Analogue of 2β (phase of B^0 mixing) but in the B_s system
- Interference between B^0 decay to $J/\psi \phi$ directly and via $B^0 - \bar{B}^0$ oscillation gives rise to a CP violating phase in the SM : a time-dependent measurement

$$\phi_S = \phi_{\text{Mixing}} - 2 \phi_{\text{Decay}} = -2\beta_s$$
- ϕ_S is expected to be very small in the SM and precisely predicted:
 $\phi_{\text{SM}} = -0.037 \pm 0.001 \text{ rad}$ (see eg Charles et al PRD84 (2011) 033005)

- ϕ_S fitted value correlated with $\Delta\Gamma_S =$ width diff. of the B_S mass eigenstates \rightarrow plot as contours in $(\phi_S \text{ vs } \Delta\Gamma_S)$ plane
- ϕ_S is 0.1σ from Standard Model and 1.6σ from zero

$$\Delta\Gamma_S = 0.0813 \pm 0.0048 \text{ ps}^{-1}$$

$$\text{CP-violating phase: } \phi_S = -0.040 \pm 0.025 \text{ rad}$$




HFLAV combination all experiments

CP-violating phase:

$$\phi_S = -0.050 \pm 0.019 \text{ rad}$$

$$(\phi_S^{\text{SM}} = -0.037 \pm 0.001 \text{ rad})$$




CP violation in charm

■ Direct CP violation

Measure asymmetry

$$\left| D^0 \rightarrow f \right|^2 \neq \left| \bar{D}^0 \rightarrow \bar{f} \right|^2 \quad A(D \rightarrow f) = \frac{N(D \rightarrow f) - N(\bar{D} \rightarrow \bar{f})}{N(D \rightarrow f) + N(\bar{D} \rightarrow \bar{f})}$$

- Most promising channels are *Cabibbo-suppressed* (CS) decays where CPV may arise from the *interference* between the **tree** and the **penguin** amplitudes



- SM prediction is very small $O(10^{-4}) \rightarrow O(10^{-3})$

Reminder of the “ ΔA_{CP} ” measurement

- Tag D^0 and \bar{D}^0 via “prompt” and “semileptonic” decays:

- ◆ Prompt: coming from primary vertex, i.e. $D^{*+-} \rightarrow \bar{D}^0 \pi^{+-}_{soft}$

- ◆ Semileptonic: coming from B-decays, i.e. $B^{+-} \rightarrow \bar{D}^0 \mu^{+} X$

- The *raw* asymmetry (**A**) in Cabibbo-suppressed $D^0 \rightarrow h^- h^+$ decays ($h = K$ or π) defined as

$$A(D \rightarrow f) = \frac{N(D \rightarrow f) - N(\bar{D} \rightarrow \bar{f})}{N(D \rightarrow f) + N(\bar{D} \rightarrow \bar{f})}$$

Phys. Rev. Lett. 122
(2019) 211803

includes physics and detector effects:

$$A = A_{CP} + A_D + A_P$$

Detection asymmetry
from π^{+}_{soft} or μ^{+}

Production asymmetry
from D^{*+} or B decays

- To eliminate these contributions and cancel the systematics measure :

$$\Delta A_{CP} = A(K^- K^+) - A(\pi^- \pi^+) = A_{CP}(K^- K^+) - A_{CP}(\pi^- \pi^+)$$



Observation of CPV in charm decays

- Measurement performed with combined Run-1 and Run-2 data-set

Phys. Rev. Lett. 122 (2019) 211803


$$\Delta A_{\text{CP}} = [-15.4 \pm 2.9] \times 10^{-4}$$

- A 5.3σ measurement of CPV in the charm system !



Charm CPV : more recent measurements


- Direct CPV : $A_{CP}(D^0 \rightarrow K_S^0 K_S^0)$ arXiv:2105.01565 (2021)
- Use $D^0 \rightarrow K^+ K^-$ channel as control for A_D & A_P
- $A_{CP} = (-3.1 \pm 1.2 \pm 0.4 \pm 0.2)\%$ [last uncertainty : CP violation of control channel]
- Consistent with no violation at the 2.4σ level



$A_{CP}(D^+ (s) \rightarrow h^+ \pi^0, h^+ \eta)$

JHEP 06 (2021) 019



$$\begin{aligned} \mathcal{A}_{CP}(D^+ \rightarrow \pi^+ \pi^0) &= (-1.3 \pm 0.9 \pm 0.6) \%, \\ \mathcal{A}_{CP}(D^+ \rightarrow K^+ \pi^0) &= (-3.2 \pm 4.7 \pm 2.1) \%, \\ \mathcal{A}_{CP}(D^+ \rightarrow \pi^+ \eta) &= (-0.2 \pm 0.8 \pm 0.4) \%, \\ \mathcal{A}_{CP}(D^+ \rightarrow K^+ \eta) &= (-6 \pm 10 \pm 4) \%, \\ \mathcal{A}_{CP}(D_s^+ \rightarrow K^+ \pi^0) &= (-0.8 \pm 3.9 \pm 1.2) \%, \\ \mathcal{A}_{CP}(D_s^+ \rightarrow \pi^+ \eta) &= (0.8 \pm 0.7 \pm 0.5) \%, \\ \mathcal{A}_{CP}(D_s^+ \rightarrow K^+ \eta) &= (0.9 \pm 3.7 \pm 1.1) \%, \end{aligned}$$



- All compatible with no CP violation
- More data needed !
- Note that LHCb is now regularly extracting measurements with neutrals in the final state ($K_s K_s$ and $h^0 h^+$)

D^0 mixing parameters in $D^0 \rightarrow K_S^0 \pi^+ \pi^-$

- Mass eigenstates $|D_{1,2}\rangle = p|D^0\rangle \pm q|\overline{D}^0\rangle$
- $x = (m_1 - m_2)/\Gamma$; $y = (\Gamma_1 - \Gamma_2)/2\Gamma$, $\phi = \arg(q/p)$
until now x measured only at $\sim 3\sigma$ (HFLAV)
- 30.6×10^6 of $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays with very small background. \overline{D} or D flavour tagging using $D^* \rightarrow D \pi$ decays
- Use the bin-flip method
 - ◆ Measure ratios between D^0 and \overline{D}^0 candidates in symmetric bins of Dalitz plot $m^2(K_S^0 \pi^-)$ vs $m^2(K_S^0 \pi^+)$
 - ◆ 2 (flavour) \times 16 (Dalitz bin) \times 13 (decay time bin) subsamples
 - ◆ In each bin, strong-phase difference approx. constant for D^0 and \overline{D}^0 amplitudes (input from CLEOc and BESIII)





$$m_{\pm}^2 \equiv \begin{cases} m^2(K_S^0 \pi^{\pm}) & \text{for } D^0 \rightarrow K_S^0 \pi^+ \pi^- \\ m^2(K_S^0 \pi^{\mp}) & \text{for } \overline{D}^0 \rightarrow K_S^0 \pi^+ \pi^- \end{cases}$$

ΔY in $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays

arXiv:2105.09889

- ΔY is the slope of the time-dependent asymmetry of the decay rates of D^0 and \bar{D}^0 mesons
- It is a measure of **CP violation in mixing and interference**
- Strategy: measure asymmetry in bins of decay time and measure the linear slope



$$\Delta Y_{K^+K^-} = (-2.3 \pm 1.5 \pm 0.3) \times 10^{-4}$$


$$\Delta Y_{\pi^+\pi^-} = (-4.0 \pm 2.8 \pm 0.4) \times 10^{-4}$$

Combining


$$\Delta Y = (-2.7 \pm 1.3 \pm 0.3) \times 10^{-4}$$

- Compatible with 0 within 2σ
- This result improves by nearly a factor 2 the precision of the previous world average


LHCb new (exotic) spectroscopy measurements




Baryon



Meson




Tetraquark




Pentaquark

New hadron discoveries at the LHC



With thanks to Partick Koppenberg

New hadron discoveries at LHCb



With thanks to Partick Koppenberg

Pentaquark discovery by LHCb

- Discovery of $X(3872)$ - now $\chi_{c1}(3872)$ - by Belle in 2003 started new era in exotic spectroscopy
- First observation of $P_c(4312)^+$, $P_c(4440)^+$ and $P_c(4457)^+$ as narrow resonances in the mass spectrum of $(J/\psi p)$ in $\Lambda_b \rightarrow (J/\psi p) K^-$ decays PRL 115 (2015) 072001
- Consistent with $c\bar{c}uud$ pentaquarks : allowed by QCD, but not observed in 50 years of searching.








Diagram from Liming Zhang



Nature of pentaquarks ?

Possible models describing the observed pentaquark states :

- Tightly bounded states
- Re-scattering models
- Meson-baryon molecules





- Molecular-state model favoured : bound mesons and baryons are expected to form narrow resonances just below mass thresholds
- More work needed


Evidence for more pentaquark states

- Amplitude analysis using 800 $B_s^0 \rightarrow J/\psi p \bar{p}$ decays
- Observe additional structure in $J/\psi p$ and $J/\psi \bar{p}$ spectra
- Significance of 3.1σ to 3.7σ depending on J^P assignment
- Evidence for new $P_c(4337)^+$ state consistent with another $(cc\bar{u}ud)$ pentaquark
- Amplitude analysis using 1750 $\Xi_b^- \rightarrow J/\psi \Lambda K^-$ decays
- Observe structure in $J/\psi \Lambda$ spectrum
- Evidence for new $P_{cs}(4459)^0$ state with significance of 3.1σ
- Consistent with $(cc\bar{u}uds)$ pentaquark

arXiv:2108.04720



	$M[\text{MeV}]$	$\Gamma[\text{MeV}]$
$P_c(4337)^+$	$4337_{-4}^{+7} \pm 2$	$29_{-12}^{+26} \pm 14$
$P_{cs}(4459)^0$	$4458.8 \pm 2.9_{-1.1}^{+4.7}$	$17.3 \pm 6.5_{-5.7}^{+8.0}$



Sci.Bull.66(2021)1278

New doubly charmed tetraquark T_{cc}^+

- Study $D^0 D^0 \pi^+$ mass spectrum near $D^{*+}D^0$ and $D^{*0}D^+$ thresholds



$$\delta m \equiv m_{T_{cc}^+} - (m_{D^{*+}} + m_{D^0})$$

- Very narrow state in $D^0 D^0 \pi^+$ mass spectrum consistent with $ccu\bar{d}$ tetraquark, with significance 10σ . Manifestly exotic state.

- Very close to $D^{*+}D^0$ mass thresholds

$$\delta m_{BW} = -273 \pm 61 \text{ keV}/c^2$$


$$\Gamma_{BW} = 410 \pm 165 \text{ keV}$$



- Possible evidence for molecular bound state, but jury still out.

More observations of new tetraquark states


- $B^+ \rightarrow J/\psi \phi K^+$ sample
- Observe structure in $J/\psi K$
- Observation of two new $c \bar{c} u \bar{s}$ tetraquark states $Z_{cs}(4000)^+$ and $Z_{cs}(4220)^+$
- Significance of 15σ and 6σ respectively, I^+ assignment



Phys. Rev. Lett. 127 (2021) 082001

$Z_{cs}(4000)$	15 (16)	$4003 \pm 6^{+4}_{-14}$	$131 \pm 15 \pm 26$
$Z_{cs}(4220)$	5.9 (8.4)	$4216 \pm 24^{+43}_{-30}$	$233 \pm 52^{+97}_{-73}$


- $B^+ \rightarrow J/\psi \phi K^+$ sample
- Observe structure in $J/\psi \phi$
- Observation of two new $c \bar{c} s \bar{s}$ tetraquark states $X(4630)$ and $X(4685)$ as well as previously confirmed states
- Significance of 5.5σ and 15σ respectively




$X(4630)$	5.5 (5.7)	$4626 \pm 16^{+18}_{-110}$	$174 \pm 27^{+134}_{-73}$
$X(4685)$	15 (15)	$4684 \pm 7^{+13}_{-16}$	$126 \pm 15^{+37}_{-41}$

The upgraded LHCb detector and outlook

LHCb Upgrade planning


WE ARE
HERE

2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	203+	
		Run III						Run IV					Run V		
LS2						LS3						LS4			
LHCb 40 MHz UPGRADE I		$L = 2 \times 10^{33}$			LHCb Consolidate: UPGRADE Ib			$L = 2 \times 10^{33}$ 50 fb^{-1}			LHCb UPGRADE II		$L = 1-2 \times 10^{34}$ 300 fb^{-1}		
ATLAS Phase I Upgr		$L = 2 \times 10^{34}$			ATLAS Phase II UPGRADE			HL-LHC $L = 5 \times 10^{34}$					HL-LHC $L = 5 \times 10^{34}$		
CMS Phase I Upgr		300 fb^{-1}			CMS Phase II UPGRADE								3000 fb^{-1}		
Belle II		5 ab^{-1}			$L = 6 \times 10^{35}$			50 ab^{-1}							




Luminosity $4 \times 10^{32} \text{ cm}^{-2}\text{s}^{-1}$
 ~1.1 visible interactions/crossing
 ~9 fb^{-1} collected

Luminosity $2 \times 10^{33} \text{ cm}^{-2}\text{s}^{-1}$
 ~5.5 visible interactions/crossing
 Up to 50 fb^{-1} collected

Luminosity $2 \times 10^{34} \text{ cm}^{-2}\text{s}^{-1}$
 ~55 visible interactions/crossing
 300 fb^{-1} collected

LHCb Upgrade I



New silicon upstream tracker (UT)

New scintillating fibre tracker (SciFi)

New electronics for muon and calorimeter systems

New PIXEL vertex detector (VELO)


New RICH1 and photodetectors

New RICH2 photodetectors

Plus new trigger

Construction & Installation – Upgrade I

SciFi tracker



Corfu Summer Institute

RICH 2




UT stove




N. Harnew

γ prospects : Run II \rightarrow Upgrade I

- Post Run II target of 4° almost surpassed ($\sim 9 \text{ fb}^{-1}$) and analyses still in progress 


- LHCb Upgrade I : target 0.9° ($\sim 50 \text{ fb}^{-1}$)

$$\gamma = (65.4^{+3.8}_{-4.2})^\circ$$




EPJC (2013) 73:2373

... and beyond 2030 : Upgrade II





Evolution of the Unitarity Triangle

LHCB-PUB-2018-009



LHCb : 2021
Run 2 ($\sim 9 \text{ fb}^{-1}$)

LHCb Upgrade I
2025 ($\sim 23 \text{ fb}^{-1}$)



LHCb Upgrade II
2035 (300 fb^{-1})

2021

N. Harnew

39

Summary and Outlook

- The LHCb experiment has performed spectacularly well :
→ $\sim 9 \text{ fb}^{-1}$ of recorded data up to $\sqrt{s} = 13 \text{ TeV}$
- So far all Unitarity Triangle measurements are consistent with the Standard Model
→ New Physics is becoming constrained
- LHCb is a fantastic platform for spectroscopy measurements: many measurements were never foreseen in LHCb's original physics portfolio.
- Still much room for New Physics, but higher precision required
→ preparing for LHCb Upgrades beyond 2022 and the decade afterwards!