



The AION (& AEDGE) Project

A UK Atom Interferometer Observatory and Network

to explore Ultra-Light Dark Matter
and Mid-Frequency Gravitational Waves.

O. Buchmueller, Imperial College London



Project executed in national partnership with **UK National Quantum Technology Hub in Sensors and Timing, Birmingham, UK,** and international partnership with **The MAGIS Collaboration and The Fermi National Laboratory, US**

The AION Programme consists of 4 Stages

❑ **Stage 1:** to build and commission the 10 m detector, develop existing technology and the infrastructure for the 100 m.

L ~ 10m

❑ **Stage 2:** to build, commission and exploit the 100 m detector and carry out a design study for the km-scale detector.

L ~ 100m

- AION was selected in 2018 by STFC as a high-priority medium-scale project.
- AION will work in equal partnership with MAGIS in the US to form a “LIGO/Virgo-style” network & collaboration, providing a pathway for UK leadership.

Stage 1 is now funded with about £9.6M by the QTFP Programme and other sources and Stage 2 could be placed at national facility in Boulby or Daresbury (UK), possibly also at CERN (France/Switzerland).



❑ **Stage 3:** to build a kilometre-scale terrestrial detector.

L ~ 1km

❑ **Stage 4:** long-term objective a pair of satellite detectors (thousands of kilometres scale) [AEDGE proposal to ESA Voyage2050 call]

- AION has established science leadership in AEDGE, bringing together collaborators from European and Chinese groups (e.g. MIGA, MAGIA, ELGAR, ZAIGA).


Stage 3 and 4 will likely require funding on international level (ESA, EU, etc) and AION has already started to build the foundation for it.



AION-10 @ Beecroft building, Oxford Physics

AION Corfu Workshop on Standard Model and Beyond


- New purpose-built building (£50M facility)
- AION-10 on basement level with 14.7m headroom (stable concrete construction)
- World-class infrastructure
- Experienced Project Manager:
- Engineering support from RAL (Oxfordshire)



10m



Laser lab for AION
vibration criterion, VC-G =
10nm@10Hz. Temperature
(22±0.1)° C





AION-10 @ Beecroft building, Oxford Physics

AION Corfu Workshop on Standard Model and Beyond

- New purpose-built building (£50M facility)
- AION-10 headroom
- World-class
- Experienced
- Engineering

For the first 30 months of the project, we will focus on the prerequisites for the 10m detector:



- Establish the Cold Atom infrastructure (e.g. build UltraCold Sr Laser Labs) and expertise
- Develop full design for 10m detector, ready for physics exploitation
- Partner AION with the MAGIS experiment in the US



Laser lab for AION
vibration criterion, VC-G =
10nm@10Hz. Temperature
(22±0.1)° C



AION: Ultra-Cold Strontium Laboratories in UK




To push the state-of-the-art single photon Sr Atom Interferometry, the AION project builds dedicated Ultra-Cold Strontium Laboratories in:
Birmingham, Cambridge, Imperial College, Oxford, and RAL

The laboratories are expected to be fully operational in fall 2021.

Example: New Ultra-Cold Strontium Laboratory at Imperial College


AION Corfu Workshop on Standard Model and Beyond

- In the context of the AION project, Imperial College is building a **brand new Ultra-Cold Strontium Laboratory on about 100m²** of high-quality real state in the Blakett Laboratory in central London.
- This Laboratory will extent significant the already very strong Ultra-Cold Atom facilities at Imperial, exploiting strong synergies with existing laboratory space in the same building.
- The new Laboratory will be **operated by world leading experts in Ultra-Cold Strontium Atomic Clocks**, and complements the already established expertise in Rb Atom Interferometry and Magneto-optical trapping and sympathetic cooling of molecules.
- **The new Laboratory will be ready for full operation by the end of 2021.**





Ground Based Large Scale O(100m) Projects


MIGA: Terrestrial detector using atom interferometer at O(100m)
(France)



ZIGA: Terrestrial detector for large scale atomic interferometers, gyros and clocks at O(100m)
(China)



AION: Terrestrial shaft detector using atom interferometer at 10m – O(100m) planned
(UK)





MAGIS: Terrestrial shaft detector using atom interferometer at O(100m)
(US)

Planned network operation

Ground Based Large Scale O(100m) Projects

MIGA: Terrestrial detector using atom interferometer at O(100m)
(France)



AION: Terrestrial shaft detector using atom interferometer at 10m – O(100m) planned
(UK)

Example of large-scale CA projects that act as demonstrators for GW mid-frequency band and DM detectors.



All these projects are represented in the AEDGE consortium.

Each project requires an investment of O(10M) currency units.

All projects (AION, MAGIS, MIGA, ZIGA) are funded by national funding agencies and foundations.

ZIGA:
interf

Timeline 2020 to 2025ish





MAGIS: Terrestrial shaft detector using atom interferometer at O(100m)
(US)

Planned network operation



QUANTUM TECHNOLOGY: ATOM INTERFEROMETRY IN A NUTSHELL

Light vs. Cold Atoms: Atom Interferometry

Light interferometer




Atom interferometer




Simple Example: Two Atomic Clocks

AION Corfu Workshop on Standard Model and Beyond




Simple Example: Two Atomic Clocks

AION Corfu Workshop on Standard Model and Beyond



Phase Noise from the Laser


The phase of the laser is imprinted onto the atom.





*Laser phase is **common** to both atoms – rejected in a differential measurement.*

AION: A Different Kind of Atom Interferometer

Hybrid “clock accelerometer”



Clock transition in candidate atom ^{87}Sr



$$\Delta\phi \sim \omega_A (2L/c)$$

Two ways for phase to vary:

$$\delta\omega_A \quad \text{Dark matter}$$

$$\delta L = hL \quad \text{Gravitational wave}$$

Clock: measure light travel time \rightarrow remove laser noise with *single baseline*

Sensitivity Scenario	L [m]	T_{int} [sec]	$\delta\phi_{noise}$ [$1/\sqrt{\text{Hz}}$]	LMT [number n]
AION-10 (initial)	10	1.4	10^{-3}	100
AION-10 (goal)	10	1.4	10^{-4}	1000
AION-100 (initial)	100	1.4	10^{-4}	1000
AION-100 (goal)	100	1.4	10^{-5}	40000
AION-km	2000	5	0.3×10^{-5}	40000

Used for sensitivity projections


For ultimate sensitivity we need to push each basic parameter by $\sim O(10)$.

The project aims to demonstrate in funding period e.g.

- LMT: ~ 1000 hbar*k
- Squeezing $\sim 20\text{dB}$ for $> 1e6$ Atoms

Main AION Physics Goals: Dark Matter and Gravitational Waves

AION Corfu Workshop on Standard Model and Beyond



Scientific Leadership in phenomenology already established:

The AION Physics Case:

AION Collaboration, AION: An Atom Interferometer Observatory and Network, arXiv:1911.11755. [accepted for publication in JCAP]

AEDGE


Y. El-Neaj, ..., O. Buchmueller *et al.*
AEDGE: Atomic Experiment for Dark Matter And Gravity Exploration in Space, arXiv:1908.00802, *EPJ Quantum Technol.* 7, 6 (2020). [Submitted to ESA Voyage2050 call]

Working with leading theorists:

J. Ellis, M. Haehnelt, C. McCabe, J. March-Russell (AION), C. Burrage, ...


Main Physics Goals:

- Search for Ultra-Light Dark Matter
 - Explore new parameter space and complement other searches.
 - Focus on Scalar DM with Vector and Pseudoscalar DM also under study.
- Gravitational Waves in mid-frequency band
 - Explore frequencies between LISA and LIGO/VIRGO, KAGRA and Einstein Telescope
 - Targets: Black hole mergers, phase transitions and cosmic string collisions



Main AION Physics Goals: Dark Matter and Gravitational Waves

AION Corfu Workshop on Standard Model and Beyond



Scientific Leadership in phenomenology already established:

The AION Physics Case:

AEDGE

AION Collaboration, AION: An Atom Interferometer, Y. El-Neai, O. Buchmueller et al.


In this talk I will focus on Ultra-Light Dark Matter but AION also has a very promising Gravitation Wave programme.

More information about the science is provided in:
 AION: [arXiv:1911.11755](https://arxiv.org/abs/1911.11755), *JCAP05(2020)011*,
 AEDGE: [arXiv:1908.00802](https://arxiv.org/abs/1908.00802), *EPJ Quantum Technol.* 7, 6 (2020).

Dark Matter
arXiv:1908.00802,

Main Physics

- Search for Ultra-Light Dark Matter
 - Explore new physics
 - Focus on Standard Model particles under study.
- Gravitational Waves in mid-frequency band
 - Explore frequencies between LISA and LIGO/VIRGO, KAGRA and Einstein Telescope
 - Targets: Black hole mergers, phase transitions and cosmic string collisions



SEARCHES FOR ULTRA-LIGHT DARK MATTER

The Landscape of Ultra-Light Dark Matter Detection

Very light dark matter and gravitational wave detection similar when detecting coherent effects of entire field, not single particles.

Example: Ultra-Light Dark Matter:




Diagram taken from P. Graham's talk at HEP Front 2018

The Landscape of Ultra-Light Dark Matter Detection

Very light dark matter and gravitational wave detection similar when detecting coherent effects of entire field, not single particles.

Example: Ultra-Light Dark Matter:




Diagram taken from P. Graham's talk at HEP Front 2018

The Landscape of Ultra-Light Dark Matter Detection

Very light dark matter and gravitational wave detection similar when detecting coherent effects of entire field, not single particles.

Example: Ultra-Light Dark Matter:




Diagram taken from P. Graham's talk at HEP Front 2018


Ultra-Light Spin-0 Dark Matter

Ultra-light spin 0 particles are expected to form a coherently oscillating classical field

$$\phi(t) = \phi_0 \cos(E_\phi t / \hbar)$$

as $E_\phi \approx m_\phi c^2$ with an energy density of

$$\langle \rho_\phi \rangle \approx m_\phi^2 \phi_0^2 / 2 \quad (\rho_{DM,local} \approx 0.4 \text{ GeV/cm}^3).$$



Ultralight scalar dark matter


Ultralight dilaton DM acts as a background field (e.g., mass $\sim 10^{-15}$ eV)

$$\mathcal{L} = + \underbrace{\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m_\phi^2 \phi^2}_{\text{DM scalar field}} - \sqrt{4\pi G_N} \phi \left[\underbrace{d_{m_e} m_e \bar{e} e}_{\text{Electron coupling}} - \underbrace{\frac{d_e}{4} F_{\mu\nu} F^{\mu\nu}}_{\text{Photon coupling}} \right] + \dots$$

e.g., QCD

$$\phi(t, \mathbf{x}) = \phi_0 \cos [m_\phi(t - \mathbf{v} \cdot \mathbf{x}) + \beta] + \mathcal{O}(|\mathbf{v}|^2) \quad \phi_0 \propto \sqrt{\rho_{\text{DM}}} \quad \text{DM mass density}$$

DM coupling causes time-varying atomic energy levels:





Courtesy of Jason Hogan!

AION Corfu Workshop on Standard Model and Beyond

Ultra-Light Scalar Dark Matter


AION Corfu Workshop on Standard Model and Beyond



The AION staged programme will have unprecedented sensitivity to DM with scalar couplings to matter, which cause time variation of fundamental constants such as the electron mass.


Based on: Arvanitaki et al., PRD **97**, 075020 (2018).

← **Linear scalar DM interactions**
Quadratic scalar DM interactions →



Ultra-Light Scalar Dark Matter

AION Corfu Workshop on Standard Model and Beyond





AION will be probing new territory in ULD scalar scenarios.
 With different configurations of the Atom Interferometer
 it will be also possible to search for
Axions (pseudo-scalar) and **Vector** DM candidates!
 [studies are ongoing]

A Space based version of AION (i.e. AEDGE), would cover an enormous region of parameter space with unprecedented discovery potential!

Quadratic scalar DM interactions \longrightarrow photo

AEDGE





A
E
D
G
E

AEDGE:
Atomic Experiment
for Dark Matter
and Gravity
Exploration

The Space Version of AION – Stage 4 of the Programme

AEDGE: Atomic Experiment for Dark Matter and Gravity Exploration



Informal Workshop
CERN, July 22/23 2019

Organizers:

Kai Bongs(CA), Philippe Bouyer(CA), Oliver Buchmueller(PP),
Albert De Roeck(PP), John Ellis(PP, Theory), Peter Graham (CA, Theory),
Jason Hogan (CA), Wolf von Klitzing(CA), Guglielmo Tino(CA), and AtomQT
PP=Particle Physics
CA=Cold Atoms

AEDGE: Atomic Experiment for Dark Matter and Gravity Exploration

**With more than 130 participants
the workshop was very well attended!**

**The full agenda can be accessed via:
<https://indico.cern.ch/event/830432/timetable/>**

**The main scope was to review the
landscape of Cold Atom
experiments on ground AND in
space to eventually establish a
roadmap for technology readiness
for space.**

**Informal Workshop
CERN, July 22/23 2019**

Organizers:

**Kai Bongs(CA), Philippe Bouyer(CA), Oliver Buchmueller(PP),
Albert De Roeck(PP), John Ellis(PP, Theory), Peter Graham (CA, Theory),
Jason Hogan (CA), Wolf von Klitzing(CA), Guglielmo Tino(CA), and AtomQT**
PP=Particle Physics
CA=Cold Atoms

AEDGE Mission Concept

AEDGE: Atomic Experiment for Dark Matter and Gravity Exploration in Space

Yousef Abou El-Neaj,¹ Cristiano Alpigiani,² Sana Amairi-Pyka,³ Henrique Araújo,⁴ Antun Balaž,⁵ Angelo Bassi,⁶ Lars Bathe-Peters,⁷ Baptiste Battelier,⁸ Aleksandar Belić,⁵ Elliot Bentine,⁹ José Bernabeu,¹⁰ Andrea Bertoldi,^{8,*} Robert Bingham,¹¹ Diego Blas,¹² Vasiliki Bolpasi,¹³ Kai Bongs,^{14,*} Sougato Bose,¹⁵ Philippe Bouyer,^{8,*} Themis Bowcock,¹⁶ William Bowden,¹⁷ Oliver Buchmueller,^{4,*} Clare Burrage,¹⁸ Xavier Calmet,¹⁹ Benjamin Canuel,^{8,*} Laurentiu-Ioan Caramete,^{20,*} Andrew Carroll,¹⁶ Giancarlo Cella,^{21,22} Vassilis Charmandaris,²³ Swapan Chattopadhyay,^{24,25} Xuzong Chen,²⁶ Maria Luisa Chiofalo,^{21,22} Jonathon Coleman,^{16,*} Joseph Cotter,⁴ Yanou Cui,²⁷ Andrei Derevianko,²⁸ Albert De Roeck,^{29,30,*} Goran Djordjevic,³¹ Peter Dornan,⁴ Michael Doser,³⁰ Ioannis Drougkakis,¹³ Jacob Dunningham,¹⁹ Ioana Dutan,²⁰ Sajjan Easo,¹¹ Gedminas Elertas,¹⁶ John Ellis,^{12,32,33,*} Mai El Sawy,³⁴ Farida Fassi,³⁵ Daniel Felea,²⁰ Chen-Hao Feng,⁸ Robert Flack,¹⁵ Chris Foot,⁹ Ivette Fuentes,¹⁸ Naceur Gaaloul,³⁶ Alexandre Gauguier,³⁷ Remi Geiger,³⁸ Valerie Gibson,³⁹ Gian Giudice,³³ Jon Goldwin,¹⁴ Oleg Grachov,⁴⁰ Peter W. Graham,^{41,*} Dario Grasso,^{21,22} Maurits van der Grinten,¹¹ Mustafa Gündogan,³ Martin G. Haehnel,^{42,*} Tiffany Harte,³⁹ Aurélien Hees,^{38,*} Richard Hobson,¹⁷ Bodil Holst,⁴³ Jason Hogan,^{41,*} Mark Kasevich,⁴¹ Bradley J. Kavanagh,⁴⁴ Wolf von Klitzing,^{13,*} Tim Kovachy,⁴⁵ Benjamin Kriker,⁴⁶ Markus Krutzik,^{3,*} Marek Lewicki,^{12,47,*} Yu-Hung Lien,¹⁵ Miaoyuan Liu,²⁶ Giuseppe Gaetano Luciano,⁴⁸ Alain Magnon,⁴⁹ Mohammed Mahmoud,⁵⁰ Sarah Malik,⁴ Christopher McCabe,^{12,*} Jeremiah Mitchell,²⁴ Julia Pahl,³ Debapriya Pal,¹³ Saurabh Pandey,¹³ Dimitris Papazoglou,⁵¹ Mauro Paternostro,⁵² Bjoern Penning,⁵³ Achim Peters,^{3,*} Marco Prevedelli,⁵⁴ Vishnupriya Puthiya-Veetil,⁵⁵ John Quenby,⁴ Ernst Rasel,^{36,*} Sean Ravenhall,⁹ Haifa Rejeb Sfar,²⁹ Jack Ringwood,¹⁶ Albert Roura,^{56,*} Dylan Sabulsky,^{8,*} Muhammed Sameed,⁵⁷ Ben Sauer,⁴ Stefan Alaric Schäffer,⁵⁸ Stephan Schiller,^{59,*} Vladimir Schkolnik,³ Dennis Schlippert,³⁶ Christian Schubert,^{3,*} Armin Shayeghi,⁶⁰ Ian Shipsey,⁹ Carla Signorini,^{21,22} Marcelle Soares-Santos,⁵³ Fiodor Sorrentino,^{61,*} Yajpal Singh,^{14,*} Timothy Sumner,⁴ Konstantinos Tassis,¹³ Silvia Tentindo,⁶² Guglielmo Maria Tino,^{63,64,*} Jonathan N. Tinsley,⁶³ James Unwin,⁶⁵ Tristan Valenzuela,¹¹ Georgios Vasilakis,¹³ Ville Vaskonen,^{12,32,*} Christian Vogt,⁶⁶ Alex Webber-Date,¹⁶ André Wenzlawski,⁶⁷ Patrick Windpassinger,⁶⁷ Marian Woltmann,⁶⁶ Michael Holynski,¹⁴ Efe Yazgan,⁶⁸ Ming-Sheng Zhan,^{69,*} Xinhao Zou,⁸ Jure Zupan⁷⁰



132 Authors, from **70** institutions,
based in **23** different counties!

The authors represent several science communities ranging from Cold Atoms, & Gravitational Waves, over Cosmology and Astrophysics to fundamental Particle Physics.

<https://arxiv.org/abs/1908.00802>

The paper is now published in **EPJ Quantum Technology**:
EPJ Quantum Technol. 7, 6 (2020).

Potential Mission Design




Using two cold-atom interferometers that perform a relative measurement of differential phase shift, a potential mission profile would be using a pair of satellites separated by a very long baseline L .

Assumed basic parameters:

- Pair of satellites in medium earth orbit (MEO)
- Satellite separation $L = 4.4 \times 10^7$ m

Note: as Laser noise is common-mode suppressed only two satellites are required

Community Workshop on Cold Atom in Space: Sep 23/24



AION Corfu Workshop on Standard Model and Beyond

➤ The purpose of this community workshop is to discuss options for a quantum technology development programme and to develop a community roadmap and milestones to demonstrate the readiness of cold atom technologies in space, as proposed in the Voyage 2050 recommendations.

➤ This event will bring together the cold atom, astrophysics, cosmology, fundamental physics, and earth observation communities to shape this development programme.

Link to the Workshop Webpage:
<https://indico.cern.ch/event/1064855/>

Workshop Registration:
<https://indico.cern.ch/event/1064855/registrations/74939/>

We strongly encourage you to register, even if you are unable to attend the virtual workshop, as this list will also be used to coordinate the Community Roadmap Process that will follow the workshop.

Summary: AION & AEDGE


- New window on gravitational physics, astrophysics & cosmology using atom interferometers, leveraging already existing investment in quantum technologies, providing new opportunities for science communities.
- AION-10 was funded by the QTFP programme and will explore parameter space **of ultra-light dark matter (ULDM)** models, partnership with MAGIS in US.
- Preparation for AION-100 (km-scale) with **unique capabilities for detecting gravitational waves** is key deliverable.
 - Funding required would be similar to that for AION-10, assuming a suitable site.
 - Possible 100m sites under investigation: Boulby, Daresbury (UK), CERN (France/Switzerland).
- AEDGE is a uniquely interdisciplinary mission that will harness cold atom technologies, as developed for AION, to address key issues in fundamental physics, astrophysics and cosmology that can be realized within the Voyage 2050 Science Programme of ESA.
 - AEDGE is currently under review by ESA and we are planning to host another AEDGE workshop when the results of the review are available.

BACKUP

BOULBY AND CERN SITE OPTION

Possible CERN Site for AION 100m

AION Corfu Workshop on Standard Model and Beyond



PX46 – P4 Support shaft
Lengths 143m
D = 10.10m


➤ **Ideal basic parameters for AION100**

First radiation studies are also looking promising but more work is needed to determine if PX46 could be a valid option for AION 100.

We are working with PBC Team on this feasibility study (K Balazs and before J. Gall)

Next step: looking with RP and HE department into the feasibility of the shielding arrangement.

Other site options that are currently investigated are the ***national facility in Boulby and Daresbury (UK).***



Location of a possible short shaft at Boulby for AION-100

AION Corfu Workshop on Standard Model and Beyond



Length 143m
Inner circle diameter 5 m

Two possible locations for AION-100

- next to the staircase : ~ 1.2 m x ~ 0.8 m
- on the other side of the cage - more space but no access from the staircase


Possible
location
for
AION-100

Inner circle diameter : 5m

Possible
location for
AION-100
1.2 m x 0.8 m

Staircase


Working cage
2 m x 2 m



AION Corfu Workshop on Standard Model and Beyond


Basic Differential Measurement

AION Corfu Workshop on Standard Model and Beyond



Basic Differential Measurement

AION Corfu Workshop on Standard Model and Beyond




Laser 2: π pulse [high p]

Laser 1: $\pi/2$ pulse [split]

Basic Differential Measurement

AION Corfu Workshop on Standard Model and Beyond




“Mirror”
 3π pulse
 [low-high/low-high]
 [Doppler shift to select]

Laser 2: π pulse [high p]

Laser 1: $\pi/2$ pulse [split]


Basic Differential Measurement

AION Corfu Workshop on Standard Model and Beyond



Basic Differential Measurement

AION Corfu Workshop on Standard Model and Beyond



Laser 1: $\pi/2$ pulse [split]

Laser 2: π pulse [low p]

“Mirror”
 3π pulse
 [low-high/low-high]
 [Doppler shift to select]

Laser 2: π pulse [high p]

Laser 1: $\pi/2$ pulse [split]

Each AI spends time L/c in excited state but at different periods in the sequence

Team roles and linkages in AION and MAGIS

AION Corfu Workshop on Standard Model and Beyond

MAGIS-100

Joint work includes:-

- Jon Coleman (Liverpool) is a founding member of the MAGIS project: Design and fabrication of key parts by Liverpool Physics.
- Hardware deliverables to MAGIS: Cameras (Oxf.), Electronics (Cam.)
- Assisting in construction, commissioning and data-taking at Fermilab site.
- Participation in data analysis and first results.
- Kavli-funded PDRA (Cam.)



UK laser company:
Unique systems
for Q Tech. with Sr

King's + Imperial Colleges:
Theory and publication office

UoB, Cambridge, Imperial:
modelling system parameters


Imperial: (clock) laser
stabilisation, squeezing

RAL: Vacuum + support
structure. Design AION-100



- Impact
- Technology transfer


10m





EXPERIMENTAL LANDSCAPE

Ground Based Large Scale O(100m) Projects

MIGA: Terrestrial detector using atom interferometer at O(100m)
(France)




ZIGA: Terrestrial detector for large scale atomic interferometers, gyros and clocks at O(100M)
(China)



MAGIS: Terrestrial shaft detector using atom interferometer at O(100m)
(US)

AION: Terrestrial shaft detector using atom interferometer at 10m – O(100m) planned
(UK)



Planned network operation

STATE-OF-THE-ART DESIGN SPECIFICATIONS

THE PHYSICS CASE

Based on DM workshop at KCL:

<https://indico.cern.ch/event/797031/timetable/>

and AION workshop at Imperial:

<https://indico.cern.ch/event/802946/>

Using Material from. M. Bauer, J. Hogan, J. March-Russel, C. McCabe, and Y. Stadnik

DARK MATTER PHYSICS @AION

Ultralight scalar dark matter


Ultralight dilaton DM acts as a background field (e.g., mass $\sim 10^{-15}$ eV)

$$\mathcal{L} = + \underbrace{\frac{1}{2} \partial_\mu \phi \partial^\mu \phi - \frac{1}{2} m_\phi^2 \phi^2}_{\text{DM scalar field}} - \sqrt{4\pi G_N} \phi \left[\underbrace{d_{m_e} m_e \bar{e} e}_{\text{Electron coupling}} - \underbrace{\frac{d_e}{4} F_{\mu\nu} F^{\mu\nu}}_{\text{Photon coupling}} \right] + \dots$$

e.g., QCD

$$\phi(t, \mathbf{x}) = \phi_0 \cos [m_\phi(t - \mathbf{v} \cdot \mathbf{x}) + \beta] + \mathcal{O}(|\mathbf{v}|^2) \quad \phi_0 \propto \sqrt{\rho_{\text{DM}}} \quad \text{DM mass density}$$



DM coupling causes time-varying atomic energy levels:



Courtesy of Jason Hogan!

AION Corfu Workshop on Standard Model and Beyond


Ultra-Light Scalar Dark Matter



The AION staged programme will have unprecedented sensitivity to DM with scalar couplings to matter, which cause time variation of fundamental constants such as the electron mass.

Based on: Arvanitaki et al., PRD **97**, 075020 (2018).

← Linear scalar DM interactions
 Quadratic scalar DM interactions →



Ultra-





Table 1. List of basic parameters: length of the detector L ; interrogation time of the atom interferometer T_{int} ; phase noise $\delta\phi_{noise}$; and number of momentum transfers LMT . The choices of these parameters largely determine the sensitivities of the projection scenarios. It should be noted that at a 100m detector it will be conceptually possible to increase the interrogation time of the atom interferometer beyond 1.4 sec.

Sensitivity Scenario	L [m]	T_{int} [sec]	$\delta\phi_{noise}$ [$1/\sqrt{\text{Hz}}$]	LMT [number n]
AION-10 (initial)	10	1.4	10^{-3}	100
AION-10 (goal)	10	1.4	10^{-4}	1000
AION-100 (initial)	100	1.4	10^{-4}	1000
AION-100 (goal)	100	1.4	10^{-5}	40000
AION-km	2000	5	0.3×10^{-5}	40000


programme will have unprecedented sensitivity to DM with scalar couplings to matter, which cause time variation of fundamental constants such as the electron mass.

Based on: Arvanitaki et al., PRD **97**, 075020 (2018).


← **Linear scalar DM interactions**
Quadratic scalar DM interactions →



Ultra-Light Scalar Dark Matter



AION will be probing new territory in ULD scalar scenarios.
With different configurations of the Atom Interferometer it will be also possible to search for
Axions (pseudo-scalar) and Vector DM candidates!
[studies are ongoing]



References:


- On the Maximal Strength of a First-Order Electroweak Phase Transition and its Gravitational Wave Signal, [1809.08242](#)
- Cosmic Archaeology with Gravitational Waves from Cosmic Strings, [1711.03104](#)
- Probing the pre-BBN universe with gravitational waves from cosmic strings, [1808.08968](#)
- Formation and Evolution of Primordial Black Hole Binaries in the Early Universe, [1812.01930](#)
- Primordial Black Holes from Thermal Inflation, [1903.09598](#)

GW PHYSICS @ AION

AION: Pathway to the GW Mid-(Frequency) Band

AION Corfu Workshop on Standard Model and Beyond


Experimental GW Landscape



AION: Pathway to the GW Mid-(Frequency) Band

AION Corfu Workshop on Standard Model and Beyond

Experimental GW Landscape




Mid-Band currently NOT covered

Mid-band science

- Detect sources BEFORE they reach the high frequency band [LIGO, ET]
- Optimal for sky localization: predict when and where events will occur (for multi-messenger astronomy)
- Search for Ultra-light dark matter in a similar frequency [i.e. mass] range

Gravitational Wave Detection with Atom Interferometry

AION Corfu Workshop on Standard Model and Beyond



Sky position determination

Sky localization
precision:


$$\sqrt{\Omega_s} \sim \left(\text{SNR} \cdot \frac{R}{\lambda} \right)^{-1}$$

Mid-band advantages

- Small wavelength λ
- Long source lifetime (~months) maximizes effective R

Benchmark	$\sqrt{\Omega_s}$ [deg]
GW150914	0.16
GW151226	0.20
NS-NS (140 Mpc)	0.19


Courtesy of Jason Hogan!



Ultimate sensitivity for terrestrial based detectors is achieved by operating 2 (or more) Detectors in synchronisation mode


Ultimate Goal: Establish International Network

AION Corfu Workshop on Standard Model and Beyond



GW Detection & Fundamental Physics - Example


First-Order Electroweak Phase Transition and its Gravitational Wave Signal



arXiv:1809.08242
John Ellis, Marek Lewicki,
José Miguel No


What is the GW signal
of electroweak phase
transition in various
theories beyond
the Standard Model.

Translate strain into dimensionless energy
density $\Omega_{\text{GW}}h^2$ in GWs against frequency




The GW Experimental Landscape: 2030ish

AION Corfu Workshop on Standard Model and Beyond




The GW Experimental Landscape: 2030ish

AION Corfu Workshop on Standard Model and Beyond



The GW Experimental Landscape: 2030ish

AION Corfu Workshop on Standard Model and Beyond



Sensitivity Scenario	L [m]	T_{int} [s]	ϕ [1/√Hz]	LMP [#]
AION-100-today	100	1.4	10^{-3}	100
AION-100-ultimate	100	1.4	10^{-5}	40000
AION-km	2000	5	0.3×10^{-5}	40000
AION-space	4.4×10^7	300	10^{-5}	<1000

List of basic parameters: Lengths of the detector L , interrogation time of the atom interferometer T_{int} , phase noise ϕ , and number of momentum transfers LMP . The choice of these parameters predominately defines the sensitivity of the projection scenarios.

GW Physics: A Few Examples


- **Astrophysical Sources**

- The Black Holes (BH) whose mergers were discovered by LIGO and Virgo have masses up to several tens of solar masses. Many galaxies are known to contain super-massive black holes (SMBHs) with masses in the range between 10^6 and billions of solar masses.
- It is expected that intermediate-mass black holes (IMBHs) with masses in the range 100 to 10^5 solar masses must also exist [6]. There is some observational evidence for IMBHs, and they are thought to have played key roles in the assembly of SMBHs.

- **Cosmological Sources**

- Many extensions of the Standard Model (SM) predict first-order phase transitions in the early Universe. Examples include extended electroweak sectors, effective field theories with higher-dimensional operators and hidden sector interactions.
 - Extended electroweak model with a massive Z' boson
 - Cosmic String Model


Strain Sensitivity & BH Mergers: 2030ish




The AION frequency range is ideal for observations of mergers involving IMBHs, to which LISA and the LIGO/Virgo/KAGRA/ET experiments are relatively insensitive.

Strain Sensitivity & BH Mergers

AION Corfu Workshop on Standard Model and Beyond




Sensitivity of AION-100m for detecting GWs from the mergers of IMBHs at signal-to-noise (SNR) levels ≥ 5 , which extends to redshifts of 1.5 for BHs with masses $\sim 10^4$ solar masses.



Comparison of the sensitivities of AION and other experiments with threshold SNR = 8.

Strain Sensitivity & BH Mergers: Future




The AION frequency range is ideal for observations of mergers involving IMBHs, to which LISA and the LIGO/Virgo/KAGRA/ET experiments are relatively insensitive.

Cosmological GW Sources: Z' Model


Many extensions of the Standard Model (SM) predict first-order phase transitions in the early Universe.

Example: Extended electroweak model with a massive Z' boson



Example of the GW spectrum in a classical scale-invariant extension of the SM with a massive Z' boson compared with various experimental sensitivities. Right panel: Signal-to-noise ratio (SNR) in the parameter plane of the same model for the AION-1km stage.

Cosmological GW Sources: Cosmic Strings



Other possible cosmological sources of GW signals are cosmic strings. These typically give a very broad frequency spectrum stretching across the ranges to which the LIGO/ET, AION/MAGIS, LISA and SKA experiments are sensitive.

The impact of including the change in the number of degrees of freedom as predicted in the Standard Model and clearly shows that probing the plateau in a wide range of frequencies can give us a significant amount of information not only on strings themselves but also on the evolution of the universe.

This way we could probe both SM processes such as the QCD phase transition and BSM scenarios predicting new degrees of freedom or even more significant cosmological modifications such as early matter domination, which would all leave distinguishable features in the GW background.

Other Fundamental Physics

Ultra-high-precision atom interferometry may also be sensitive to other aspects of fundamental physics beyond dark matter and GWs, though studies of such possibilities are still at exploratory stages.

Examples may include:

- *The possibility of detecting the astrophysical neutrinos*
- *Probes of long-range fifth forces.*
- *Constraining possible variations in fundamental constants.*
- *Probing dark energy.*
- Probes of basic physical principles such as foundations of quantum mechanics and Lorentz invariance.

AION-10: 10 METER SIDE CHOSEN TO BE OXFORD


Beecroft building, Oxford Physics

The Beecroft in Oxford is the proposed site, with a backup at RAL (MICE Hall) in case show-stoppers are encountered.



Beecroft building, Oxford Physics

AION Corfu Workshop on Standard Model and Beyond



Ultralow vibration

- All plant isolated
- Thick concrete walls

Adjacent laser lab reserved for AION use

- keel slabs
- $\pm 0.1^\circ\text{C}$ stability
- Isolated mains

Vertical space


- 12m basement to ground floor
- 14.7m floor to ceiling

Stairwell is **not** a fire escape route.


Bakeout room and cleanroom nearby

Beecroft building, Oxford Physics

AION Corfu Workshop on Standard Model and Beyond




← Side view
↓ Plan view





AION-10 site: Beecroft building, Oxford Physics

Beecroft building – brand new, low-vibration laser lab and concrete stairwell



- Detailed planning of support structure by RAL (Engineering), Oxford Physics Technical Services and Liverpool Univ.
- Experienced Project Manager: Roy Preece
- Good site for long-term operation and wide accessibility (also ‘visibility’ and outreach).




Beecroft building laser lab

AION Corfu Workshop on Standard Model and Beyond

Beecroft stairwell: lowest level




Core site: feed through fibre and cables



laser lab (interior): optical table enclosure with laminar air flow and temperature-control installed.

Bake-out room next door




Assembly: extruded aluminium support structure

Scaffolding erected
from ground up.

vacuum pipe;
3.8 m long,
<100 kg.


Remove top layer
after hoisting

10 m




ATOM INTERFEROMETER CONCEPT

Simple Example: Two Atomic Clocks




Simple Example: Two Atomic Clocks

AION Corfu Workshop on Standard Model and Beyond



Phase Noise from the Laser

The phase of the laser is imprinted onto the atom.



*Laser phase is **common** to both atoms – rejected in a differential measurement.*