

# BSM and Top physics with FCC-eh

Oliver Fischer



FCC Physics Workshop  
Liverpool (online)  
10/01/22



## The Large Hadron-Electron Collider at the HL-LHC

LHeC and FCC-he Study Group



P. Agostini *et al.*, [arXiv:2007.14491 [hep-ex]]


LHeC  $E_e = 50 \text{ GeV}$ ,  $\sqrt{s} \simeq 1.2 \text{ TeV}$ ,  $\mathcal{L}_{int} = 1 \text{ ab}^{-1}$ , parallel to HL-LHC

FCC-he  $E_e = 50 \text{ GeV}$ ,  $\sqrt{s} \simeq 3.2 \text{ TeV}$ ,  $\mathcal{L}_{int} = 3 \text{ ab}^{-1}$ , parallel to FCC-hh

## The Large Hadron-Electron Collider at the HL-LHC – section 5.3

5.3	Top Quark Physics . . . . .	129
5.3.1	$Wtq$ Couplings . . . . .	129
5.3.2	Top Quark Polarisation . . . . .	131
5.3.3	Top- $\gamma$ and Top- $Z$ Couplings . . . . .	131
5.3.4	Top-Higgs Coupling . . . . .	133
5.3.5	Top Quark PDF and the Running of $\alpha_s$ . . . . .	133
5.3.6	FCNC Top Quark Couplings . . . . .	134
5.3.7	Other Top Quark Property Measurements and Searches for New Physics .	136
5.3.8	Summary of Top Quark Physics . . . . .	136

# Anomalous $Wtq$ couplings





- ▶ Flagship measurement  $V_{tb}$ ,  $\sigma = 15.3$  pb
- ▶ Generic modification of  $Wtq$  interactions

$$\begin{aligned}\delta\mathcal{L}_{Wtb} = & -\frac{g}{\sqrt{2}}\bar{b}\gamma^\mu V_{tb}(f_1^L P_L - f_1^R P_R)tW_\mu^- \\ & -\frac{g}{\sqrt{2}}\bar{b}\frac{-i\sigma^{\mu\nu}q_\nu}{M_W}(f_2^L P_L - f_2^R P_R)tW_\mu^- + \text{h.c.}\end{aligned}$$

- ▶ SM: all  $f_i$  vanish at tree level.
- ▶ Analysis of  $e^- p \rightarrow \nu\bar{t} + X$  at the detector level including background processes and systematic effects:  
 $\Rightarrow$  Precision of  $\sim 10^{-3}$  for  $f_i^L$  and  $10^{-2}$  for  $f_i^R$  at LHeC.

Mellado *et al.*, *Eur. Phys. J. C* **75** (2015) no.12, 577 [arXiv:1307.1688 [hep-ph]].

# Top- $\gamma$ couplings



- ▶ Photoproduction of  $t\bar{t}$  directly proportional to  $t\bar{t}\gamma$  vertex
- ▶ Generic effective model independent parametrisation:

$$\mathcal{L}_{t\bar{t}\gamma} = e\bar{t} \left( \lambda Q_t \gamma^\mu A_\mu + \frac{1}{4m_t} \sigma_{\mu\nu} V^{\mu\nu} (\kappa_V + i\tilde{\kappa}_V \gamma_5) \right) t$$


with anomalous magnetic and electric dipole moments  $\kappa$  and  $\tilde{\kappa}$

[Bouzas and Larios, Phys. Rev. D \*\*88\*\* \(2013\) no.9, 094007 \[arXiv:1308.5634 \[hep-ph\]\].](#)

- ▶ Recent analysis (anomalous dipole moments) at the detector level incl. several backgrounds and systematics (18%).

[New results from Bouzas and Larios, \[arXiv:2111.04723 \[hep-ph\]\].](#)

# Flavor changing neutral currents with top quarks



- ▶ Single top quark searches are sensitive to FCNC  $tqV$ :

$$\mathcal{L}_{FCNC}^{qV} = \frac{g}{\Lambda_V} \bar{t} \sigma^{\mu\nu} (\lambda_{qV}^L P_L + \lambda_{qV}^R P_R) q V_{\mu\nu} + \text{h.c.}$$

$$q = u, c, V = \gamma, Z$$

- ▶ These couplings also give rise to FCNC top decays:  $t \rightarrow Vq$
- ▶ Process  $e^- p \rightarrow e^- W^+ b + X$
- ▶ Analysis at the detector level yields sensitivity of  $\lambda_{q\gamma} = 0.0025$  and  $\lambda_{qZ} = 0.0037$  at  $2\sigma$  for  $1/\text{ab}$ .

Cakir *et al.*, Nucl. Phys. B **944** (2019), 114640 [arXiv:1809.01923 [hep-ph]].

<b>8</b>	<b>Searches for Physics Beyond the Standard Model</b>	<b>188</b>
8.1	Introduction . . . . .	188
8.2	Extensions of the SM Higgs Sector . . . . .	188
8.2.1	Modifications of the Top-Higgs interaction . . . . .	189
8.2.2	Charged scalars . . . . .	189
8.2.3	Neutral scalars . . . . .	190
8.2.4	Modifications of Higgs self-couplings . . . . .	191
8.2.5	Exotic Higgs boson decays . . . . .	192
8.3	Searches for supersymmetry . . . . .	192
8.3.1	Search for the SUSY Electroweak Sector: prompt signatures . . . . .	193
8.3.2	Search for the SUSY Electroweak Sector: long-lived particles . . . . .	194
8.3.3	R-parity violating signatures . . . . .	195
8.4	Feebly Interacting Particles . . . . .	196
8.4.1	Searches for heavy neutrinos . . . . .	196
8.4.2	Fermion triplets in type III seesaw . . . . .	197
8.4.3	Dark photons . . . . .	199
8.4.4	Axion-like particles . . . . .	200
8.5	Anomalous Gauge Couplings . . . . .	201
8.5.1	Radiation Amplitude Zero . . . . .	202
8.6	Theories with heavy resonances and contact interaction . . . . .	202
8.6.1	Leptoquarks . . . . .	203
8.6.2	$Z'$ mediated charged lepton flavour violation . . . . .	204
8.6.3	Vector-like quarks . . . . .	205
8.6.4	Excited fermions ( $\nu^*, e^*, u^*$ ) . . . . .	206
8.6.5	Colour octet leptons . . . . .	206
8.6.6	Quark substructure and Contact interactions . . . . .	206

## Beyond the Standard Model studies at $ep$

- ▶ **Electron-proton collider** ideal laboratory to study common features of electrons and quarks with EW / VBF production, LQ, multi-jet final states, forward objects
- ▶ **Upside:**
  - Small background (no QCD interaction between e and p)
  - Very low pileup
- ▶ **Downside:** low production rates for new physics processes due to small  $\sqrt{s}$
- ▶ Increased engagement from theory community in recent years, summarised in “chapter 8” (almost 100 articles).



Here: [brief overview over some of the “latest” contributions.](#)



# Searching for charged lepton flavor violation at ep colliders

S. Antusch, A. Hammad and A. Rashed, JHEP **03** (2021), 230 [arXiv:2010.08907 [hep-ph]].

# Lepton flavor violating processes




- ▶ An effective vertex couples incoming electron to a muon or a tau and a neutral scalar or vector boson.
- ▶ Flavor changing physics parametrised via an effective vertex coupling of leptons with Higgs, photon, and Z.
- ▶ Analysis for the LHeC at the detector level.

# Backgrounds: small cross sections, well separable


#	Backgrounds $\tau$ final state	$\sigma_{(LHeC)}[Pb]$
bkg1	$pe^- \rightarrow Z/\gamma^*(\rightarrow \tau^- \tau^+) \nu_l j$	0.0316
bkg2	$pe^- \rightarrow W^\pm(\rightarrow \tau^\pm \nu_\tau) e^- j$	0.2657
bkg3	$pe^- \rightarrow ZZ(\rightarrow \tau^- \tau^+) \nu_l j$	$1.1 \times 10^{-5}$
bkg4	$pe^- \rightarrow Z(\rightarrow \tau^- \tau^+) W^\pm(\rightarrow \tau^\pm \nu_\tau) \nu_l j$	$2.64 \times 10^{-5}$

#	Backgrounds $\mu$ final state	$\sigma_{(LHeC)}[Pb]$
bkg1	$pe^- \rightarrow Z/\gamma^*(\rightarrow \mu^- \mu^+) \nu_l j$	0.0316
bkg2	$pe^- \rightarrow W^\pm(\rightarrow \mu^\pm \nu_\mu) e^- j$	0.2657
bkg3	$pe^- \rightarrow Z/\gamma^*(\rightarrow \tau^- \tau^+ \rightarrow \text{leptons}) \nu_l j$	$9.1 \times 10^{-4}$
bkg4	$pe^- \rightarrow W^\pm(\rightarrow \tau^\pm \nu_\tau \rightarrow \text{leptons}) e^- j$	0.0451
bkg5	$pe^- \rightarrow ZZ(\rightarrow \mu^- \mu^+) \nu_l j$	$1.1 \times 10^{-5}$
bkg6	$pe^- \rightarrow Z(\rightarrow \mu^- \mu^+) W^\pm(\rightarrow \mu^\pm \nu_\mu) \nu_l j$	$2.64 \times 10^{-5}$



Cut-based optimisation of signal-to-background ratio.

# Sensitivity to flavor violation




- ▶ Model independent limits on form factors for LHeC.
- ▶ Recast in specific model, here: sterile neutrinos.
- ▶ Flavor violation proportional to  $|\theta_e \theta_\alpha^*|$

# Exotic Higgs decays into displaced jets at the LHeC



K. Cheung, O. Fischer, Z. S. Wang and J. Zurita, JHEP 02 (2021), 161 [arXiv:2008.09614 [hep-ph]].

# Extending the SM with a complex neutral scalar singlet $S$



- ▶  $S$  can couple to and mix with the SM Higgs field.
- ▶ Physical fields:  $h_1$  ('Higgs'),  $h_2$  with  $m_{h_2} = \mathcal{O}(10)$  GeV.
- ▶  $h_2$  production at LHeC:  $h_1 \rightarrow 2h_2$  with small branching ratio.
- ▶ Decay rate of  $h_2$  suppressed by mixing  $\Rightarrow$  long-lived particle

# Sensitivity



- ▶ Consider only CC Higgs production:  $e^- p \rightarrow \nu_e h_1 j$ .
- ▶  $h_1 \rightarrow 2h_2 \rightarrow 4b$  with two displaced vertices.
- ▶ Analysis at the detector level.
- ▶ From events with  $n_{jet} \geq 5$ , reconstruct  $m_{h_2}$ , require displacement. “Delphes with displacement.” <https://sites.google.com/site/leftrightsep/delphes>.
- ▶ Inclusive backgrounds:  $e^- p \rightarrow \nu_e + n_b b + n_j j + n_\tau \tau$



# Displaced Neutrino Jets at the LHeC

G. Cottin, O. Fischer, S. Mandal, M. Mitra and R. Padhan,  
[arXiv:2104.13578 [hep-ph]].



# Leptoquark $\tilde{R}_2$ and longlived sterile neutrino

$$\mathcal{L}_{LQ} = -Y_{ij} \bar{d}_R^i \tilde{R}_2^a \epsilon^{ab} L_L^{j,b} + Z_{ij} \bar{Q}_L^{i,a} \tilde{R}_2^a N_R^j + \text{H.c.}$$



- ▶ Heavy neutrino  $N$  with mass  $\sim GeV$ ; long lived particle.
- ▶  $\tilde{R}$  with dominant branching into  $qN$  difficult to study at LHC.
- ▶ Can be produced in  $ep$  collisions via  $\tilde{R}$ :  
 $ep \rightarrow \tilde{R} \rightarrow jN$ , with  $N \rightarrow$  displaced fat jet.
- ▶  $5\sigma$  with  $120 \text{ fb}^{-1}$  for  $M_N \sim 10 \text{ GeV}$  and  $\tilde{R}Nq$  coupling  $\sim 0.1$ .
- ▶ Significant improvement from positron-proton scattering.

# Doubly Charged Higgs Production at Future ep Colliders

X. H. Yang and Z. J. Yang, [arXiv:2103.11412 [hep-ph]].

## Extending the SM with a $SU(2)_L$ triplet scalar: $\Delta$

- ▶ Motivation: type II seesaw for neutrino masses:


$$\mathcal{L}_{Y_\Delta} = Y_\Delta \bar{l}^c i \sigma^2 \Delta l + H.c.$$

$$\Rightarrow m_\nu = Y_\Delta \sqrt{2} v_\Delta$$


- ▶ Lepton flavor violating processes  $\tau \rightarrow \bar{l}_i l_j l_k$  and  $\mu \rightarrow \bar{e} e e$  mediated at tree level and constrain  $Y_\Delta$ .
- ▶ Constraints from precision measurements:  $v_\Delta \leq 1$  GeV.
- ▶ LHC searches for doubly charged scalars only stringent when  $H^{\pm\pm} \rightarrow l^\pm l^\pm$  is the dominant decay mode.

cf. also S. Antusch et al., JHEP **02** (2019), 157 [arXiv:1811.03476 [hep-ph]].

# Searching doubly charged scalars at FCC-he




- ▶ Scalar production via vector boson fusion.
- ▶ Search for two doubly (and singly) charged scalars, decaying to  $2SS\mu$  plus jets.
- ▶ Signal: analytical calculation  $\rightarrow$  simulation with vegas.
- ▶ Background:  $e^- p \rightarrow e(\nu_2)t\bar{t}W^\pm j \rightarrow$  Madgraph5.



# Other recent articles

“Search for heavy Majorana neutrinos at electron-proton colliders,” [arXiv:2201.12997 [hep-ph]].

- ▶ By H. Gu and K. Wang.
- ▶ Analysis at detector level using boosted decision tree.
- ▶ Sensitivity similar to lepton-number conserving signatures,
- ⇒ Background free to excellent approximation.



- ▶ A. Jueid, J. Kim, S. Lee and J. Song, “Studies of nonresonant Higgs pair production at electron-proton colliders,” [arXiv:2102.12507 [hep-ph]].
- ▶ K. Cheung and Z. S. Wang, “Physics potential of a muon-proton collider,” [arXiv:2101.10476 [hep-ph]].
- ▶ G. D. Kribs, D. McKeen and N. Raj, “Breaking up the Proton: An Affair with Dark Forces,” Phys. Rev. Lett. **126** (2021) no.1, 011801 [arXiv:2007.15655 [hep-ph]].
- ▶ A. Gutiérrez-Rodríguez, M. A. Hernández-Ruíz, E. Gurkanli, V. Ari and M. Köksal, “Study on the anomalous quartic  $W^+W^-\gamma\gamma$  couplings of electroweak bosons in  $e^-p$  collisions at the LHeC and the FCC-he,” Eur. Phys. J. C **81** (2021) no.3, 210 [arXiv:2005.11509 [hep-ph]].

# Conclusions

- ▶ The LHeC generates a lot of interest in the pheno community.
- ▶ Driving factor: complementary to pp and ee colliders.
- ▶ Opportunities for precision measurements of top physics:
  - ★ Single top and  $t\bar{t}$  production;
  - ★ top couplings to  $\gamma$ ,  $Z$ ,  $W$ , and FCNC interactions.
- ▶ Opportunities for BSM that is hidden at the LHC:
  - ★ Displaced vertices from long lived particles;
  - ★ Lepton flavor violation (electron-tau);
  - ★ Not-too-heavy scalars;
  - ★ GeV-scale bosons.