

Reconstructing the Kinematics of Deep Inelastic Scattering with Deep Learning

Wednesday 4 May 2022 17:10 (20 minutes)

We present a novel method to reconstruct the kinematics of neutral-current deep inelastic scattering (DIS) using a deep neural network (DNN). Unlike traditional methods, it exploits the full kinematic information of both the scattered electron and the hadronic-final state, and it accounts for QED radiation by identifying events with radiated photons and event-level momentum imbalance. The method is studied with simulated events at HERA and the future Electron-Ion Collider (EIC). We will show that the DNN method outperforms all the traditional methods over the full phase space, improving resolution and reducing bias. The DNN-based reconstruction has the potential to extend the kinematic reach of future experiments at the EIC, and thus their discovery potential in polarized and nuclear DIS.

Submitted on behalf of a Collaboration?

No

Authors: NACHMAN, Ben (Lawrence Berkeley National Lab. (US)); BRITZGER, Daniel (Max-Planck-Institut für Physik München); ARRATIA MUNOZ, Miguel Ignacio (Lawrence Berkeley National Lab. (US)); LONG, Owen (University of California Riverside (US))

Presenters: NACHMAN, Ben (Lawrence Berkeley National Lab. (US)); BRITZGER, Daniel (Max-Planck-Institut für Physik München); ARRATIA MUNOZ, Miguel Ignacio (Lawrence Berkeley National Lab. (US)); LONG, Owen (University of California Riverside (US))

Session Classification: WG1: Structure Functions and Parton Densities

Track Classification: WG1: Structure Functions and Parton Densities