

## Heavy Neutrinos at Future Linear $e^+e^-$ Colliders

*Tuesday 3 May 2022 17:10 (20 minutes)*

Neutrinos are probably the most mysterious particles of the Standard Model. The mass hierarchy and oscillations, as well as the nature of their antiparticles, are currently being studied in experiments around the world. Moreover, in many models of New Physics, baryon asymmetry or dark matter density in the universe are explained by introducing new species of neutrinos. Among others, heavy neutrinos of the Dirac or Majorana nature were proposed to solve problems persistent in the Standard Model. Such neutrinos with masses above the EW scale could be produced at future linear  $e^+e^-$  colliders, like the Compact Linear Collider (CLIC) or the International Linear Collider (ILC).

We studied the possibility of observing production and decays of heavy neutrinos in the  $qq$  final state at ILC running at 500 GeV and 1 TeV and CLIC running at 3 TeV. The analysis is based on the WHIZARD event generation and fast simulation of the detector response with DELPHES. Dirac and Majorana neutrinos with masses from 200 GeV to 3.2 TeV are considered. Estimated limits on the production cross sections and on the neutrino-lepton coupling are compared with the current limits coming from the LHC running at 13 TeV, as well as the expected future limits from hadron colliders. Obtained results are stricter than other limit estimates published so far.

### Submitted on behalf of a Collaboration?

No

**Primary authors:** ZARNECKI, Aleksander (University of Warsaw (PL)); Mr MEKALA, Krzysztof; REUTER, Jürgen

**Presenter:** Mr MEKALA, Krzysztof

**Session Classification:** WG6: Future Experiments

**Track Classification:** WG6: Future Experiments