



Canadian Association
of Physicists

Association canadienne
des physiciens et physiciennes

Contribution ID: 3365 Type: **Oral Competition (Graduate Student) / Compétition orale (Étudiant(e) du 2e ou 3e cycle)**

(G*) Force Without Form: Delineating a Disordered Protein Complex with Single-Molecule Fluorescence Spectroscopy

Tuesday, 7 June 2022 09:00 (15 minutes)

Intrinsically disordered proteins (IDPs) play critical roles in regulatory protein interactions, but detailed structural/dynamics characterization of their ensembles remain challenging, both in isolation and they form dynamic 'fuzzy' complexes. Such is the case for mRNA cap-dependent translation initiation, which is regulated by the interaction of the predominantly folded eukaryotic initiation factor 4E (eIF4E) with the intrinsically disordered eIF4E binding proteins (4E-BPs) in a phosphorylation-dependent manner. Fluorescence spectroscopy provides crucial insights into the dimensions and dynamics of IDPs which inform the molecular basis of their function. Single-molecule Förster resonance energy transfer showed that the conformational changes of 4E-BP2 induced by binding to eIF4E are non-uniform along the sequence; while a central region containing both motifs that bind to eIF4E expands and becomes stiffer, the C-terminal region is less affected. Fluorescence anisotropy decay revealed a nonuniform segmental flexibility at different sites along the chain. Dynamic quenching of these fluorescent probes by intrinsic aromatic residues measured via fluorescence correlation spectroscopy report on transient intra- and inter-molecular contacts on ns- μ s timescales. The chain rigidity around sites in the C-terminal region far away from the two binding motifs significantly increased upon binding to eIF4E, suggesting that this region is also involved in the highly dynamic 4E-BP2:eIF4E complex. Our time-resolved fluorescence data paint a sequence-level rigidity map of three states of 4E-BP2 differing in phosphorylation or binding status and distinguish regions that form contacts with eIF4E. We are now conducting single-molecule experiments aimed at resolving site-specific interactions and kinetics of the eIF4E:4E-BP2 complex. Our results constitute an important step towards a mechanistic understanding of the biological function of IDPs via integrative modelling.

Primary author: SMYTH, Spencer

Co-authors: ZHANG, Zhenfu (University of Toronto); Prof. BAH, Alaji (SUNY Upstate Medical University); Mr TSANGARIS, Thomas (University of Toronto); Dr DAWSON, Jennifer (Cleveland Clinic); Prof. FORMAN-KAY, Julie (The Hospital for Sick Children); GRADINARU, Claudiu (University of Toronto)

Presenter: SMYTH, Spencer

Session Classification: T1-1 Advances in Physics in Biology and Medicine Symp.: Protein system dynamics (DPMB) | Symposium sur les progrès en physique dans la biologie et la médecine: dynamique des systèmes de protéines (DPMB)

Track Classification: Symposia Day (Tues. June 7) / Journée de symposiums (mardi, le 7 juin): Symposia Day (DPMB) - Advances in Biological and Medical Physics Symposium