

Contribution ID: **3211** Type: **Oral Competition (Graduate Student)** / **Compétition orale (Étudiant(e) du 2e ou 3e cycle)** 

## (G\*) Inverse Laplace transform of NMR spin-lattice relaxation data

Wednesday 8 June 2022 10:45 (15 minutes)

Traditional NMR data analysis techniques, such as the stretched exponential fit, are used to determine the sample-averaged nuclear spin-lattice relaxation rate  $1/T_1$ . However, they face difficulty when dealing with heterogeneous materials with NMR signals coming from distinct local environments, especially those with large, overlapping distributions of their Knight shifts and  $1/T_1$ .

To overcome this, we perform inverse Laplace transform (ILT) to obtain the histogram  $P(1/T_1)$  of the  $1/T_1$  distribution from the nuclear spin recovery curve M(t). We apply this technique to <sup>63</sup>Cu and <sup>79</sup>Br NQR data of kagome lattice materials herbertsmithite (ZnCu<sub>3</sub>(OD)<sub>6</sub>Cl<sub>2</sub>) and Zn-barlowite (ZnCu<sub>3</sub>(OD)<sub>6</sub>FBr) as well as <sup>19</sup>F NMR data of the latter.

From the  ${}^{63}$ Cu data, we were able to use ILT to observe the gradual emergence of spin singlets with spatially varying excitation gaps below  $\sim$ 30<sup>°</sup>K in both materials. We also performed ILT across the  ${}^{19}$ F NMR spectrum to obtain 3-dimensional ILT-resolved NMR lineshapes, which allowed us to separate the signals coming from two distinct, overlapping sites.

[1] J. Wang et al., Nat. Phys. 17, 1109–1113 (2021)

[2] J. Wang, W Yuan et al., Phys. Rev. Lett. (in press)

Primary author: WANG, Jiaming

Co-authors: YUAN, Weishi; IMAI, Takashi (McMaster University)

**Presenter:** WANG, Jiaming

Session Classification: W1-9 Quantum Magnetism (DCMMP) | Magnétisme quantique (DPMCM)

**Track Classification:** Technical Sessions / Sessions techniques: Condensed Matter and Materials Physics / Physique de la matière condensée et matériaux (DCMMP-DPMCM)