



University  
of Granada

# Solar Atmospheric Neutrinos Searches with the ANTARES Neutrino Telescope

ANTARES-KM3NeT

Dark Ghost-2022

April 1<sup>st</sup>, 2022

On behalf of the ANTARES collaboration

D. Lopez-Coto

S. Navas

J.D. Zornoza


# Solar Atmospheric Neutrinos

ANTARES – Solar Atmospheric Neutrinos

- CRs blocked by the Sun yield  $\nu$  as final state particles.
- The majority of the neutrinos are absorbed in the inner part.
- $\nu$  produced at the solar corona can escape and reach the Earth.
- Important for understanding the solar composition as well as the background for indirect solar DM searches.

Figure from:


J. Edsjö et al JCAP06(2017)033





# The ANTARES Detector

- First undersea Neutrino Telescope.
- Located in the Mediterranean Sea, near Toulon, at 2500 m depth.
- Construction 2006-2008.
- Continuously taking data.
- Switched off on February 2022. Dismantling foreseen by summer 2022.
- 12 lines (885 PMTs)
- 25 storeys/line
- 3 PMTs/storey







# The ANTARES Detector

Median angular resolution, track channel.


- 2 main topologies
  - Track like  $\rightarrow$  From  $\nu_\mu$  and  $\nu_\tau$  CC.
  - Shower like  $\rightarrow$  From all-flavours NC and  $\nu_e$  and  $\nu_\tau$  CC.
- Angular resolution  $< 0.4^\circ$  for  $E_\nu > 10$  TeV).



# The ANTARES Detector



- ANTARES effective area for this analysis.
- Sun tracking taken into account.



# Analysis Outlook



- Only track channel considered ( $\nu_{\mu}$  CC).
- Data taking period from 2008 to 2018 (both included)  $\rightarrow$  lifetime of 3022 days.
- Main background  $\rightarrow$  Atmospheric  $\mu$  and atmospheric  $\nu$ .
- Selection quality cuts to optimize  $SA_{\nu}$  Sensitivity and reject background.
  - $\Lambda > -5.2$ , reconstruction fit parameter
  - $\beta < 1^{\circ}$ , error estimate in the reconstructed angle
  - $\cos\theta > 0 \rightarrow$  upward-going events.
- Unbinned likelihood search.



# Analysis


- Solar Atmospheric Neutrino flux from WimpSim 5.0  
From: **J. Edsjö et al JCAP06(2017)033**
  - 2 Cosmic Ray (CR) models (**H3a** and **GST4**).
  - 2 Solar composition models. (**Ser+Stein** and **Ser+GS98**).
  - Oscillation and Normal Ordering parameters. From global-best fit: **JHEP 01 (2017) 087**
  - Solar Magnetic Field Effect is neglected.

$$\theta_{12} = 33.56^\circ \quad \delta = 261^\circ$$

$$\theta_{13} = 8.46^\circ \quad \Delta m_{21}^2 = 7.5 \cdot 10^{-5} \text{eV}^2$$

$$\theta_{23} = 41.6^\circ \quad \Delta m_{31}^2 = 2.524 \cdot 10^{-3} \text{eV}^2$$


- Sun as a point source, filled disk and ring shape.





# Analysis

- Unbinned Likelihood search. 
$$\mathcal{L}(n_{\text{sig}}) = e^{-(n_{\text{sig}}+n_{\text{bkg}})} \prod_i^N [n_{\text{sig}} \cdot \mathcal{S}(\Psi_{\odot,i}, \beta_i, E_i) + n_{\text{bkg}} \cdot \mathcal{B}(\Psi_{\odot,i}, \beta_i, E_i)]$$
- Signal and Background PDFs from MC weighted events and scrambled data respectively.







# Analysis


$$\mathcal{L}(n_{\text{sig}}) = e^{-(n_{\text{sig}} + n_{\text{bkg}})} \prod_i^N [n_{\text{sig}} \cdot \mathcal{S}(\Psi_{\odot,i}, \beta_i, E_i) + n_{\text{bkg}} \cdot \mathcal{B}(\Psi_{\odot,i}, \beta_i, E_i)]$$

- Likelihood ratio test.

$$\text{TS} = \log_{10} \left( \frac{\mathcal{L}(\hat{n}_{\text{sig}})}{\mathcal{L}(0)} \right)$$

- Natural statistical fluctuations and 15% uncertainty in the number of detected events are included.
- Sensitivity computation.
- 90% CL upper limit computation.

Schematic Sensitivity Calculation





# Analysis


$$\mathcal{L}(n_{\text{sig}}) = e^{-(n_{\text{sig}} + n_{\text{bkg}})} \prod_i^N [n_{\text{sig}} \cdot \mathcal{S}(\Psi_{\odot,i}, \beta_i, E_i) + n_{\text{bkg}} \cdot \mathcal{B}(\Psi_{\odot,i}, \beta_i, E_i)]$$

- Likelihood ratio test.


$$\text{TS} = \log_{10} \left( \frac{\mathcal{L}(\hat{n}_{\text{sig}})}{\mathcal{L}(0)} \right)$$

- Natural statistical fluctuations and 15% uncertainty in the number of detected events are included.
- Sensitivity computation.
- 90% CL upper limit computation.

Schematic Upper-Limit Calculation



# Results



- Event distribution as a function of the angular distance around the source.
- Expected signal magnified for comparison (blue histogram).
- Expected background (green).
- Data (black points).



# Results

- The flux limit is computed as:

$$\frac{d\phi_{\nu_\mu + \bar{\nu}_\mu}^{90\%CL}(E)}{dE} = \frac{\bar{\mu}_{sg}^{90\%CL}}{n_{sg}^{theor}} \cdot \frac{d\phi_{\nu_\mu + \bar{\nu}_\mu}^{theor}(E)}{dE}$$

Where:

$$n_{sg}^{theor} = T_{live} \int \sum_{l \in \nu_\mu, \bar{\nu}_\mu} \left( \frac{d\phi_l^{theor}(E')}{dE'} A_{eff}^l(E') \right) dE'$$

- Is the expected number of signal events for the considered lifetime (3022 days).


-  $n_{sig}^{theor} = 0.36$

- Unblinded results:

-  $\mu_{90} = 3.15 \rightarrow C_{90} \approx 8.6$

- p-value = 0.41

- Base line:
  - H3a + Ser-Stein
  - Sun as a Point Source





# Summary

- 11 years of ANTARES data.
- Unbinned Likelihood Method is used.
- No signal evidence is observed.
- A flux upper limit is established to be  $7 \times 10^{-11}$  [ $\text{TeV}^{-1} \text{ cm}^{-2} \text{ s}^{-1}$ ] at 1 TeV neutrino Energy.
- Pre-print: [arXiv:2201.11642](https://arxiv.org/abs/2201.11642)