

Analysis Update: p_T spectra as a function of R_T for pp collisions at $\sqrt{s} = 5.02$ TeV

Tutor: Dr. Antonio Ortiz Velásquez



Luz Elena Tiscareño Montoya
Paola Vargas Torres

Tuesday, September 13th 2021

1D Unfolding of distributions $N_{\text{ch}}^{\text{TS}}$



Purpose: To obtain a better estimate of the true distribution of the multiplicity of charged particles from the development of experimental distributions.

$N_{\text{ch},\text{true}}^{\text{TS}}$: True multiplicity in the Transverse region.
 $N_{\text{acc}}^{\text{TS}}$: Multiplicity distribution of measured events in the Transverse region.
 $P(N_{\text{acc}}^{\text{TS}} | N_{\text{ch},\text{true}}^{\text{TS}})$: Multiplicity response matrix.





Multiplicity correlation matrices



$$5 < p_{\text{T}}^{\text{leading}} < 40 \text{ GeV}/c$$

Left: 2015 track cuts and right: hybrid track cuts

2D Unfolding of p_T spectra

NEW

First: Apply the tracking efficiency and secondary particle contamination


Second:

(Multiplicity response matrix, $P(N_{\text{acc}}^{\text{TS}} | N_{\text{ch,true}}^{\text{TS}})$)  (Weight of the number of measured particles)


MC closure test


Normal and reverse method

Near Side ¶





Reverse Method




Reverse Method








Reverse Method





Away Side





Reverse Method




Reverse Method








Reverse Method





Transverse Side ¶





Reverse Method




Reverse Method



Reverse Method

Thank you

