



Contribution ID: 118

Type: not specified

## Antiproton Flux and Properties of Elementary Particle Fluxes in Primary Cosmic Rays Measured with the Alpha Magnetic Spectrometer on the ISS

*Wednesday 10 August 2022 16:50 (20 minutes)*

Latest results by AMS on the fluxes and flux ratios of charged elementary particles in the absolute rigidity range from 1 up to 2000 GV reveal unique properties of cosmic charged elementary particles. In the absolute rigidity range  $\sim 60$  to  $\sim 500$  GV, the antiproton flux and proton flux have nearly identical rigidity dependence. This behavior indicates an excess of high energy antiprotons compared with secondary antiprotons produced from the collision of cosmic rays. More importantly, from  $\sim 60$  to  $\sim 500$  GV the antiproton flux and positron flux show identical rigidity dependence. The positron-to-antiproton flux ratio is independent of energy and its value is determined to be a factor of 2 with percent accuracy. This unexpected observation indicates a common origin of high energy antiprotons and positrons in the cosmos.

### Collaboration name

AMS

**Primary author:** WENG, Zhili (Massachusetts Inst. of Technology (US))

**Presenter:** WENG, Zhili (Massachusetts Inst. of Technology (US))

**Session Classification:** Cosmic Rays

**Track Classification:** Cosmic Rays