



Contribution ID: 233

Type: **Parallel Talk**

## Ab initio calculations of structure factors for dark matter searches

*Tuesday 9 August 2022 16:50 (20 minutes)*

We present converged ab initio calculations [1] of structure factors for elastic spin-dependent WIMP scattering off all nuclei used in dark matter direct-detection searches:  $^{19}\text{F}$ ,  $^{23}\text{Na}$ ,  $^{27}\text{Al}$ ,  $^{29}\text{Si}$ ,  $^{73}\text{Ge}$ ,  $^{127}\text{I}$ ,  $^{129}\text{Xe}$ , and  $^{131}\text{Xe}$ . From a set of established two- and three-nucleon interactions derived within chiral effective field theory, we construct consistent WIMP-nucleon currents at the one-body level, including effects from axial- vector two-body currents. We then apply the in-medium similarity renormalization group to construct effective valence-space Hamiltonians and consistently transformed operators of nuclear responses. Combining the recent advances of natural orbitals with three-nucleon forces expressed in large spaces, we obtain basis-space converged structure factors even in heavy nuclei. Generally, results are consistent with previous calculations, but in certain cases can differ by as much as 80-90% at low momentum transfer.

[1]. Ab initio structure factors for spin-dependent dark matter direct detection. B.S. Hu, et al. Phys. Rev. Lett. 128 (2022) 072502. arXiv:2109.00193.

### Collaboration name

**Primary author:** Dr HU, Baishan (TRIUMF)

**Presenter:** Dr HU, Baishan (TRIUMF)

**Session Classification:** Dark Matter

**Track Classification:** Dark Matter