

Neutrino point source searches for dark matter spikes black hole remnants of the first stars under the lamp shade



SISSA



Patrick Stengel

SISSA

May 10, 2022



2202.01126 with Katherine Freese, Irina Galstyan and Pearl Sandick

Distribution depends on star formation history, BH mergers

Termination of first star formation

- More minihalos become sufficiently massive to host Pop III.1 stars over time
- History is not well constrained, assume benchmark termination redshifts $z_f = 20, 15, 10$




Figure: 1008.3552

BH mergers can destroy spikes

- Fraction of suitable minihalos $f_{DS} = f_{DS}^0 (1 - f_{\text{merged}})$ which host spikes
- For $M_{BH} \lesssim 10^3 M_{\odot}$ or $f_{DS}^0 \ll 1$ mergers are not relevant

Total number of spikes for $f_{DS} = 1$

$$\int d^3\mathbf{R} N_{\text{sp}}^{z_f}(R) = 409, 7983, 12416$$

Density profiles and annihilation rate for DM spikes

Density of spike depends on M_{BH}

- More DM contracts around more massive central objects
- Standard models of stellar evolution suggest the collapse of fusion-powered Pop III.1 stars yields 10 , $10^2 M_{\odot}$ BHs
- Extended Dark Star phase powered by DM annihilation can yield 10^3 , 10^4 , $10^5 M_{\odot}$ BHs






Figure: 100 GeV WIMPs around IMBH formed in $10^6 M_{\odot}$ subhalo at $z = 15$

$$\Gamma = \frac{\langle \sigma v \rangle}{2m_{\chi}^2} \int_{r_{\min}}^{r_{\max}} dr 4\pi r^2 \rho_{DM}^2$$

For spike 10 pc away, $> 99\%$ of flux from within 0.01° , i.e. $r \lesssim 0.002$ pc

Calculate the neutrino signal from DM spikes



Observable neutrino flux


- Normalization depends on m_χ , M_{BH} , $z_f = 20, 15, 10$
- Shape from source after vacuum oscillations
- Depends on $\nu_{\ell'}$ flavor and annihilation channel

Plot different channels for fixed Γ and D

$$\frac{d\Phi}{dE_\nu} = \frac{\Gamma}{4\pi D^2} \sum_{\ell=e,\mu,\tau} P(\nu_\ell \rightarrow \nu_{\ell'}) \frac{dN}{dE_\nu}$$


$$\ell' = \mu \text{ for } \rightarrow \bar{b}b, W^+W^-, \tau^+\tau^-, \mu^+\mu^-$$

PS searches sensitive to DM spikes under the lamp shade







D_{\min} surface above the galactic plane, fix $z_f = 10$ and $m_\chi = 5 \text{ TeV}$

- **Line** from GC to solar system
- Compare to spherical shells from $N_{\text{sp}}^{10}(R)$, random (θ, ϕ)





N_{sp} contained within D_{\min} surface

$$f_{\text{DS}} \int d^3\mathbf{s} N_{\text{sp}}^{z_f}(\mathbf{s} < D_{\min}) < 1$$

Limits on f_{DS} given lack of confirmed PS by AN or IC

Neutrino PS searches ideal for detecting DM spikes



Probe nature of DM, substructure and models of stellar evolution

- $f_{\text{DS}} \lesssim 10\%$ for 5 TeV WIMPs around $\gtrsim 10^3 M_{\odot}$ BHs
- Robust to BH mergers and tidal disruption near GC
- Complement γ -ray limits

Interesting (?) ideas for future work

- KM3Net combines visibility of AN with larger exposure of IC
- Consider heavier (non-thermal) DM for $E_{\nu} \gtrsim 100$ TeV
- Diffuse emission around SMBHs

N-body simulations suggest there are many $\sim 10^6 M_{\odot}$ halos




Figure: DM density in 800 kpc of VL-II, insets show inner 40 kpc (0805.1244)

Thermal WIMP production motivates ID searches





Figure: 1003.0904

EW coupling \Rightarrow weak scale mass

Can look for products of WIMP dark matter annihilation today



Thermal freeze-out relic abundance

$$\Omega_\chi h^2 \sim 0.1 \left(\frac{1 \text{ pb}}{\langle \sigma_{AV} \rangle} \right)$$

Interact around the detector, measure charged products





Figure: 1805.11112

Estimate number of events for a source flux and declination

Event rate depends on effective area

$$\frac{dN_{EV}}{dt} = \int dE_\nu A_{\text{eff}}(E_\nu, \delta) \frac{d\Phi}{dE_\nu}$$

$$A_{\text{eff}} = V_{\text{gen}} \times \frac{N_{\text{det}}}{N_{\text{gen}}} \times \rho N_A \times \sigma \times P_{\text{Earth}}$$



Calculated from event simulation

- Simulated target volume, detection efficiency
- Target nucleon density
- ν -nucleon cross section, absorption in Earth

Visibility of sources below horizon

$$= \begin{cases} 0, & \text{if } |\delta| > \pi/2 - |\phi|, \delta \cdot \phi > 0 \\ V_\phi(\delta), & \text{if } |\delta| \leq \pi/2 - |\phi| \\ 1, & \text{if } |\delta| > \pi/2 - |\phi|, \delta \cdot \phi < 0 \end{cases}$$

Neutrino telescopes typically only most sensitive for $\rightarrow \nu\nu$




Figure: 1912.09486

γ -ray experiments more sensitive to visible channels





Figure: ANTARES limits on WIMP annihilation in GC (1912.05296)

Limits on annihilation weaken as final state becomes more hadronic



γ -ray signals are easier to detect

- GC limits 10-100 stronger
- Limits from dSphs can exclude $\lesssim 100$ GeV thermal WIMP
- What about DM spikes?

Density profiles and distribution of subhalo masses




Figure: Compare density of **MW halo** to 8 larger **subhalos**. $\gtrsim 40\text{k}$ total evenly distributed by decade $10^6 - 10^9 M_{\odot}$




Figure: Peak height for circular velocity $\propto \sqrt{M^{\text{halo}}}$ of subhalos contained within **402**, 100, 50 kpc of galactic center

Minihalos contract around BH remnants of the first stars

Criteria for Pop III.1 star formation

$$M^{\text{halo}} \gtrsim 10^5 M_{\odot} \left(\frac{1+z}{31} \right)^{-2}$$

$\gtrsim 10^2 M_{\odot}$ star collapses into IMBH

- NFW for initial DM (85%) and baryon (15%) densities
- Gravitational potential pulls DM into central object
- Estimate adiabatic contraction following Blumenthal et al. (1986) prescription




Figure: Comparison of SHM to **DM spike** formed around SMBH, with **central density cutoff** $\simeq m_{\chi} / \langle \sigma v \rangle t_{BH}$ (astro-ph/0504422)

Search for clusters in $1^\circ \times 1^\circ$ grid across visible sky






Figure: 1706.01857

Limits on PS flux normalization Φ_0 for E_ν^{-2} spectrum



Signal events = Acceptance \times Φ_0

$$A_S = \Phi_0^{-1} T_{\text{exp}} \int dE_\nu A_{\text{eff}} \frac{d\Phi}{dE_\nu}$$

Focus on $E_\nu \lesssim 100 \text{ TeV}$ for WIMPs

- AN better visibility for $\delta \lesssim 0^\circ$
- IC very sensitive for $\delta \gtrsim 0^\circ$
- Need N_{EV} limits to estimate sensitivity to DM spikes

Effective area after track (shower) selection cuts








Figure: IceCube 1910.08488

Compare to AN and IC limits in different sky regions





EXi	δ Range
AN1	$[-90^\circ, -45^\circ]$
AN2	$[-45^\circ, 0^\circ]$
AN3	$[0^\circ, 45^\circ]$
IC1	$[-30^\circ, -5^\circ]$
IC2	$[-5^\circ, 0^\circ]$
IC3	$[0^\circ, 30^\circ]$
IC4	$[30^\circ, 60^\circ]$
IC5	$[60^\circ, 90^\circ]$







Number of TR or SH events from a DM spike in EXi

$$N_{\text{EV}}^{\text{EX}i} = \sum_{j \in \text{EX}i} T_{\text{exp}}^j \sum_{\ell' \in \text{EV}} \int_{E_{\text{th}}}^{m_\chi} dE_\nu \frac{d\Phi_{f \rightarrow \nu \ell'}}{dE_\nu} A_{\ell' \text{EV}}^j(E_\nu)$$



Minimum distance to have not been detected as a PS



Minimum distance to have not been detected as a PS



Combine D_{\min} from AN and IC sky regions







D_{\min} surface across full sky for
 $M_{BH} = 10^3 M_{\odot}$ $\chi\chi \rightarrow W^+W^-$
 $m_{\chi} = (500 \text{ GeV})$ and 5 TeV





Compare to $N_{\text{sp}}^{\text{zf}}$ with α towards GC

- GC visible for larger M_{BH}
- Many spikes away from GC

D_{\min} surface compared to $N_{\text{sp}}^{Z_f}$ at different α



Limits on f_{DS} given lack of confirmed PS by AN or IC



Compare to gamma-ray limits from Fermi-LAT

