

Probing ν_R -philic Z' at the DUNE near detector

Garv Chauhan

Centre for Cosmology, Particle Physics and Phenomenology (CP3),
UC Louvain, Belgium

Based on **G.C.**, Bhupal Dev (WUSTL) and Xun-Jie Xu (IHEP-CAS) [arXiv: 2204.11876]

PHENO 2022
University of Pittsburgh
May 9, 2022



- The discovery of neutrino oscillations implies non-zero neutrino masses.

- The discovery of neutrino oscillations implies non-zero neutrino masses.
- Monumental progress to understand neutrino mixing paradigm but yet to understand the neutrino mass mechanism.

- The discovery of neutrino oscillations implies non-zero neutrino masses.
- Monumental progress to understand neutrino mixing paradigm but yet to understand the neutrino mass mechanism.
- The minimal scenario includes the introduction of right-handed neutrinos (ν_R).

$$\mathcal{L}_{mass} \supset m_D \nu_L \nu_R + \frac{M_R}{2} \nu_R \nu_R + h.c.$$

- The discovery of neutrino oscillations implies non-zero neutrino masses.
- Monumental progress to understand neutrino mixing paradigm but yet to understand the neutrino mass mechanism.
- The minimal scenario includes the introduction of right-handed neutrinos (ν_R).

$$\mathcal{L}_{mass} \supset m_D \nu_L \nu_R + \frac{M_R}{2} \nu_R \nu_R + h.c.$$

- ν_R 's are motivated BSM candidates - ν masses, dark matter and matter-antimatter asymmetry (η_B)

- Since ν_R is dark, it might have abundant new interactions beyond the SM.

- Since ν_R is dark, it might have abundant new interactions beyond the SM.
- In this talk, I consider the possibility where ν_R is charged under a hidden $U(1)_R$.

- Since ν_R is dark, it might have abundant new interactions beyond the SM.
- In this talk, I consider the possibility where ν_R is charged under a hidden $U(1)_R$.
- Although SM particles are uncharged under $U(1)_R$, this Z' can still interact through loop-level couplings.

- Since ν_R is dark, it might have abundant new interactions beyond the SM.
- In this talk, I consider the possibility where ν_R is charged under a hidden $U(1)_R$.
- Although SM particles are uncharged under $U(1)_R$, this Z' can still interact through loop-level couplings.
- Here, couplings to neutrinos are generically larger than its loop-induced couplings to electrons.

- Since ν_R is dark, it might have abundant new interactions beyond the SM.
- In this talk, I consider the possibility where ν_R is charged under a hidden $U(1)_R$.
- Although SM particles are uncharged under $U(1)_R$, this Z' can still interact through loop-level couplings.
- Here, couplings to neutrinos are generically larger than its loop-induced couplings to electrons.
- We'll explore how this feature is useful for probing this scenario at DUNE.

- We consider Z' coupled to ν_R with relevant lagrangian,



$$\mathcal{L} \supset g' Z'_\mu \nu_R \gamma^\mu \nu_R + m_D \nu_L \nu_R + \frac{M_R}{2} \nu_R \nu_R + \dots$$

How dark is this Dark Photon ?

- We consider Z' coupled to ν_R with relevant lagrangian,

$$\mathcal{L} \supset g' Z'_\mu \nu_R \gamma^\mu \nu_R + m_D \nu_L \nu_R + \frac{M_R}{2} \nu_R \nu_R + \dots$$


- In absence of kinetic mixing with SM, this Z' can interact with SM particles through these loop-level diagrams,




$$g_f^{(W)} = \frac{\sqrt{2} G_F m_D^2}{8\pi^2} g', \quad g_f^{(Z)} = \frac{\sqrt{2} G_F m_D^2}{8\pi^2} g' Q_f^{(Z)}.$$

where $Q_f^{(Z)} \equiv I_3 - Q_{\text{em}} s_W^2$.

- DUNE is a long-baseline neutrino experiment, with a high intensity neutrino beam originating at Fermilab with a massive far detector (1300 km away) in South Dakota.




- DUNE is a long-baseline neutrino experiment, with a high intensity neutrino beam originating at Fermilab with a massive far detector (1300 km away) in South Dakota.



- The far detector consists of 4 massive liquid argon detectors each about 10 kilotons.

- DUNE is a long-baseline neutrino experiment, with a high intensity neutrino beam originating at Fermilab with a massive far detector (1300 km away) in South Dakota.




- The far detector consists of 4 massive liquid argon detectors each about 10 kilotons.
- Uses liquid argon time projection chamber (LArTPC) technology, which provide excellent particle identification and energy measurements.

- The ND will measure the initial unoscillated ν_μ and ν_e energy spectra.


DUNE Near Detector (ND)

- The ND will measure the initial unoscillated ν_μ and ν_e energy spectra.
- Comparing the energy spectra before oscillation and again at the far site allows to disentangle different energy-dependent effects and systematics.



DUNE Near Detector (ND)


- The ND will measure the initial unoscillated ν_μ and ν_e energy spectra.
- Comparing the energy spectra before oscillation and again at the far site allows to disentangle different energy-dependent effects and systematics.



- The ND consists of

DUNE Near Detector (ND)


- The ND will measure the initial unoscillated ν_μ and ν_e energy spectra.
- Comparing the energy spectra before oscillation and again at the far site allows to disentangle different energy-dependent effects and systematics.



- The ND consists of
 - a LArTPC called ArgonCube

DUNE Near Detector (ND)

- The ND will measure the initial unoscillated ν_μ and ν_e energy spectra.
- Comparing the energy spectra before oscillation and again at the far site allows to disentangle different energy-dependent effects and systematics.




- The ND consists of
 - a LArTPC called ArgonCube
 - a high-pressure gaseous argon TPC (HPgTPC) surrounded by an electromagnetic calorimeter (ECAL) in a 0.5 T magnetic field

- Proton beam can have upto 120 GeV of energy

- Proton beam can have upto 120 GeV of energy
- We assume data collection of 5 years each in ν (“forward horn current”) and $\bar{\nu}$ (“reverse horn current”) modes.

- Proton beam can have upto 120 GeV of energy
- We assume data collection of 5 years each in ν (“forward horn current”) and $\bar{\nu}$ (“reverse horn current”) modes.
- DUNE produces a large flux of charged mesons (mostly π^\pm and K^\pm) that decay leptonically, leading to a large flux of SM neutrinos.



- The effective couplings of Z' to normal matter and neutrinos:

$$\mathcal{L} = \bar{\psi} [g_{eL}\gamma^\mu P_L + g_R\gamma^\mu P_R] Z'_\mu \psi + \bar{\psi}_\nu [g_\nu\gamma^\mu P_L] Z'_\mu \psi_\nu,$$

- The effective couplings of Z' to normal matter and neutrinos:


$$\mathcal{L} = \bar{\psi} [g_{eL}\gamma^\mu P_L + g_R\gamma^\mu P_R] Z'_\mu \psi + \bar{\psi}_\nu [g_\nu\gamma^\mu P_L] Z'_\mu \psi_\nu,$$

- For later use, we define

$$g \equiv \sqrt{g_L^2 + g_R^2},$$

and

$$r \equiv \frac{g_\nu}{g_e}, \quad (g_L, g_R) = (\cos \beta, \sin \beta)g_e.$$



- Differential cross section for elastic neutrino-electron scattering including both the SM and the new physics contributions:

$$\frac{d\sigma}{dT} = \frac{2m_e G_F^2}{\pi} \left[c_L^2 + c_R^2 \left(1 - \frac{T}{E_\nu} \right)^2 - c_L c_R \frac{m_e T}{E_\nu^2} \right],$$

where

$$c_L = c_L^{(\text{SM})} + \frac{g_{eL} g_\nu}{2\sqrt{2} G_F (2m_e T_e + m_{Z'}^2)}, \quad c_L^{(\text{SM})} = -\frac{1}{2} + s_W^2 + \delta_{\alpha e},$$
$$c_R = c_R^{(\text{SM})} + \frac{g_{eR} g_\nu}{2\sqrt{2} G_F (2m_e T_e + m_{Z'}^2)}, \quad c_R^{(\text{SM})} = s_W^2.$$

- Differential cross section for elastic neutrino-electron scattering including both the SM and the new physics contributions:

$$\frac{d\sigma}{dT} = \frac{2m_e G_F^2}{\pi} \left[c_L^2 + c_R^2 \left(1 - \frac{T}{E_\nu} \right)^2 - c_L c_R \frac{m_e T}{E_\nu^2} \right],$$

where


$$c_L = c_L^{(\text{SM})} + \frac{g_{eL} g_\nu}{2\sqrt{2} G_F (2m_e T_e + m_{Z'}^2)}, \quad c_L^{(\text{SM})} = -\frac{1}{2} + s_W^2 + \delta_{\alpha e},$$



$$c_R = c_R^{(\text{SM})} + \frac{g_{eR} g_\nu}{2\sqrt{2} G_F (2m_e T_e + m_{Z'}^2)}, \quad c_R^{(\text{SM})} = s_W^2.$$

- The event rate of elastic neutrino-electron scattering at the detector is computed by:


$$\frac{dN}{dT} = N_e \lambda_{\text{POT}} \int \Phi(E_\nu) \frac{d\sigma(T, E_\nu)}{dT} \Theta(T_{\text{max}} - T) dE_\nu,$$

- assuming $m_{Z'} = 100$ MeV and $\sqrt{g_e g_\nu} = 10^{-4}$





- At the neutrino production site of DUNE, Z' can be produced from the proton beam striking the target. Due to its weak loop-level couplings to SM fermions, the produced Z' boson can be long-lived.
- assuming $r = 100$,



- The number of events are calculated as

$$N_{\text{det.}} = \int dp_{Z'} \frac{dN_{\text{prod.}}(p_{Z'})}{dp_{Z'}} P_{\text{decay}}(p_{Z'}) \text{BR}_{Z' \rightarrow \text{vis.}},$$

$$P_{\text{decay}} = e^{-L_1/L_{Z'}} \left(1 - e^{-L_2/L_{Z'}} \right)$$

- The number of events are calculated as

$$N_{\text{det.}} = \int dp_{Z'} \frac{dN_{\text{prod.}}(p_{Z'})}{dp_{Z'}} P_{\text{decay}}(p_{Z'}) \text{BR}_{Z' \rightarrow \text{vis.}},$$


$$P_{\text{decay}} = e^{-L_1/L_{Z'}} \left(1 - e^{-L_2/L_{Z'}} \right)$$

- The visible decay width is

$$\text{BR}_{Z' \rightarrow \text{vis.}} \equiv 1 - \frac{\Gamma_{Z' \rightarrow \nu \bar{\nu}}}{\Gamma_{Z'}}.$$


when $g_\nu \gg g_e$ and g_q , we have $\text{BR}_{Z' \rightarrow \text{vis.}} \ll 1$


- The sensitivity reach of DUNE MPD to the ν_R -philic Z' with loop-induced couplings. The results depend on the ratio r




Combined Results

- For $r = 1$, DUNE-MPD exhibits a significant advantage over other beam dump experiments in the mass range $0.1\text{GeV} \lesssim m_{Z'} \lesssim 1\text{GeV}$.





- For larger r such as $r = 10$ or 100 , DUNE-LArTPC will be able to generate the leading constraints, exceeding collider bounds from BaBar, LHCb, etc.



- Hidden $U(1)$ symmetries in ν_R sector give rise to dark gauge boson : ν_R -philic Z' .
- Loop-suppressed couplings to SM and larger couplings to ν , neutrino experiments are the most suited to probe this scenario.
- We consider two complementary near DUNE detectors LArTPC and DUNE-MPD (HPgTPC); could be sensitive to Z' signals via elastic ν - e scattering and via Z' decay.
- Larger ν couplings lead to higher elastic ν - e scattering rates in DUNE-LArTPC but make Z' decay less visible in DUNE-MPD due to the enhanced invisible decay width.
- Excellent prospect of DUNE probing new physics hidden in the sector of ν_R .

Thank you!

Additional Slide

- The ArgonCube shares same aspects of form and functionality with the FD, reduces sensitivity to nuclear effects and detector-driven systematic uncertainties in extracting the oscillation signal at the FD.
- Muons with momentum higher than 0.7 GeV/c will not be contained in the LArTPC volume. Since muon momentum is critical to determining the incoming neutrino's energy, a magnetic spectrometer is needed downstream of the LArTPC to measure the momentum and charge of the muons i.e. MPD.
- Both ArgonCube and MPD can move off-axis relative to the beam, providing access to different neutrino energy spectra.

