

Gamma-ray emission from young supernova remnants in dense, structured circumstellar environments

Monday 10 October 2022 16:45 (15 minutes)

Supernova remnants are known to accelerate cosmic rays from the detection of non-thermal emission of radio waves, X-rays, and gamma rays. The presence of cut-offs in the gamma-ray spectra of several young SNRs led to the idea that the highest energies might only be achieved during the very initial stages of a remnant's evolution. Unfortunately, the gamma-ray luminosity is assumed to peak in the first weeks after the Supernova explosion where strong $\gamma\gamma$ -absorption attenuates the observable signal. Here, we investigate to which extent the interaction of SNR-shocks with dense structures in the medium around red supergiant (RSG) and luminous blue variable (LBV) stars can boost the gamma-ray emission months to years after the explosion.

We use the time-dependent acceleration code RATPaC to study the acceleration of cosmic rays in supernovae expanding into dense environments around massive stars. We performed spherically symmetric 1-D simulations in which we simultaneously solve the transport equations for cosmic rays, magnetic turbulence, and the hydrodynamical flow of the thermal plasma in the test-particle limit.

We investigated typical parameters of the circumstellar medium (CSM) in the freely expanding winds around RSG and LBV stars and added dense structures that arise from episodes of highly-enhanced mass-loss in case of LBV or photoionization-shells in the case of RSG progenitors.

We find that the interactions with the dense structures happens typically after a few months for LBV progenitors and a few years for RSG progenitors. During the interaction stage, the $\gamma\gamma$ -absorption by photons emitted from the Supernova's photosphere became negligible. The gamma-ray luminosity of the interacting SNRs can surpass the internal/unabsorbed peak-luminosity that arises shortly after the explosion. As a consequence, the observable flux can be considerably higher compared to the signal expected shortly after the explosion where $\gamma\gamma$ -absorption is important and where most gamma-ray observatories search for transient signals from these Supernovae.

Track

SNR/PWNe

Primary author: B., Robert

Co-authors: SUSHCH, Iurii (North-West University); Dr MACKEY, Jonathan (Dublin Institute for Advanced Studies)

Presenter: B., Robert

Session Classification: Parallel 2