

Double Beta Decay as a Probe of New Physics


Frank Deppisch

`f.deppisch@ucl.ac.uk`


University College London

Dirac versus Majorana


- ▶ Origin of neutrino masses beyond the Standard Model
- ▶ Two possibilities to define neutrino mass



Dirac mass analogous to other fermions but with $m_\nu / \Lambda_{EW} \approx 10^{-12}$ couplings to Higgs



Majorana mass, using only a left-handed neutrino
 → Lepton Number Violation



Dirac versus Majorana

- ▶ Origin of neutrino masses beyond the Standard Model
- ▶ Crucial role of total lepton number L symmetry
 - Arises accidentally as global $U(1)_L$ in SM from particle content and gauge symmetry
 - L broken non-perturbatively but $B - L$ conserved
 - Global symmetries expected to be broken gravitational effects?

$$m_\nu \approx \frac{v^2}{M_{\text{Planck}}} \approx 10^{-5} \text{ eV}$$

- Too small to explain oscillations but too large as subdominant splitting
- Connection to matter–antimatter asymmetry

Beta Decays and Neutrinos

► Single beta decay

$$(A, Z) \rightarrow (A, Z + 1) + e^- + \bar{\nu}_e$$

- Kinematic neutrino mass measurement


► Allowed double beta ($2\nu\beta\beta$) decay

$$(A, Z) \rightarrow (A, Z + 2) + 2e^- + 2\bar{\nu}_e$$

► Neutrinoless double beta ($0\nu\beta\beta$) decay

$$(A, Z) \rightarrow (A, Z + 2) + 2e^-$$

- Violation of lepton number
- Mediated by Majorana neutrinos
- Alternatives:
 - $0\nu\beta^+\beta^+$: $(A, Z) \rightarrow (A, Z - 2) + 2e^+$
 - $0\nu\beta^+EC$: $(A, Z) + e^- \rightarrow (A, Z - 2) + e^+$
 - $0\nuECEC$: $(A, Z) + 2e^- \rightarrow (A, Z - 2)$




Neutrinoless Double β Decay

► Half-life

$$T_{1/2}^{-1} = |\mathbf{m}_{\beta\beta}|^2 \mathbf{G}^{0\nu} |\mathbf{M}^{0\nu}|^2$$

► Particle Physics



$$\mathcal{A}_{\mu\nu}^{lep} = \frac{1}{4} \sum_{i=1}^3 U_{ei}^2 \gamma_\mu (1 + \gamma_5) \frac{\cancel{q} + m_{\nu_i}}{q^2 - m_{\nu_i}^2} \gamma_\nu (1 - \gamma_5) \approx \frac{\gamma_\mu (1 + \gamma_5) \gamma_\nu}{4q^2} \sum_{i=1}^3 U_{ei}^2 m_{\nu_i} \rightarrow \mathbf{m}_{\beta\beta}$$


► Atomic Physics

- Leptonic phase space $\mathbf{G}^{0\nu} \propto Q^5$

► Nuclear Physics


- Nuclear transition matrix element $\mathbf{M}^{0\nu} \approx 1$
but large uncertainties, factor 2–3

$$\frac{10^{25} \text{ y}}{T_{1/2}} \approx \left(\frac{|m_{\beta\beta}|}{\text{eV}} \right)^2$$




Three Active Neutrinos

- ▶ Effective $0\nu\beta\beta$ Mass



Three Active Neutrinos

► Current Best Limit from KamLAND-Zen (arXiv:2203.02139)



Theory Predictions:

- (a) Minimal seesaw with texture zeros, Harigaya, Ibe, Yanagida, PRD 86 (2012) 013002
- (b) Modular $A(4)$ in large volume limit, Asaka, Heo, Yoshida, PLB 811 (2020) 135956
- (c) $U(1)_{L_e-L_\mu}, U(1)_{L_\mu-L_\tau}, U(1)_{B-L}$ gauge symmetries, Asai, EPJC 80 (2020) 76

Sterile Neutrinos


- ▶ Masses lighter than ≈ 100 MeV

$$|m_{\beta\beta}| = |c_{12}^2 c_{13}^2 m_{\nu_1} + s_{12}^2 c_{13}^2 m_{\nu_2} e^{i\phi_{12}} + s_{13}^2 m_{\nu_3} e^{i\phi_{13}} + s_{14}^2 m_{\nu_4} e^{i\phi_{14}} + \dots|$$


- ▶ Masses heavier than ≈ 100 MeV

$$\mathcal{A}_{\mu\nu}^{lep} = \frac{1}{4} \sum_{i=1}^3 V_{ei}^2 \gamma_\mu (1 + \gamma_5) \frac{\cancel{q} + M_{N_i}}{q^2 - M_{N_i}^2} \gamma_\nu (1 - \gamma_5) \approx \frac{-\gamma_\mu (1 + \gamma_5) \gamma_\nu}{4} \sum_{i=1}^3 \frac{V_{ei}^2}{M_{N_i}} \rightarrow \left\langle \frac{1}{M_N} \right\rangle_{\beta\beta}$$

- ▶ Short-distance on nuclear scale




Sterile Neutrinos




New Physics and $0\nu\beta\beta$



► Plethora of New Physics scenarios



$$T_{1/2}^{-1} = \epsilon_{NP}^2 G_{NP}^{0\nu} |M_{NP}^{0\nu}|^2$$



Left–Right Symmetry



R-Parity
Violating SUSY

Extra Dimensions




Majorons

Leptoquarks

...

New Physics and $0\nu\beta\beta$

► Examples in Left-Right Symmetry





$$\epsilon_{V-A}^{V+A} = \sum_{i=1}^3 U_{ei} W_{ei} \tan \zeta_W$$

$$\approx \frac{10^{-9}}{(\Lambda/10 \text{ TeV})^3}$$

$$\epsilon_3^{RRZ} = \sum_{i=1}^3 V_{ei}^2 \frac{m_p}{m_N} \frac{m_W^4}{m_{W_R}^4}$$

$$\approx \frac{10^{-8}}{(\Lambda/1 \text{ TeV})^5}$$





► $0\nu\beta\beta$ probes LNV at the TeV scale and above

Heavy New Physics

FFD, Graf, Iachello, Kotila, PRD 102 (2020)

- ▶ Limits on short-range operators
 - NMEs from IBM-2 with $g_A = 1.0$ and short-range correlations in Argonne parametrization






Pion-mediated contributions

- ▶ R-parity violating SUSY (Faessler, Kovalenko, Simkovic, Schwieger, Phys.Rev.Lett. 78 (1997) 183)
- ▶ Chiral EFT with Pion operators from Lattice QCD (Cirigliano, Dekens, de Vries, Graesser, Mereghetti, JHEP 1812 (2018) 097)

Exotic Particle Emission

Cepedello, FFD, González, Hati, Hirsch, PRL 122 (2019)

- ▶ Majoron(-like) J emission
- ▶ Majoron-like ϕ emission assisted by RH current
- ▶ Electron energy distribution



Exotic Particle Emission



Cepedello, FFD, González, Hati, Hirsch, PRL 122 (2019)

- ▶ Majoron(-like) J emission
- ▶ Majoron-like ϕ emission assisted by RH current
- ▶ Sensitivity to Left–Right symmetric model with Dirac neutrinos

$$\frac{T_{1/2}^{\text{Xe}}}{10^{25} \text{ y}} \approx \left(\frac{1.4 \times 10^{-4}}{g_R^2 \kappa y_N y_\nu} \right)^2 \left(\frac{m_{W_R}}{25 \text{ TeV}} \right)^4 \left(\frac{m_N}{100 \text{ MeV}} \right)^4$$

- ▶ Searched for in EXO-200
(Phys.Rev.D 104 (2021) 11, 112002)

$$T_{1/2}^{Xe} > 4 \times 10^{24} \text{ y}$$




New Physics in $2\nu\beta\beta$

Bolton, FFD, Graf, Simkovic, PRD 103 (2021)

- ▶ Sterile neutrino search through energy endpoint (also Agostini, Bossio, Ibarra, Marcano, PLB 815 (2021))

- Emission of one sterile neutrino in double beta decay: $\nu N \beta\beta$
- Same principle as endpoint searches in single β decays
- Observed limit at GERDA: $|V_{eN}|^2 < 0.013$ (Krause, NDM22)




New Physics in $2\nu\beta\beta$

FFD, Graf, Simkovic, PRL 125 (2020)

- ▶ Lepton-number conserving right-handed currents
 - Exotic charged currents probed e.g.
 - in neutron and single β decay
 - at LHC in $pp \rightarrow eX + MET$
 - Limits on RH currents

$$\frac{G_F \cos \theta_C}{\sqrt{2}} ((1 + \delta_{SM} + \epsilon_{LL}) j_L^\mu J_{L\mu} + \epsilon_{RL} j_L^\mu J_{R\mu} + \epsilon_{LR} j_R^\mu J_{L\mu} + \epsilon_{RR} j_R^\mu J_{R\mu})$$





less severe due to lack of interference with SM

- ▶ Modification of angular and energy distribution in $2\nu\beta\beta$ decay
 - Current limit $\epsilon_{XR} < 3 \times 10^{-2}$ from NEMO3 competitive to other searches

New Physics in $2\nu\beta\beta$

FFD, Graf, Simkovic, PRL 125 (2020)

- Lepton-number conserving right-handed currents






New Physics in $2\nu\beta\beta$

FFD, Graf, Rodejohann, Xu, PRD 102 (2020)

► Neutrino self-interactions

- Same signature as SM $2\nu\beta\beta$ decay
- Potential interference with SM $2\nu\beta\beta$ decay
- Non-observation of enhanced rate excludes regime $G_S \approx 4 \times 10^9 G_F$ suggested to resolve Hubble tension

(Kreisch, Cyr-Racine, Doré,
PRD 101 (2020) 12, 123505)



New Physics in $2\nu\beta\beta$



FFD, Graf, Rodejohann, Xu, PRD 102 (2020)

► Neutrino self-interactions

- Same signature as SM $2\nu\beta\beta$ decay
- Potential interference with SM $2\nu\beta\beta$ decay
- Non-observation of enhanced rate excludes regime $G_S \approx 4 \times 10^9 G_F$ suggested to resolve Hubble tension

(Kreisch, Cyr-Racine, Doré,
PRD 101 (2020) 12, 123505)


- Excludes quadruple double beta decay



Heeck, Rodejohann,
EP Lett. 103 (2013) 32001

Conclusion

- ▶ **Neutrinos much lighter than other fermions**
 - Dirac or Majorana? Lepton Number Violation?
 - Determination of absolute mass scale
- ▶ **$0\nu\beta\beta$ is crucial probe for BSM physics**
 - Universal probe of LNV physics
 - LNV physics near GUT scale
 - Direct sensitivity to LNV physics at scales $m_N \approx 1 \text{ eV} - 100 \text{ TeV}$
 - Light exotic particles
- ▶ **$2\nu\beta\beta$ is sensitive to New Physics**
 - Ongoing and future searches probe $2\nu\beta\beta$ decay with high statistics
 - Exotic (right-handed) currents
 - Neutrino self-interactions
 - Endpoint searches for sterile neutrinos



$$\frac{T_{1/2}^{0\nu\beta\beta}}{10^{28} \text{ y}} \approx \left(\frac{\Lambda_{\text{NP}}}{10^{15} \text{ GeV}} \right)^2$$