

Quantum aspects of stimulated Hawking radiation in an analog white-black hole pair

Dimitrios Kranas

Louisiana State University, Baton Rouge, Louisiana, USA

LOOPS'22, ENS de Lyon, France

19 July 2022

This talk is based on:

Ivan Agullo, Anthony J. Brady, and Dimitrios Kranas. “Quantum Aspects of Stimulated Hawking Radiation in an Optical Analog White-Black Hole Pair”. In: Phys. Rev. Lett. 128 (9 Feb. 2022), p. 091301.

- **Hawking effect:** Creation of entangled pairs of particles by black hole event horizons. S. W. Hawking, Black hole explosions, Nature 248, 30 .
- **Direct observation?** For typical black hole masses: $T_H = T_{CMB} \ll 1$! signal buried under CMB.
- **Analog black holes:** A viable alternative for the experimental observation of the Hawking effect.

Optical black holes: The Kerr effect

- Ingredients:
 1. Dielectric medium.
 2. Strong EM pulse (pump).
 3. Weak EM pulse (probe).
- The mechanism: Kerr effect, i.e. non-linear interaction between the strong pulse and the medium.


$$n_e(\omega; x; t) = n(\omega) + n(x; t); \quad n(x; t) = E_S^2(x; t)$$

Modes and emergent causal structure

- In the comoving frame (;) w.r.t to the strong pulse, the system has time symmetry ! Conservation of frequency.
- For $\omega < \omega_h$, $\omega > \omega_h$: 4 modes (1 right-moving and 3 left-moving).
- For $\omega_h < \omega < \omega_h$: 2 modes (both left-moving).


Modes and emergent causal structure

- In the comoving frame (;) w.r.t to the strong pulse, the system has time symmetry ! Conservation of frequency.
- For $\omega < \omega_h$, $\omega > +\omega_h$: 4 modes (1 right-moving and 3 left-moving).
- For $\omega_h < \omega < +\omega_h$: 2 modes (both left-moving).



Modes and emergent causal structure

- In the comoving frame (;) w.r.t to the strong pulse, the system has time symmetry ! Conservation of frequency.
- For $\omega < \omega_h$, $\omega > +\omega_h$: 4 modes (1 right-moving and 3 left-moving).
- For $\omega_h < \omega < +\omega_h$: 2 modes (both left-moving).



- Hawking particle creation at the WH and BH horizons out of vacuum fluctuations!

Observations?

- T. G. Philbin et al, (2008), Science 319, 1367.
- J. Drori et al, (2019), Phys. Rev. Lett. 122, 010404.

Observations?

- T. G. Philbin et al, (2008), Science 319, 1367.
- J. Drori et al, (2019), Phys. Rev. Lett. 122, 010404.

Successes

- Generation of horizons.
- Particle production via the stimulated process.

Observations?

- T. G. Philbin et al, (2008), Science 319, 1367.
- J. Drori et al, (2019), Phys. Rev. Lett. 122, 010404.

Successes

- Generation of horizons.
- Particle production via the stimulated process.

Open questions



- Observation of a blackbody spectrum (dispersion the main obstacle and absence of perfect horizons).
- Generation of entanglement (fragile to background thermal noise).

The effect of background noise

- As input states, we restrict ourselves to the family of Gaussian quantum states and we quantify entanglement by means of Logarithmic Negativity (LN).
- We assume that the input modes are in thermal equilibrium with a blackbody bath of "environment" photons.

The effect of background noise

- As input states, we restrict ourselves to the family of Gaussian quantum states and we quantify entanglement by means of Logarithmic Negativity (LN).
- We assume that the input modes are in thermal equilibrium with a blackbody bath of "environment" photons.



- We find that the presence of the thermal background degrades the entanglement generated in the Hawking process.

The stimulated process

Stimulated Hawking emission: Input non-vacuum initial state (on top of the environmental noise).

The stimulated process

Stimulated Hawking emission: Input non-vacuum initial state (on top of the environmental noise).

- Coherent state:

$$\exp(-\hat{a}^\dagger \alpha) \hat{a} |0\rangle = e^{-\frac{1}{2}|\alpha|^2} \sum_{m=0}^{\infty} \frac{\alpha^m}{\sqrt{m!}} |m\rangle$$

The stimulated process

Stimulated Hawking emission: Input non-vacuum initial state (on top of the environmental noise).

- Coherent state:

$$\exp(\hat{a}^\dagger \alpha) |0\rangle = e^{-\frac{1}{2}|\alpha|^2} \sum_{m=0}^{\infty} \frac{\alpha^m}{\sqrt{m!}} |m\rangle$$

- Increase intensity of the Hawking radiation? !
- Enhances the generated entanglement? %

The stimulated process

Stimulated Hawking emission: Input non-vacuum initial state (on top of the environmental noise).

- Coherent state:

$$\exp(\alpha \hat{a}^\dagger) |0\rangle = e^{-\frac{1}{2}|\alpha|^2} \sum_{m=0}^{\infty} \frac{\alpha^m}{\sqrt{m!}} |m\rangle$$

- Increase intensity of the Hawking radiation? !
- Enhances the generated entanglement? %

- Single-mode squeezed state:

$$\exp\left[\frac{1}{2} r_1 e^{-i\theta} \hat{a}^2 - r_1 e^{i\theta} \hat{a}^{\dagger 2}\right] |0\rangle = \frac{1}{\cosh r_1} \sum_{m=0}^{\infty} \frac{(-1)^m (2m)!}{2^m m!} e^{im\theta} (\tanh r_1)^m |2m\rangle$$

The stimulated process

Stimulated Hawking emission: Input non-vacuum initial state (on top of the environmental noise).

- Coherent state:

$$\exp(\hat{a}^\dagger \alpha) |0\rangle = \sum_{m=0}^{\infty} \frac{\alpha^m}{m!} |m\rangle$$

- Increase intensity of the Hawking radiation? !
- Enhances the generated entanglement? %

- Single-mode squeezed state:

$$\exp\left(\frac{1}{2} r_1 e^{-i\theta} \hat{a}^2 - r_1 e^{i\theta} \hat{a}^{\dagger 2}\right) |0\rangle = \sum_{m=0}^{\infty} \frac{1}{\cosh r_1} \left(\frac{1}{2}\right)^m \frac{(2m)!}{2^m m!} e^{im\theta} (\tanh r_1)^m |2m\rangle$$


- Increase intensity of the Hawking radiation? !
- Enhances the generated entanglement? !

Entanglement, noise, and single-mode squeezed input

- We consider the ingoing k_3 mode to be in a single-mode squeezed state.

Entanglement, noise, and single-mode squeezed input

- We consider the ingoing k_3 mode to be in a single-mode squeezed state.



- Single-mode squeezed inputs **enhance** the **entanglement** generated by the horizons.

The protocol to extract T_H from observations

Intensities (classical signal)

- $\langle \hat{n}_3(t) \rangle = A(t) (e^{t/T_H} - 1)^{-1}$.

The protocol to extract T_H from observations

Intensities (classical signal)

- $\langle \hat{n}_3(t) \rangle = A(t)(e^{t/T_H} - 1)^{-1}$.

Entanglement (quantum signal)

- Theoretically, we compute the function $LN_{1j4}(t) = f(t; T_H; T_{env}; r_l)$.
- Measure all mode correlations $\langle \hat{a}_i^{out}(t); \hat{a}_j^{out}(t) \rangle$, $\langle \hat{a}_i^{out}(t); \hat{a}_j^{out y}(t) \rangle$, $\langle \hat{a}_i^{out y}(t); \hat{a}_j^{out y}(t) \rangle$.
- Construct $LN_{1j4}(t)$.
- Obtain T_H from $LN_{1j4}(t)$.

Take home messages

- The black hole causal structure can be reproduced in dispersive, inhomogeneous media.
- Stimulating the process with a single-mode squeeze state is a promising strategy for the detection of the Hawking effect as it enhances both the Hawking intensity and the entanglement generated by the Hawking process.
- Connecting observations (intensities and entanglement) with the causal horizon is essential for the experimental confirmation of the Hawking effect.

Additional Slides

- **Hawking effect:** Creation of entangled pairs of particles by black hole event horizons. S. W. Hawking, Black hole explosions, Nature 248, 30 .
- **Direct observation?** For typical black hole masses $T_H = T_{CMB} \ll 1$! signal buried under CMB.
- **Analog black holes:** Dispersive media where propagating perturbations experience causal horizons mimicking the structure of black hole/white hole spacetimes. Popular analog models include: 1) Hydrodynamic systems, Bose Einstein Condensates, Optical media, etc.

River-analog of black holes



River-analog of black holes



River metric in 1+1D:

$$ds^2 = u^2 dt^2 + (dx - V(x)dt)^2$$

Acoustic horizon condition $|V(x)| = u$.

River-analog of black holes



River metric in 1+1D:

$$ds^2 = u^2 dt^2 + (dx - V(x)dt)^2$$

Acoustic horizon condition $|V(x)| = u$.

$$ds^2 = - \left(1 - \frac{r_s}{r}\right) c^2 dt^2 + \frac{r^2}{\left(1 - \frac{r_s}{r}\right)} dr^2$$

$$ds^2 = c^2 d\tilde{t}^2 + \frac{r^2}{c^2} \left(\frac{dr}{r} + \frac{r_s}{r^2} d\tilde{t}\right)^2$$

Flow velocity: $V(r) = c \sqrt{\frac{r_s}{r}}$.

Optical black holes: The (microscopic) model

$$L = \underbrace{\frac{1}{2} \int d^3x \left(j_{@t} A^2 + j_{@x} A^2 \right)}_{\text{EM field}} + \underbrace{\frac{1}{2} \int d^3x j_{@t}^2 \Omega^2}_{\text{medium}} + \underbrace{g \text{Re} \left[\int d^3x \dot{A} \right]}_{\text{linear interaction}}$$

$$\Omega(x; t) = \Omega_0 + \underbrace{\frac{1}{2} \int d^3x S T j E_s(x; t)^2}_{\text{nonlinear interaction}}$$

$$n_e(\omega_{lab}; x; t) = \frac{S}{1 + \frac{g^2}{\Omega^2(x; t) \omega_{lab}^2}}$$

Wave equation

$$\frac{\partial^2 A(x;t)}{\partial t^2} - \frac{\partial^2 A(x;t)}{\partial x^2} = g \frac{\partial A(x;t)}{\partial t}$$


$$\frac{\partial^2 A(x;t)}{\partial t^2} + \Omega^2(x;t) A(x;t) = g \frac{\partial A(x;t)}{\partial t}$$



$$\nabla^2 u + \Omega^2(x;t) u = g \frac{\partial u}{\partial t}$$

Dispersion relation in the comoving frame


$$(\omega + uk) = \Omega(\omega) \frac{S}{1 + \frac{g^2}{\omega^2 k^2}}$$



Particle creation: a simple example

Consider a time dependent quantum harmonic oscillator.

$$\hat{H} = \frac{1}{2}\hat{p}^2 + \frac{1}{2}\omega^2(t)\hat{x}^2 \quad \frac{d^2\hat{x}}{dt^2} + \omega^2(t)\hat{x} = 0$$
$$[\hat{x}; \hat{p}] = i; \quad [\hat{a}; \hat{a}^\dagger] = 1$$




Particle creation: a simple example

Consider a time dependent quantum harmonic oscillator.

$$\hat{H} = \frac{1}{2}p^2 + \frac{1}{2}\omega^2(t)x^2 \quad \frac{d^2 \hat{x}}{dt^2} + \omega^2(t)\hat{x} = 0$$

$$[x; p] = i; \quad [a; \hat{a}^\dagger] = 1$$



For $t < 0$:

- $\omega(t) = \omega_1$; $\hat{x}(t) = \sqrt{\frac{1}{2\omega_1}} \hat{a}_1 e^{-i\omega_1 t} + \sqrt{\frac{1}{2\omega_1}} \hat{a}_1^\dagger e^{+i\omega_1 t}$
- $\hat{a}_1 |0\rangle_{i_1} = 0$; $\langle 0| \hat{a}_1^\dagger \hat{a}_1 |0\rangle_{i_1} = 0$

For $t > 0$:


- $\omega(t) = \omega_2$; $\hat{x}(t) = \sqrt{\frac{1}{2\omega_2}} \hat{a}_2 e^{-i\omega_2 t} + \sqrt{\frac{1}{2\omega_2}} \hat{a}_2^\dagger e^{+i\omega_2 t}$
- $\hat{a}_2 |0\rangle_{i_1} \neq 0$; $\langle 0| \hat{a}_2^\dagger \hat{a}_2 |0\rangle_{i_1} \neq 0$

Particle creation: a simple example

Consider a time dependent quantum harmonic oscillator.

$$\hat{H} = \frac{1}{2}p^2 + \frac{1}{2}\omega^2(t)x^2 \quad \frac{d^2x}{dt^2} + \omega^2(t)x = 0$$

$$[x; p] = i; \quad [a; a^\dagger] = 1$$



For $t < 0$:

- $\omega(t) = \omega_1$; $x(t) = \frac{1}{\sqrt{2\omega_1}} (a_1 e^{-i\omega_1 t} + a_1^\dagger e^{+i\omega_1 t})$
- $a_1 |0\rangle = 0$; $\langle 0| a_1^\dagger a_1 |0\rangle = 0$


For $t > 0$:

- $\omega(t) = \omega_2$; $x(t) = \frac{1}{\sqrt{2\omega_2}} (a_2 e^{-i\omega_2 t} + a_2^\dagger e^{+i\omega_2 t})$
- $a_2 |0\rangle \neq 0$; $\langle 0| a_2^\dagger a_2 |0\rangle \neq 0$

Time evolution:

$$e^{-i\omega_1 t} \quad e^{-i\omega_2 t} + e^{+i\omega_2 t}; \quad a_1 \quad a_2 \quad a_2^\dagger \quad \langle 0| a_2 a_2^\dagger |0\rangle = 1$$

WH-BH scattering via a quantum circuit



The scattering process at the black hole can be modeled via a **two-mode-squeezer** followed by a **beam splitter**.

Squeezer

$$\hat{a}_3^{ext} = \cosh r_H \hat{a}_4^{in} + e^i \sinh r_H (\hat{a}_1^{in})^y$$


$$\hat{a}_1^{int} = \cosh r_H \hat{a}_1^{in} + e^i \sinh r_H (\hat{a}_4^{in})^y$$

Beam splitter

$$\hat{a}_3^{out} = \cos \hat{a}_3^{ext} \sin \hat{a}_2^{in}$$

$$\hat{a}_2^{int} = \cos \hat{a}_2^{in} + \sin \hat{a}_3^{ext}$$

White hole-Black hole scattering




In the asymptotic regions ($r \rightarrow \infty$):

$$\hat{A}(r, t) = \sum_{j=1}^4 d_j e^{ik_j(t)} e^{-i\omega t} + \hat{a}_{j;l}^y e^{ik_j(t)} e^{+i\omega t}$$

$$\hat{a}_i^{out} = \sum_j \hat{a}_i^{in} + \sum_j \hat{a}_j^{in y}$$


Hawking radiation

Hawking radiation emitted via the particle creation process at the analog BH horizon.



Hawking radiation


Hawking radiation emitted via the particle creation process at the analog BH horizon.



$$\hat{a}_3^{out} = \alpha_{31} \hat{a}_1^{in} + \alpha_{32} \hat{a}_2^{in} + \alpha_{33} \hat{a}_3^{in} + \alpha_{34} \hat{a}_4^{in}$$

Hawking radiation

Hawking radiation emitted via the particle creation process at the analog BH horizon.




$$\hat{a}_3^{\text{out}} = \beta_{31} \hat{a}_1^{\text{in}} + \beta_{32} \hat{a}_2^{\text{in}} + \beta_{33} \hat{a}_3^{\text{in}} + \beta_{34} \hat{a}_4^{\text{in}}$$

$$\hat{N}_3^{\text{out}} = \sum_j \beta_{3j}^2 \hat{N}_j^{\text{in}}$$

Particle creation from quantum nothing!

Hawking radiation

Hawking radiation emitted via the particle creation process at the analog BH horizon.



$$\hat{a}_3^{\text{out}} = \alpha_{31} \hat{a}_1^{\text{in}} + \alpha_{32} \hat{a}_2^{\text{in}} + \alpha_{33} \hat{a}_3^{\text{in}} + \alpha_{34} \hat{a}_4^{\text{in}}$$


$$\hat{N}_3^{\text{out}} = \sum_{i \in \text{D}} \sum_{j \in \text{E}} \alpha_{3i}^{\text{out}} \alpha_{3j}^{\text{out}} \hat{N}_{i \text{ in}} = \sum_{i \in \text{D}} \alpha_{3i}^2$$

Particle creation from quantum nothing!

$$\sum_{i \in \text{D}} \alpha_{3i}^2 = \frac{1}{e^{f(!)/T_H}}; \quad T_H = \frac{u}{2}$$


Modes and emergent causal structure

- In the comoving frame (;) w.r.t to the strong pulse, the system has time symmetry ! Conservation of frequency.
- For $j > j_h$: 4 modes (1 right-moving and 3 left-moving).
- For $j < j_h$: 2 modes (both left-moving).



Modes and emergent causal structure

- In the comoving frame (;) w.r.t to the strong pulse, the system has time symmetry ! Conservation of frequency.
- For $j > j_h$: 4 modes (1 right-moving and 3 left-moving).
- For $j < j_h$: 2 modes (both left-moving).





- Hawking particle creation at the WH and BH horizons out of vacuum fluctuations!

Gaussian states and Logarithmic Negativity

- We work with the family Gaussian states, where all the information about the quantum state is encoded in the first moments $\tilde{\mu} = \begin{matrix} D \\ E \end{matrix} A$ and the covariance matrix $\Sigma = \begin{matrix} fA & \\ & g \end{matrix}$, where $A = (\hat{a}_1; \hat{a}_1^y; \hat{a}_2; \hat{a}_2^y; \hat{a}_3; \hat{a}_3^y; \hat{a}_4; \hat{a}_4^y)^T$.
- The evolution of the state is given by: $A_{out} = S A_{in}$, $\tilde{\mu}_{out} = S \tilde{\mu}_{in}$, $\Sigma_{out} = S \Sigma_{in} S^T$.
- We use Logarithmic negativity (LN), a well-known entanglement monotone in quantum information theory, to study and quantify the entanglement produced in the Hawking process. LN can be easily computed from the first and second moments.

Entanglement structure of the analog WH-BH




- The BH and WH horizons behave as frequency-dependent two-mode squeezers generating entanglement in the modes $(k_1^{\text{out}}|k_3^{\text{out}})$ and $(k_1^{\text{out}}|k_4^{\text{out}})$, respectively.
- At low frequencies: $\text{LN}_{1/4} > \text{LN}_{1/3}$.
- At larger frequencies: $\text{LN}_{1/4} < \text{LN}_{1/3}$.

Generation of entanglement from vacuum fluctuations

- To study the evolution, we use Gaussian states and we quantify entanglement by means of Logarithmic Negativity (LN).
- The BH and WH horizons behave as frequency-dependent two-mode squeezers generating entanglement in the bipartitions $(k_1^{out} j k_2^{out})$, $(k_1^{out} j k_3^{out})$ and $(k_1^{out} j k_4^{out})$, respectively.
- The strongest correlated couple is the WH Hawking pair of modes $(k_1^{out} j k_4^{out})$.

Generation of entanglement from vacuum fluctuations

- To study the evolution, we use Gaussian states and we quantify entanglement by means of Logarithmic Negativity (LN).
- The BH and WH horizons behave as frequency-dependent two-mode squeezers generating entanglement in the bipartitions $(k_1^{out} j k_2^{out})$, $(k_1^{out} j k_3^{out})$ and $(k_1^{out} j k_4^{out})$, respectively.
- The strongest correlated couple is the WH Hawking pair of modes $(k_1^{out} j k_4^{out})$.



Observations?

- T. G. Philbin et al, (2008), Science 319, 1367.
- S. Weinfurtner et al, (2011), Phys. Rev. Lett. 106, 021302
- J. Steinhauer, (2016), Nature Phys. 12, 959-965.
- J. Drori et al, (2019), Phys. Rev. Lett. 122, 010404.

Successes

- Generation of horizons.
- Particle production via the stimulated process.

Open questions

- Observation of a blackbody spectrum (dispersion the main obstacle and absence of perfect horizons).
- Generation of entanglement (fragile to background thermal noise).

The protocol to extract T_H from observations

Intensities (classical signal)

- $\langle \hat{n}_3(t) \rangle = A(t) (e^{t/T_H} - 1)^{-1}$

Entanglement (quantum signal)

$$LN_{1j4}(t) = \log_2 \frac{1}{4} [7 - 4 \cosh 2r(t) + 8 \cosh 4r(t) + 4 \cosh 6r(t) + \cosh 8r(t)] \\ 16 \cosh^2 r(t) \cosh 2r(t)^{3=2} (9 + 6 \cosh 2r(t) + \cosh 4r(t))^{1=2} \sinh r(t)^{1=2}$$

- Measure all mode correlations $\langle \hat{a}_i(t) \hat{a}_j(t) \rangle$
- Construct $LN_{1j4}(t)$
- Obtain T_H from $LN_{1j4}(t)$

Take home messages

- The black hole causal structure can be reproduced in dispersive, inhomogeneous media.
- Frequency conservation and dispersion single out a finite number of degrees of freedom interacting with each other, allowing us to import theoretically rigorous and experimentally accessible tools from quantum information theory to study the entanglement in the Hawking process.
- Stimulating the process with a single-mode squeeze state is a promising strategy for the detection of the Hawking effect as it enhances both the Hawking intensity and the entanglement generated by the Hawking process.