


Amplitudes, Angular Momentum, and the Binary Inspiral Problem

Chia-Hsien Shen 沈家賢 (UC San Diego)

2203.04283 with Manohar and Ridgway


“Can we solve the full binary inspiral problem?”



[PC: BBC Science focus]

Symmetry

Lorentz
invariance




$$\gamma = \sigma \equiv \frac{p_1 \cdot p_2}{m_1 m_2}$$

$$A(t, s)$$

$$\theta(J, E)$$

Conservative
Dynamics



$$V(r, \vec{p}^2, \vec{p} \times \hat{r})$$

Effective Field Theory

[Goldberger, Rothstein '04]

Scattering Amplitudes


Generalized unitarity [Bern, Dixon, Dunbar, Kosower]...

GR=YM² [Kawai, Lewellen, Tye][Bern, Carrasco, Johansson]...

Symmetry



Dissipative
 Dynamics



Starts at 2.5PN!!

State of the art: (partial) 4.5PN

2.5PN		3.5PN		4.5PN					
v^3	+	v^5	+	v^7	+	v^9	+	\dots	G^2
v	+	v^3	+	v^5	+	v^7	+	\dots	G^3
		v	+	v^3	+	v^5	+	\dots	G^4

$$F_{\text{RR}}(r, \vec{p}^2, \vec{p} \cdot \hat{r})$$


[Burke, Throne '69]

Pioneering idea: [Iyer, Will]

similar idea in B2B [Cho, Kalin, Porto]


Symmetry

Poincare
invariance



$$E, J$$

Dissipative
Dynamics




$$F_{RR}(r, \vec{p}^2)$$

[Manohar, Ridgway, CHS, 2203.04283]

Symmetry


Poincare
invariance



$$E, J$$

[Herrmann, Parra-Martinez, Ruf, Zeng] [This talk]

Dissipative
Dynamics



$$F_{RR}(r, \vec{p}^2)$$

2.5PN	3.5PN	4.5PN		
$(v^3 + v^5 + v^7 + v^9 + \dots)$				G^2
$(v + v^3 + v^5 + v^7 + \dots)$				G^3
	$(v + v^3 + v^5 + \dots)$			G^4

Radiated Angular Momentum

*We do not intend to resolve the BMS subtlety here

Radiated Angular Momentum

- Textbook formula for angular momentum

$$J^i = \frac{c^2}{32\pi G} \int d^3x \left[-\epsilon^{ikl} \dot{h}_{ab}^{\text{TT}} x^k \partial^l h_{ab}^{\text{TT}} + 2\epsilon^{ikl} h_{ak}^{\text{TT}} \dot{h}_{al}^{\text{TT}} \right]. \quad (2.51)$$

Why not covariant?


How to see gauge invariance?

Radiated Angular Momentum

- Consider the final state of scattering.

The radiated linear and angular momentum are

$$P^\mu = \int d^3x T^{\mu 0}$$
$$J^{\mu\nu} = \int d^3x \underline{x}^{[\mu} T^{\nu]0}$$




- Need to include the $1/r^2$ of the field for angular momentum

Sources \longrightarrow Fields $\longrightarrow T^{\mu\nu}$ \longrightarrow linear/angular momentum

Sources

- EM: currents \mathcal{J}^μ
- **3 degrees** of freedom under conservation $k_\nu \mathcal{J}^\nu(k) = 0 \quad \partial_\mu \mathcal{J}^\mu(x) = 0$
- “on-shell” part can be projected to transverse mode
 $\mathcal{J}^\nu(k) \rightarrow \mathcal{J}^\nu(k) + \alpha k^\nu$
- Gravity: stress-energy pseudotensor $\mathcal{T}^{\mu\nu}$
- **6 degrees** of freedom under conservation $k_\mu \mathcal{T}^{\mu\nu}(k) = 0 \quad \partial_\mu \mathcal{T}^{\mu\nu}(x) = 0$
- “on-shell” part can be projected to traceless and transverse mode
 $\mathcal{T}^{\mu\nu}(k) \rightarrow \mathcal{T}^{\mu\nu}(k) + k^\mu \epsilon^\nu(k) + k^\nu \epsilon^\mu(k)$



Fields

- Solved by fixing a gauge, e.g., under Lorentz gauge $\square A_\mu = 4\pi J_\mu$
- No ambiguity after gauge fixing, even for static sources

- Position space:

Needed for angular momentum

$$A^\mu(x) = \int d\omega e^{-i\omega u} \left(\frac{1}{r} \mathcal{J}^\mu(k = \omega(1, \hat{r})) + \frac{1}{r^2} \mathcal{O}(\partial_k J) \right)$$

residual gauge redundancy
→ **2 degrees of freedom**
→ amplitudes!

No gauge redundancy
→ **≥ 2 degrees of freedom**

$$\vec{E} = \frac{Q\hat{r}}{r^2}$$

Fields

- Solved by fixing a gauge, e.g., under Lorentz gauge $\square A_\mu = 4\pi J_\mu$
- No ambiguity after gauge fixing, even for static sources
- Momentum space: $k^2 = 0$

$$A_\mu(x) = \int \underline{d\tilde{k}} \left(\underline{\mathcal{J}_\mu(k)} e^{-ik \cdot x} + \text{c.c.} \right)$$

Lorentz-invariant phase space sources (current or stress-energy pseudotensor)

Stress-energy Tensor

- EM: gauge-invariant

$$T^{\mu\nu} = -F^{\mu}_{\rho} F^{\nu\rho} + \eta^{\mu\nu} \frac{1}{4} F^{\rho\sigma} F_{\rho\sigma}$$

- Gravity: *not* gauge invariant

$$T_{\mu\nu} = -\frac{1}{8\pi G} \left(R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R \right)$$

- But global charges are gauge invariant


Radiated Poincare Charges

- Radiated linear and angular momentum are

$$P^\mu = \int d^3x T^{\mu 0}$$

$$J^{\mu\nu} = \int d^3x x^{[\mu} T^{\nu]0}$$

- Need to include the $1/r^2$ of the field, so we use sources instead of amplitudes



Radiated Linear Momentum

- Linear momentum:

same as cross section weighted by momentum

- EM: $P^\mu = \int \widetilde{d}k k^\mu (-\mathcal{J}^{*\rho}(k)\mathcal{J}_\rho(k))$

Phase space integral momentum cross section

- GR: $P^\mu = 8\pi G \int \widetilde{d}k k^\mu \left(\mathcal{T}^{*\rho\sigma}(k)\mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k)\mathcal{T}_\sigma^\sigma(k)}{D-2} \right)$

- Covariance and gauge invariance are obvious

- Directly related to on-shell amplitudes via KMOC

Radiated Angular Momentum

- New covariant formula in GR for angular momentum

$$P^\mu = 8\pi G \int \widetilde{d\mathbf{k}} k^\mu \left(\mathcal{T}^{*\rho\sigma}(k) \mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k) \mathcal{T}_\sigma^\sigma(k)}{D-2} \right), \quad \mathcal{L}^{\mu\nu} = ik^\mu \frac{\partial}{\partial k_\nu} - ik^\nu \frac{\partial}{\partial k_\mu}$$
$$J^{\mu\nu} = 8\pi G \int \widetilde{d\mathbf{k}} \left(\mathcal{T}^{*\rho\sigma}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_\sigma^\sigma(k)}{D-2} + 2i \mathcal{T}^{*\rho[\mu}(k) \mathcal{T}^{\nu]\rho}(k) \right)$$

[Manohar, Ridgway, CHS]

Radiated Angular Momentum

- $$P^\mu = 8\pi G \int \widetilde{d}k k^\mu \left(\mathcal{T}^{*\rho\sigma}(k) \mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k) \mathcal{T}_\sigma^\sigma(k)}{D-2} \right), \quad \mathcal{L}^{\mu\nu} = ik^\mu \frac{\partial}{\partial k_\nu} - ik^\nu \frac{\partial}{\partial k_\mu}$$

$$J^{\mu\nu} = 8\pi G \int \widetilde{d}k \left(\mathcal{T}^{*\rho\sigma}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_\sigma^\sigma(k)}{D-2} + 2i \mathcal{T}^{*\rho[\mu}(k) \mathcal{T}^{\nu]\rho}(k) \right)$$

- Poincare algebra:**

$$x^\mu \rightarrow x^\mu + a^\mu \qquad P^\mu \rightarrow P^\mu$$

$$\mathcal{T}^{\mu\nu}(k) \rightarrow \mathcal{T}^{\mu\nu}(k) e^{ik \cdot a} \longrightarrow J^{\mu\nu} \rightarrow J^{\mu\nu} + a^{[\mu} P^{\nu]}$$

- Gauge invariance:** $\mathcal{T}^{\mu\nu}(k) \rightarrow \mathcal{T}^{\mu\nu}(k) + k^\mu \epsilon^\nu(k) + k^\nu \epsilon^\mu(k)$

- No physical separation b/w “orbital” and “spin” parts

Zero-Frequency Limit & Leading order

(Weinberg soft theorem, memory effect, ...)


Stress-energy Pseudotensor

- Zoom out the time scale so collision occurs at $t=0$ (zero-frequency limit)



$$\mathcal{T}^{\mu\nu}(k)|_{\omega \rightarrow 0^+} = \underbrace{-i\pi\delta(\omega) \sum_a \frac{p_a^\mu p_a^\nu}{E_a - \hat{\mathbf{k}} \cdot \mathbf{p}_a}}_{\text{Free particles (Coulomb mode)}} + \underbrace{\frac{1}{\omega + i0} \sum_a \left(\frac{p_a^\mu p_a^\nu}{E_a - \hat{\mathbf{k}} \cdot \mathbf{p}_a} \right)}_{\text{deflection turned on at } t=0 \text{ valid to all orders}} \Big|_i^f$$

Free particles
(Coulomb mode)

deflection turned on at $t=0$
valid to all orders



$$\delta(u = t - r)$$



Radiated angular momentum

- Zero-frequency limit

$$P^\mu = 0,$$

$$J^{\mu\nu} = 8\pi G \int \frac{d\tilde{k}}{d\omega \omega \delta(\omega)} \left(\mathcal{T}^{*\rho\sigma}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_\sigma^\sigma(k)}{D-2} + 2i \mathcal{T}^{*\rho[\mu}(k) \mathcal{T}^{\nu]\rho}(k) \right) \frac{1}{\omega}$$

- Nothing wrong w/ zero energy but non-zero angular momentum

$$\mathbf{J} = \mathbf{r} \times \mathbf{p} \sim \int d\Omega r^2 \mathbf{r} \times (\mathbf{E} \times \mathbf{B}) \frac{1}{r^2} \frac{1}{r}$$

Radiated angular momentum

- Leading order in deflection angle θ

$$\frac{J_{\text{CM},2}^{12}}{J_{\text{CM}}} = 2 \times \frac{J_{\text{rest},2}^{12}}{J_{\text{rest}}} = 2m_1m_2 \mathcal{I}(\sigma) \theta$$

$$\mathcal{I}(\sigma) = -\frac{16}{3} + \frac{2\sigma^2}{\sigma^2 - 1} + \frac{4(2\sigma^2 - 3)}{\sigma^2 - 1} \frac{\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2 - 1}}$$


- Model independent
(GR, with spin, dilaton gravity, supergravity, etc)
- Radiated Angular momentum *is positive* when scattering is *attractive*

Soft Finiteness to all orders

- Total radiated angular momentum is finite

$$J^{\mu\nu} = 8\pi G \int \widetilde{dk} \left(\mathcal{T}^{*\rho\sigma}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_\sigma^\sigma(k)}{D-2} + 2i \mathcal{T}^{*\rho[\mu}(k) \mathcal{T}^{\nu]\rho}(k) \right)$$

$$\int d\Omega \int d\omega \omega \frac{1}{\omega} \frac{1}{\omega}$$



- soft divergence cancels after integrating the full sphere

Comparison

- J^{12} at G^2 agrees with [Damour]
- Fully agrees arbitrary deflection [Di Vecchia, Heissenberg, Russo]
- J^{0i} at G^2 agrees with [Gralla, Lobo] (modulo a potential extra term)
- **Disagree with the textbook formula in the initial rest frame by x^2**
[Jakobsen, Mogull, Plefka, Steinhoff][Mougiakakos, Riva, Vernizzi]

Leading $O(G^2)$ results

Textbook formula

TT part of metric

$$\frac{J_{\text{rest},2}^{12}}{J_{\text{rest}}} = \frac{J_{\text{CM},2}^{12}}{J_{\text{CM}}}$$

Our formula

stress-energy pseudotensor


$$\frac{J_{\text{rest},2}^{12}}{J_{\text{rest}}} = \frac{1}{2} \times \frac{J_{\text{CM},2}^{12}}{J_{\text{CM}}}$$

- Both agree in the CM frame
- Independent checks: general covariance and 3.5PN RR force

[Jaranowski, Schafer; Nissanke, Blanchet]

Non-perturbative Parametrization

- Form factors: Lorentz covariance + Poincare algebra



$$P^\mu = \underline{F_1 p_1^\mu + F_2 p_2^\mu + F_3 \Delta b^\mu},$$

$$J^{\mu\nu} = \underline{\bar{b}^{[\mu} (F_1 p_1^{\nu]} + F_2 p_2^{\nu]})} + \underline{\Delta b^{[\mu} (G_1 p_1^{\nu]} - G_2 p_2^{\nu]})} + \underline{H_{12} p_2^{[\mu} p_1^{\nu]}}$$

CM Frame

$$\left. \frac{J_{\text{CM}}^{12}}{J_{\text{CM}}} \right|_{\omega=0} = G_1 + G_2$$


Rest Frame

$$\left. \frac{J_{\text{rest}}^{12}}{J_{\text{rest}}} \right|_{\omega=0} = G_2$$

- Since $G_1 = G_2$ at leading order, our answer agrees with this general prediction

Crosscheck with Burke-Throne

- Burke-Throne force at G^2 : $\mathbf{a}_1 = -\mathbf{a}_2 = \frac{4G^2 m_1 m_2}{5r^3} (3v^2 v_r \hat{\mathbf{r}} - v^2 \mathbf{v})$




- Final energy is the same as initial
- Impact parameter shrinks equally
- Non-decoupling of heavy particle

Crosscheck with Burke-Throne


- Burke-Throne force at G^2 : $\mathbf{a}_1 = -\mathbf{a}_2 = \frac{4G^2 m_1 m_2}{5r^3} (3v^2 v_r \hat{\mathbf{r}} - v^2 \mathbf{v})$

CM Frame:



$$\frac{J_{\text{CM},2}^{12}}{J_{\text{CM}}} = \frac{2p\Delta b}{pb} = 2 \times \frac{\Delta b}{b}$$

Rest Frame:



$$\frac{J_{\text{rest},2}^{12}}{J_{\text{rest}}} = \frac{p'\Delta b}{p'b} = \frac{\Delta b}{b}$$

Non-decoupling of heavy particle. Back reaction is important!

$$J^{\mu\nu} = 8\pi G \int \widetilde{d^4k} \left(\mathcal{T}^{*\rho\sigma}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_{\rho\sigma}(k) - \frac{\mathcal{T}_\rho^{*\rho}(k) \mathcal{L}^{\mu\nu} \mathcal{T}_\sigma^\sigma(k)}{D-2} + 2i \mathcal{T}^{*\rho[\mu}(k) \mathcal{T}^{\nu]\rho}(k) \right)$$

Our formula agrees with covariance and Burke-Throne force

Standard formula is incomplete for scattering because
of the presence of soft mode

Precision Frontier:

Radiated angular momentum at G^3


Radiated Poincare Charges

- State of the art precision at G^3

$P^\mu \rightarrow$ Known [Herrmann, Parra-Martinez, Ruf, Zeng]

$J^{\mu\nu} \rightarrow$ Both **zero** and **finite frequency** contributions


Soft Theorem



- Same as before, just use G^2 impulses

[Westpfahl 80's]

Double Copy & Generalized Unitarity



- Waveform from 2-to-3 amplitude via KMOC

- Resum velocity expansion from $O(v^{60})$ series

[See Carlos Heissenberg's poster]

New Results in General Relativity

- New results for G^3 radiated angular momentum

$$\frac{J_3}{\pi} = \frac{28}{5} p_\infty^2 + \left(\frac{739}{84} - \frac{163}{15} \nu \right) p_\infty^4 + \left(-\frac{5777}{2520} - \frac{5339}{420} \nu + \frac{50}{3} \nu^2 \right) p_\infty^6 + \left(\frac{115769}{126720} + \frac{1469}{504} \nu + \frac{9235}{672} \nu^2 - \frac{553}{24} \nu^3 \right) p_\infty^8 + \dots$$

[Bini, Damour, Geralico '21]

2.5PN	3.5PN	4.5PN			
$(v^3 + v^5 + v^7 + v^9 + \dots)$					G^2
$(v + v^3 + v^5 + v^7 + \dots)$					G^3
	$(v + v^3 + v^5 + \dots)$				G^4

[Manohar, Ridgway, CHS]

$$\begin{aligned} \mathcal{I}(\sigma) &= -\frac{16}{3} + \frac{2\sigma^2}{\sigma^2 - 1} + \frac{(2\sigma^2 - 3)\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2 - 1}} \\ \frac{\mathcal{E}(\sigma)}{\pi} &= f_1 + f_2 \log\left(\frac{\sigma+1}{2}\right) + f_3 \frac{\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2 - 1}} \\ \frac{\mathcal{C}(\sigma)}{\pi} &= g_1 + g_2 \log\left(\frac{\sigma+1}{2}\right) + g_3 \frac{\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2 - 1}} \\ \mathcal{D}(\sigma) &= \frac{3\pi(5\sigma^2 - 1)}{8} \mathcal{I}(\sigma) \\ f_1 &= \frac{210\sigma^6 - 552\sigma^5 + 339\sigma^4 - 912\sigma^3 + 3148\sigma^2 - 3336\sigma + 1151}{48(\sigma^2 - 1)^{3/2}} \\ f_2 &= -\frac{35\sigma^4 + 60\sigma^3 - 150\sigma^2 + 76\sigma - 5}{8\sqrt{\sigma^2 - 1}} \\ f_3 &= \frac{(2\sigma^2 - 3)(35\sigma^4 - 30\sigma^2 + 11)}{8(\sigma^2 - 1)^{3/2}} \\ g_1 &= \frac{105\sigma^7 - 411\sigma^6 + 240\sigma^5 + 537\sigma^4 - 683\sigma^3 + 111\sigma^2 + 386\sigma - 237}{24(\sigma^2 - 1)^2} \\ g_2 &= \frac{35\sigma^5 - 90\sigma^4 - 70\sigma^3 + 16\sigma^2 + 155\sigma - 62}{4(\sigma^2 - 1)} \\ g_3 &= \frac{-(2\sigma^2 - 3)(35\sigma^5 - 60\sigma^4 - 70\sigma^3 + 72\sigma^2 + 19\sigma - 12)}{4(\sigma^2 - 1)^2} \end{aligned}$$

Applications: impulses at G^4

- Predict for G^4 odd-in- v impulses [Bini, Damour, Geralico]

$$\Delta p_{\perp,4} = \nu M^5 \left(\frac{G}{b} \right)^4 (c_{b,4}^{\text{cons}} + c_{b,4}^{\text{rr,even}} + c_{b,4}^{\text{rr,odd}})$$

Conservative; even in v

Dissipative; even in v

Dissipative; odd in v

Unknown!!

[Bern, Parra-Martinez, Roiban, Ruf, CHS, Solon, Zeng, '21]

[Manohar, Ridgway, CHS '22]


[Dlapa, Kalin, Liu, Porto '21]

- Simplify a G^4 problem into mostly leading order inputs via **factorizations in EFT** [Bini, Damour]

$$2\chi^{\text{rr}} = \frac{\partial \chi^{\text{cons}}}{\partial J} \Delta J + \frac{\partial \chi^{\text{cons}}}{\partial E} \Delta E$$

Precision Binary Dynamics

$m_1=m_2, G=0.01, E[0]=-0.0176, J[0]=0.4, v[0]=0.128$
 $t = 0.0$
 $\{E, J\} = \{-0.0176, 0.400\}$



$$\begin{aligned}
 \dot{\mathbf{x}} &= \frac{\partial H}{\partial \mathbf{p}} \\
 \dot{\mathbf{p}} &= -\frac{\partial H}{\partial \mathbf{x}} + \mathbf{F}_{RR}
 \end{aligned}$$

Conservative

All orders in v to G^4
 from the scattering angle

[Bern, Cheung, Roiban, CHS, Solon, Zeng, '19]

[Bern, Parra-Martinez, Roiban, Ruf, CHS, Solon, Zeng, '21]

[Dlapa, Kalin, Liu, Porto '21]

Dissipative

All orders in v to G^3
 from radiated E and J

[Manohar, Ridgway, CHS '22]


*Caveat: illustration only

Don't trust a plot made by theorists

Precision Binary Dynamics

Only conservative effect included in PM so far

Can dissipation bring closer to numerical data?



Summary & Outlook

- **Symmetries** are powerful
- **State-of-the-art dissipative force in GR purely from YM**
- New formulae, perturbative and non-perturbative results of **angular momentum**
- **Higher order** dissipative effects?
 - Interesting functions & Tail effect. Mapping to bound states?
- Applications of angular momentum in **particle physics**?

Thank you

Backups

New Results in General Relativity

- New results for G^3 radiated angular momentum

$$J_{\text{rest},3}^{12} = bm_1m_2^2 \left(\underline{m_1\mathcal{C}(\sigma)} + \underline{(m_1 + m_2)\mathcal{D}(\sigma)} \right)$$

- As the form factors show, **radiated energy** enters when translating from rest to CM frame

$$\frac{J_{\text{CM},3}^{12}}{J_{\text{CM}}} = \frac{m_1m_2(m_1 + m_2)}{\sqrt{\sigma^2 - 1}} \left[\underline{\mathcal{C}(\sigma)} + \underline{2\mathcal{D}(\sigma)} - \frac{m_1m_2\sqrt{\sigma^2 - 1}}{E^2} \underline{\mathcal{E}(\sigma)} \right]$$

- Elucidate the relation originally found by Bini, Damour, Geralico when considering G^4 scattering

$$\begin{aligned} \mathcal{I}(\sigma) &= -\frac{16}{3} + \frac{2\sigma^2}{\sigma^2 - 1} + \frac{(2\sigma^2 - 3)\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2 - 1}} \\ \frac{\mathcal{E}(\sigma)}{\pi} &= f_1 + f_2 \log\left(\frac{\sigma+1}{2}\right) + f_3 \frac{\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2 - 1}} \\ \frac{\mathcal{C}(\sigma)}{\pi} &= g_1 + g_2 \log\left(\frac{\sigma+1}{2}\right) + g_3 \frac{\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2 - 1}} \\ \mathcal{D}(\sigma) &= \frac{3\pi(5\sigma^2 - 1)\mathcal{I}(\sigma)}{8} \\ f_1 &= \frac{210\sigma^6 - 552\sigma^5 + 339\sigma^4 - 912\sigma^3 + 3148\sigma^2 - 3336\sigma + 1151}{48(\sigma^2 - 1)^{3/2}} \\ f_2 &= -\frac{35\sigma^4 + 60\sigma^3 - 150\sigma^2 + 76\sigma - 5}{8\sqrt{\sigma^2 - 1}} \\ f_3 &= \frac{(2\sigma^2 - 3)(35\sigma^4 - 30\sigma^2 + 11)}{8(\sigma^2 - 1)^{3/2}} \\ g_1 &= \frac{105\sigma^7 - 411\sigma^6 + 240\sigma^5 + 537\sigma^4 - 683\sigma^3 + 111\sigma^2 + 386\sigma - 237}{24(\sigma^2 - 1)^2} \\ g_2 &= \frac{35\sigma^5 - 90\sigma^4 - 70\sigma^3 + 16\sigma^2 + 155\sigma - 62}{4(\sigma^2 - 1)} \\ g_3 &= \frac{-(2\sigma^2 - 3)(35\sigma^5 - 60\sigma^4 - 70\sigma^3 + 72\sigma^2 + 19\sigma - 12)}{4(\sigma^2 - 1)^2} \end{aligned}$$

Comparison


	Scalar	EM	GR
Spacetime	Minkowski	Minkowski	Dynamical
Gauge freedom	No	Yes	Yes
zero E but nonzero J	yes	yes	yes
RR force	[SEP] agree	Abraham-Lorentz-Dirac [SEP] agree	Burke-Thorne [SEP] agree
BMS ambiguity	No	No	Maybe?

CM Frame v.s. Rest frame

- They are related by boost and translation

CM Frame:

$$p_2^\mu = (E_2, -|\mathbf{p}|, 0, 0)$$




$$p_1^\mu = (E_1, |\mathbf{p}|, 0, 0)$$

$$J_{\text{CM}} = |\mathbf{p}|b$$

Rest Frame:

$$p_2^\mu = (\sigma m_2, -\sqrt{\sigma^2 - 1}m_2, 0, 0)$$



$$p_1^\mu = (m_1, 0, 0, 0)$$

$$J_{\text{rest}} = \sqrt{\sigma^2 - 1}m_2b$$

Precision Binary Dynamics


- State-of-the-art EOM all orders in v to G^3

$$H(r, p^2)$$

$$c_1 = \frac{\nu^2 m^2}{\gamma^2 \xi} (1 - 2\sigma^2),$$

$$c_2 = \frac{\nu^2 m^3}{\gamma^2 \xi} \left[\frac{3}{4} (1 - 5\sigma^2) - \frac{4\nu\sigma(1 - 2\sigma^2)}{\gamma\xi} - \frac{\nu^2(1 - \xi)(1 - 2\sigma^2)^2}{2\gamma^3\xi^2} \right],$$

$$c_3 = \frac{\nu^2 m^4}{\gamma^2 \xi} \left[\frac{1}{12} (3 - 6\nu + 206\nu\sigma - 54\sigma^2 + 108\nu\sigma^2 + 4\nu\sigma^3) - \frac{4\nu(3 + 12\sigma^2 - 4\sigma^4) \operatorname{arcsinh}\sqrt{\frac{\sigma-1}{2}}}{\sqrt{\sigma^2-1}} \right. \\ \left. - \frac{3\nu\gamma(1 - 2\sigma^2)(1 - 5\sigma^2)}{2(1 + \gamma)(1 + \sigma)} - \frac{3\nu\sigma(7 - 20\sigma^2)}{2\gamma\xi} + \frac{2\nu^3(3 - 4\xi)\sigma(1 - 2\sigma^2)^2}{\gamma^4\xi^3} \right. \\ \left. - \frac{\nu^2(3 + 8\gamma - 3\xi - 15\sigma^2 - 80\gamma\sigma^2 + 15\xi\sigma^2)(1 - 2\sigma^2)}{4\gamma^3\xi^2} + \frac{\nu^4(1 - 2\xi)(1 - 2\sigma^2)^3}{2\gamma^6\xi^4} \right].$$



$$\mathbf{F}_{RR} = c_r p_r \hat{\mathbf{r}} + c_p \mathbf{p}$$

$$c_r = \frac{G^2}{r^3} c_{r,2}(\mathbf{p}^2) + \frac{G^3}{r^4} c_{r,3}(\mathbf{p}^2) + \dots,$$

$$c_p = \frac{G^2}{r^3} c_{p,2}(\mathbf{p}^2) + \frac{G^3}{r^4} c_{p,3}(\mathbf{p}^2) + \dots,$$

$$c_{r,2}(\mathbf{p}^2) = -3c_{p,2}(\mathbf{p}^2),$$

$$c_{p,2}(\mathbf{p}^2) = -\frac{\nu^2 M^4}{E_1 E_2} (2\sigma^2 - 1) \mathcal{I}(\sigma)$$

$$c_{p,3}(\mathbf{p}^2) = -\frac{2p_\infty J_{CM,3}^{12}}{\pi\xi E J_0} + \left(2\xi E c'_{p,2}(\mathbf{p}^2) - \left(2 - \frac{p_\infty^2(1 - 3\xi)}{\xi^2 E^2} \right) \frac{J_{CM,2}^{12}}{2p_\infty J_0} \right) c_{H,1}(\mathbf{p}^2) - p_\infty c'_{H,1}(\mathbf{p}^2) \frac{J_{CM,2}^{12}}{J_0}$$

$$c_{r,3}(\mathbf{p}^2) = \frac{8}{\pi p_\infty} \left(\frac{p_\infty^2}{J_0 E \xi} J_{CM,3}^{12} - E_{CM,3} \right) + \left(-6\xi E c'_{p,2}(\mathbf{p}^2) + 2 \left(1 + \frac{p_\infty^2(1 - 3\xi)}{\xi^2 E^2} \right) \frac{J_{CM,2}^{12}}{p_\infty J_0} \right) c_{H,1}(\mathbf{p}^2) + 4p_\infty c'_{H,1}$$

$$\mathcal{I}(\sigma) = -\frac{16}{3} + \frac{2\sigma^2}{\sigma^2-1} + \frac{(2\sigma^2-3)\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2-1}}$$

$$\frac{\mathcal{E}(\sigma)}{\pi} = f_1 + f_2 \log\left(\frac{\sigma+1}{2}\right) + f_3 \frac{\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2-1}}$$

$$\frac{\mathcal{C}(\sigma)}{\pi} = g_1 + g_2 \log\left(\frac{\sigma+1}{2}\right) + g_3 \frac{\sigma \operatorname{arcsinh}\left(\sqrt{\frac{\sigma-1}{2}}\right)}{\sqrt{\sigma^2-1}}$$

$$\mathcal{D}(\sigma) = \frac{3\pi(5\sigma^2-1)\mathcal{I}(\sigma)}{8}$$

$$f_1 = \frac{210\sigma^6 - 552\sigma^5 + 339\sigma^4 - 912\sigma^3 + 3148\sigma^2 - 3336\sigma + 1151}{48(\sigma^2-1)^{3/2}}$$

$$f_2 = -\frac{35\sigma^4 + 60\sigma^3 - 150\sigma^2 + 76\sigma - 5}{8\sqrt{\sigma^2-1}}$$

$$f_3 = \frac{(2\sigma^2-3)(35\sigma^4 - 30\sigma^2 + 11)}{8(\sigma^2-1)^{3/2}}$$

$$g_1 = \frac{105\sigma^7 - 411\sigma^6 + 240\sigma^5 + 537\sigma^4 - 683\sigma^3 + 111\sigma^2 + 386\sigma - 237}{24(\sigma^2-1)^2}$$

$$g_2 = \frac{35\sigma^5 - 90\sigma^4 - 70\sigma^3 + 16\sigma^2 + 155\sigma - 62}{4(\sigma^2-1)}$$

$$g_3 = \frac{-(2\sigma^2-3)(35\sigma^5 - 60\sigma^4 - 70\sigma^3 + 72\sigma^2 + 19\sigma - 12)}{4(\sigma^2-1)^2}$$

Common concerns

- Can zero-energy radiation carries angular momentum? **Yes**
 - Is radiated angular momentum infrared finite (due to $1/r$ potential in 4D)? **Yes**
 - Are distribution functions (e.g. delta functions) well-defined? **Yes**
-

All of above can be answered in scalar theory

- Is there BMS ambiguity on angular momentum? **Maybe, I don't know**
 - But need to explain the match to ALD force in electromagnetism
 - Need to explain the match to Burke-Throne force in GR

See Snowmass white paper [Buonanno, Khalil, O'Connell, Roiban, Solon, Zeng, 2204.05194]
and many other talks in this conference

Conservative Dynamics

- Impressive progress from both traditional and new methods
- **Higher order potential**
[Bern, Cheung, Parra-Martinez, Roiban, Ruf, **CHS**, Solon, Zeng]
[Bini, Damour, Geralico] [Blumlein, Maier, Marquard, Schafer] [Dlapa, Kalin, Liu, Porto]
[Bjerrum-Bohr, Cristofoli, Damgaard, Festuccia, Plante, Vanhove] [di Vecchia, Heissenberg, Russo, Veneziano]
[Kosower, Maybee, O'Connell] [Damgaard, Haddard, Helset] [Jakobsen, Mogull, Plefka, Steinhoff]
[Brandhuber, Chen, Travaglini, Wen] [Kol, O'Connell, Telem]....
- **Spin**
[Vaidya] [Vines] [Guevara, Ochirov, Vines] [Chung, Huang, Kim, Lee] [Aoude, Haddard, Helset]
[Bern, Luna, Roiban, **CHS**, Zeng][Bern, Kosmopoulos, Luna, Roiban, Teng]
[Steinhoff, Levi] [Levi, Von Hippel, McLeod] [Liu, Porto, Yang]
[Maybee, O'Connell, Vines] [Jakobsen, Mogull, Plefka, Steinhoff] [Chiodaroli, Johansson, Pichini]...
- **Tidal effects**
[Bini, Damour][Cheung, Solon][Kalin, Liu, Porto][Aoude, Haddard, Helset]
[Bern, Parra-Martinez, Roiban, **CHS**, Sawyer] [Cheung, Shah, Solon]...

Dissipative Dynamics in Scattering

- **Double copy structure**
[Goldberger, Ridgway] [CHS] [Vazquez-Holm, Carrasco]...
- **Radiative contribution to binary deflections**
[Amati, Ciafaloni, Veneziano][Bini, Damour] [Damour] [Bini, Damour, Geralico]
[Di Vecchia, Heissenberg, Russo, Veneziano][Herrmann, Parra-Martinez, Ruf, Zeng]...
- **Radiated energy via classical or quantum (KMOC) methods**
[Herrmann, Parra-Martinez, Ruf, Zeng]
[Jakobsen, Mogull, Plefka, Steinhoff] [Mougiakakos, Riva, Vernizzi]
- **Radiated angular momentum**
[Damour][Bini, Damour, Geralico] [Di Vecchia, Heissenberg, Russo, Veneziano]
[Jakobsen, Mogull, Plefka, Steinhoff][Mougiakakos, Riva, Vernizzi][Gralla, Lobo]
- **Waveforms**
[Jakobsen, Mogull, Plefka, Steinhoff][Mougiakakos, Riva, Vernizzi]
[Cristofoli, Gonzo, Kosower, O'Connell] [Britto, Gonzo, Jehu] [Cristofoli, Gonzo, Moynihan, O'Connell, Ross]
- **Boundary to bound map**
[Cho, Kalin, Porto]