

Top quark mass corrections to NNLO double Higgs boson production

arXiv:2110.03697

SEPTA Meeting

J. Davies



December 8, 2021

Higgs self coupling

Standard Model Higgs potential:

$$V(H) = \frac{1}{2}m_H^2 H^2 + \lambda v H^3 + \frac{\lambda}{4}H^4,$$

where $\lambda = m_H^2/(2v^2) \approx 0.13$.

Want to measure λ , to determine if $V(H)$ is consistent with nature.

- ▶ Challenging! Cross-section $\approx 10^{-3} \times H$ prod.
- ▶ $-3.3 < \lambda/\lambda_{SM} < 8.5$


[CMS '21]

λ appears in various production channels:



- ▶ Gluon fusion – dominant, 10x
- ▶ $t\bar{t}$ associated production
- ▶ VBF
- ▶ H -strahlung

Higgs self coupling



Gluon Fusion

Leading order (1 loop) partonic amplitude:



$$\mathcal{M}^{\mu\nu} \sim \mathcal{A}_1^{\mu\nu}(\mathcal{F}_{tri} + \mathcal{F}_{box1}) + \mathcal{A}_2^{\mu\nu}(\mathcal{F}_{box2})$$

- ▶ \mathcal{F}_{tri} contains the dependence on λ




Form factors:

- ▶ LO: known exactly [Glover, van der Bij '88]
- ▶ Beyond LO... no fully-exact (analytic) results to date
 - ▶ numerical evaluation, expansion in various kinematic limits

$gg \rightarrow HH$ Beyond LO




$m_t \rightarrow \infty$ limit ("HEFT") used in many approximations:

► NLO "Born-improved" HTL:






$$\text{B-i: } d\sigma_{NLO}(m_t) \approx \frac{d\sigma_{LO}(m_t)}{d\sigma_{LO}(\infty)} d\sigma_{NLO}(\infty)$$

► NLO "FTapprox":



► NLO Full:

[Borowka, Greiner, Heinrich, Jones, Kerner, Schlenk, Schubert, Zirke '16]
[Baglio, Campanario, Glaus, Mühlleitner, Spira, Streicher '19]



$gg \rightarrow HH$ Beyond LO


- ▶ NLO: large- m_t + threshold expansion Padé [Gröber, Maier, Rauh '17]
- ▶ NLO: high-energy expansion [Davies, Mishima, Steinhauser, Wellmann '18,'19]
- ▶ NLO: small- p_T expansion [Bonciani, Degrassi, Giardino, Gröber '18]

- ▶ NNLO: large- m_t exp. of virt [Grigo, Hoff, Steinhauser '15][Davies, Steinhauser '19]
- ▶ HEFT + num. reals [Grazzini, Heinrich, Jones, Kallweit, Kerner, Lindert, Mazzitelli '18]

- ▶ N3LO: Wilson coefficient C_{HH} [Spira '16][Gerlach, Herren, Steinhauser '18]
- ▶ N3LO: HEFT [Chen, Li, Shao, Wang '19]


NNLO Total Cross Section

Need to compute: $\sigma_{jj}^{(2)} = \sigma_{jj,\text{virt}}^{(2)} + \sigma_{jj,\text{real}}^{(2)} + \sigma_{jj,\text{coll}}^{(2)}$ ($jj = gg, gq, q\bar{q}, qq', \dots$)



Each part is divergent, sum is finite (including also Collinear CTs).

Total XS: proceed via *Optical Theorem*. Phase-space → loop integrals:





Three- and four-particle cuts of 5-loop 2 → 2 forward diagrams.

Large- m_t expansion

Expand integrals in the region “ $m_t \gg$ other scales”.

Result: series in powers of $\{q_i \cdot q_i, q_i \cdot q_j, q_j \cdot q_j, \dots\} / m_t^2$

- ▶ $gg \rightarrow H(\rightarrow HH)$: $q_1 \cdot q_2 / m_t^2$
- ▶ $gg \rightarrow HH$: $\{q_3 \cdot q_3, q_1 \cdot q_2, q_1 \cdot q_3, q_2 \cdot q_3\} / m_t^2$



Large- m_t expansion


Expansion by sub-graph:

- ▶ sum over sub-graphs which contain m_t
- ▶ expand **hard-scaling propagators** in their small parameters

Diagrams factorize into:


- ▶ massless integrals
- ▶ expanded hard sub-graphs → **massive tadpole integrals**

Example: 2-loop $gg \rightarrow H$ diagram:



Large- m_t expansion

More explicitly:



$$\iint d^d l_1 d^d l_2 \frac{1}{l_2^2} \frac{1}{(l_2 + q_1)^2} \frac{1}{(l_2 - q_2)^2} \frac{1}{(l_1 + q_1)^2 - m_t^2} \frac{1}{(l_1 - q_2)^2 - m_t^2} \frac{1}{(l_1 - l_2)^2 - m_t^2} \rightarrow \quad (1)$$



$$\int d^d l_2 \frac{1}{l_2^2} \frac{1}{(l_2 + q_1)^2} \frac{1}{(l_2 - q_2)^2} \int d^d l_1 \left[\frac{1}{(l_1^2 - m_t^2)^3} + \frac{2(q_1 \cdot l_1 - q_2 \cdot l_1 - l_2 \cdot l_1) + l_2 \cdot l_2}{(l_1^2 - m_t^2)^4} + \dots \right] \quad (2)$$

What remains?

- ▶ massless integral over l_2
- ▶ massive tadpole integrals: $I_1^{\mu_1} I_1^{\mu_2} \dots I_1^{\mu_N} / (l_1^2 - m_t^2)^m$

Large- m_t expansion


Eg, LO \mathcal{F}_{tri} : expansion to $1/m_t^{14}$



Software:


diagram generation	qgraf	[Nogueira '93]
large- m_t expansion	q2e/exp [Harlander, Seidelsticker, Steinhauser '97] FORM 4.2 [Ruijl, Ueda, Vermaseren '17]	
tadpoles (1-3 loop)	MATAD	[Steinhauser '00]
massless integrals	FIRE 6 LiteRed	[Smirnov '19] [Lee '12]

NNLO $gg \rightarrow HH$



qgraf generates a large number of 5-loop $2 \rightarrow 2$ diagrams:

- ▶ filter for valid cuts using `gen`
 - ▶ gg channel: 16.6M → 160.1K
 - ▶ gq channel: 1.7M → 5.4K




Expansion of such graphs is very difficult, computationally.

- ▶ compute only the leading term ($1/m_t^0$) in this style

“Building blocks”


Approach: construct “building blocks”, pre-expanded effective vertices:



3600 diagrams



1 diagram



120 diagrams

Need to compute and expand various building blocks:

- ▶ ggH
- ▶ $gggH$
- ▶ $ggggH$
- ▶ $ggHH$
- ▶ $gggHH$
- ▶ $ggggHH$


Generate “building block” diagrams directly:

- ▶ gg channel: 16.6M \rightarrow 160.1K \rightarrow 4612
- ▶ gq channel: 1.7M \rightarrow 5.4K \rightarrow 336

Phase-space Integrals

After large- m_t expansion, 2- and 3-loop “phase-space integrals” remain.

IBP reduce (LiteRed) to obtain **three** and **four** particle cuts of 74 3-loop MIs belonging to 13 topologies,



and 16 2-loop MIs belonging to 3 topologies,





Compute MIs via differential equations w.r.t. $x = m_H^2/s$:

- ▶ $\partial_x \vec{I} = M(\epsilon, x) \vec{I}$
- ▶ exact solns, and via series expansion: $\delta = 1 - 4x \rightarrow 0$ (2 m_h thr.)

Phase-space Integrals

Two examples, ratio of $\delta = 1 - 4x \rightarrow 0$ expansion to exact:





- ▶ δ^{30} reproduces exact result very well up to $\delta \approx 0.9$
- ▶ $2m_t$ threshold corresponds to $\delta = 1 - m_h^2/m_t^2 = 0.48$

Exact expressions are “unpleasant” (GPLs), so produce cross sections (and compute collinear CTs) as series in δ .


NNLO $gg \rightarrow HH$, Partonic Cross Sections

Resulting expansions for channels gg ($\rho^3 = (m_h^2/m_t^2)^3$) and gq (ρ^4):



- m_t dependence is a $\sim 100\%$ correction to HEFT at NNLO

NNLO $gg \rightarrow HH$, Partonic Cross Sections



- ▶ LO \rightarrow NLO: +100%
- ▶ NLO \rightarrow NNLO: +30%

Summary

Multi-scale multi-loop amplitudes are hard:

- ▶ study them via expansions in certain kinematic limits
- ▶ direct numerical evaluation

Large- m_t expansions:

- ▶ good description of amplitudes below top quark threshold
 - ▶ above threshold: “Born improvement” – **work in progress**
- ▶ can be combined with other expansions for a better approximation
 - ▶ large- m_t + threshold Padé at NNLO – **work in progress**
- ▶ differential XS? Requires large- m_t exp. of 2-loop 2 → 3 – **planned**