

Probing double-aligned two Higgs doublet models at LHC


Michihisa Takeuchi (Osaka Univ.)

In collaboration with S. Kanemura and K. Yagyu

arXiv: 2112.13679[hep-ph] (accepted in PRD)

CP violation beyond the SM required

- Baryon Asymmetry of the Universe by EWBG : too small CPV in the SM
→ **CPV source of BSM required**
- Consider the possibility: new CPV phases exist in extended Higgs sector



Aligned CPV 2HDM and EDM

Higgs potential

$$V = -\mu_1^2 |\Phi_1|^2 - \mu_2^2 |\Phi_2|^2 - \left\{ \mu_3^2 (\Phi_1^\dagger \Phi_2) + h.c. \right\} \quad \begin{array}{l} \text{(Higgs basis)} \\ \text{[Davidson, Haber,} \\ \text{PRD72, 035004 (2005)]} \end{array}$$

$$+ \frac{1}{2} \lambda_1 |\Phi_1|^4 + \frac{1}{2} \lambda_2 |\Phi_2|^4 + \lambda_3 |\Phi_1|^2 |\Phi_2|^2 + \lambda_4 |\Phi_2^\dagger \Phi_1|^2$$

$$+ \left\{ \left[\frac{1}{2} \lambda_5 (\Phi_1^\dagger \Phi_2) + \lambda_6 |\Phi_1|^2 + \lambda_7 |\Phi_2|^2 \right] (\Phi_1^\dagger \Phi_2) + h.c. \right\}$$

Yukawa couplings

$$\mathcal{L}_{\text{yukawa}} = -\bar{Q}_L \frac{\sqrt{2}M_u}{v} (\tilde{\Phi}_1 + \zeta_u \tilde{\Phi}_2) u_R$$

$$- \bar{Q}_L \frac{\sqrt{2}M_d}{v} (\Phi_1 + \zeta_d \Phi_2) d_R$$

$$- \bar{L}_L \frac{\sqrt{2}M_e}{v} (\Phi_1 + \zeta_e \Phi_2) e_R$$

$$+ h.c.$$

Higgs basis

$$\Phi_1 = \begin{pmatrix} G^+ \\ \frac{1}{\sqrt{2}}(v + h_1^0 + iG^0) \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} H^+ \\ \frac{1}{\sqrt{2}}(h_2^0 + ih_3^0) \end{pmatrix}$$

Mass Matrix

$$\mathcal{M}^2 = v^2 \begin{pmatrix} \lambda_1 & \text{Re}[\lambda_6] & -\text{Im}[\lambda_6] \\ \text{Re}[\lambda_6] & \frac{M^2}{v^2} + \frac{1}{2}(\lambda_3 + \lambda_4 + \text{Re}[\lambda_5]) & -\frac{1}{2}\text{Im}[\lambda_5] \\ -\text{Im}[\lambda_6] & -\frac{1}{2}\text{Im}[\lambda_5] & \frac{M^2}{v^2} + \frac{1}{2}(\lambda_3 + \lambda_4 - \text{Re}[\lambda_5]) \end{pmatrix}.$$

Pheno-motivated 2 types of alignments assumed:

Higgs alignment $\lambda_6=0 (= \mu_3)$ \Rightarrow No mixing among Higgses 125GeV
 Higgs measurements indicate SM like

Yukawa alignment to avoid FCNC at tree level

\rightarrow 4 complex parameters remain $\zeta_e, \zeta_d, \zeta_u, \lambda_7$


EDM constraint :

$$-i \frac{d_e}{2} \bar{\psi}_e \sigma^{\mu\nu} \gamma^5 \psi_e F^{\mu\nu}$$

$$d_e = \sum_f^{t,b,\tau} \frac{\zeta_f}{\zeta_e} \text{ (loop diagram)} + \frac{\lambda_7}{\zeta_e} \text{ (loop diagram)} < 1.1 \times 10^{-29} e \text{ cm}$$

EW production at LHC

- In 2HDM we cannot switch off the EW production




Cross section fixed only by the masses

→ No dependence on Yukawa param.

For 6 modes

~ 10-500 fb at 13 TeV LHC
($m_H \sim 300$ GeV)

10^3 - 10^5 Events at 139fb-1



Neutral $H \rightarrow \tau\tau, bb$

Charged $H^+ \rightarrow \tau\nu, tb$

Heavy higgs also decay via

$H_2 \rightarrow Z^* H_3, H_2 \rightarrow W^{*\pm} H^\mp$

→ 4 τ lepton events expected
(BR depends on Yukawa param.)

- Latest LHC 4+ lepton (including taus) searches set very strong constraints

BR in aligned 2HDM

BR is determined by the ζ parameters

(For T parameter constraints, Charged Higgs and one of Neutral Higgses degenerated)

Easy to understand the BR behavior by separating fermion/gauge boson modes.

$$R = \frac{\sum_f \Gamma_f}{\sum_f \Gamma_f + \sum_V \Gamma_V} = \frac{1}{1 + r/\zeta^2} \simeq \sum_f BR_f$$

$$R_\tau = \frac{\Gamma_\tau}{\sum_f \Gamma_f} = \zeta_e^2 / \zeta^2 \simeq BR_\tau / R$$

$$\zeta^2 \simeq \sum_f \frac{m_f^2}{m_\tau^2} N_f^c |\zeta_f|^2$$

The corresponding parameters R^\pm, R_τ^\pm also defined for H^\pm

Neutral Higgs

$$\mathcal{B}(H \rightarrow \tau\tau) = RR_\tau$$

$$\mathcal{B}(H \rightarrow bb) = R(1 - R_\tau)$$

$$\mathcal{B}(H \rightarrow Z^{(*)}\tau\tau) = R_Z R_\tau$$


$$\mathcal{B}(H \rightarrow Z^{(*)}bb) = R_Z(1 - R_\tau)$$

$$\mathcal{B}(H \rightarrow W^{(*)}\tau\nu) = R_W R_\tau^\pm$$

$$\mathcal{B}(H \rightarrow W^{(*)}bt) = R_W(1 - R_\tau^\pm)$$

Fermion Ratio

τ -mode/Fermion Ratio



Charged Higgs

$$\mathcal{B}(H^\pm \rightarrow \tau\nu) = R^\pm R_\tau^\pm$$

$$\mathcal{B}(H^\pm \rightarrow bt) = R^\pm(1 - R_\tau^\pm)$$


$$\mathcal{B}(H^\pm \rightarrow W^{(*)}\tau\tau) = (1 - R^\pm)R_\tau$$

$$\mathcal{B}(H^\pm \rightarrow W^{(*)}bb) = (1 - R^\pm)(1 - R_\tau)$$

Current LHC bounds

Various flavor constraints make the parameter space finite

$$m_{H_3} = m_{H^\pm} = 230 \text{ GeV}, m_{H_2} = 280 \text{ GeV}, \quad |\zeta_u| = 0.1$$



At HL-LHC multilepton
 $BR_\tau \sim 0.2$ reachable

Large $\tau\tau$ BR
 constrained by LHC
 multi lepton searches

Type X interpretation:
 $(\zeta_e = \zeta_d^{-1} = \zeta_u^{-1})$


Currently,
 $\zeta_e = \tan \beta \gtrsim 5$ excluded

At HL-LHC, up to
 $\zeta_e = \tan \beta \gtrsim 1.5$
 would be sensitive

Effects of Charged Higgs spectrum


$$m_{H_3} = m_{H^\pm} \leq m_{H_2}$$

[light charged H]



$$m_{H_3} \leq m_{H_2} = m_{H^\pm}$$

[heavy charged H]



All 6 modes produced similar in size

If H^+ exists below, H_2 decay into $H^+ \rightarrow \tau \nu$: fewer leptons

\rightarrow heavier H^+ provides stronger constraints ($H \rightarrow \tau \tau$, bb , $H^+ \rightarrow \tau \nu$, tb)

At $\Delta M \sim m_W, m_Z$ the situation changes :

difference between light/heavy H^\pm more significant when open

Current LHC bounds


$m_{H_3} [\text{GeV}]$

330

$m_{H_3} = m_{H^\pm} \leq m_{H_2}$

[light charged H]

280




230

$\Delta M \sim m_W, m_Z$
situation changes

180

Multi-lepton sensitivity
weaker



180

230

280

330

$m_{H_2} [\text{GeV}]$

Type X interpretation: $\zeta_e = \tan \beta \gtrsim 2$ excluded at HL-LHC

$$\Delta m = m_{H_2} - m_{H_3}$$





Current LHC bounds

$m_{H_3} [\text{GeV}]$

330

$m_{H_3} \leq m_{H_2} = m_{H^\pm}$
[heavy charged H]

280



180


230

280

330

Heavier H+ set stronger constraints
($H \rightarrow \tau \tau, bb, H^+ \rightarrow \tau \nu, tb$)

Type X interpretation: $\zeta_e = \tan \beta \gtrsim 1$ excluded at HL-LHC



$$\Delta m = m_{H_2} - m_{H_3}$$


$\Delta M \sim m_W, m_Z$
situation changes

Multi-lepton sensitivity
stronger

$m_{H_2} [\text{GeV}]$

Current/future reaches in type X-like case

S. Kanemura, M.T., K. Yagyu [arXiv: 2112.13679 [hep-ph]]



Type X-like case, lighter charged Higgs case ($\Delta m < 0$) constrained weaker.
At HL-LHC almost all parameter space reachable below 2mt.

Mass measurements at LHC





$\text{BR}(\tau\tau) \sim 1$ already constrained

Can we use $bb\tau\tau$ mode?

H is heavy enough, collinear approx. valid

$$\vec{p}_{\nu_1} = \alpha_1 \vec{p}_{\tau_{\text{vis}1}}$$

$$\vec{p}_{\nu_2} = \alpha_2 \vec{p}_{\tau_{\text{vis}2}}$$






example) $m_{H2}, m_{H3}, m_{H^\pm} = 280, 230, 280$ GeV

Mass measurements at LHC

Can we use $b\bar{b}\tau\tau$? Large $t\bar{t}$ BG ~ 900 pb

For the masses 230 GeV, 330 GeV (signal xs ~ 10 - 50 fb)



Only 1 prong pi+ contributions
 $\tau^\pm \rightarrow \pi^\pm \nu$ (BR~10%) plotted,
other modes also usable

top BG reduced by $\min(m_{b\tau}^{\text{rec}}) > 150$ GeV: efficiency ~ 0.04 vs. 10^{-4}

We expect top BG controllable using further 2D cut

Summary

- Baryon Asymmetry of the Univ. — too small CPV phase in the SM, thus CP violation beyond the SM required
- 125GeV Higgs is SM like → **Aligned CPV 2HDM**
- discussed Heavy Higgs discovery, measurements of mass, phases at LHC
- As the first step, we identify the current/future available region by multi-lepton searches at LHC

→ (counter-intuitively) heavier H^+ cases stronger constrained
S. Kanemura, M.T., K. Yagyu [arXiv: 2112.13679 [hep-ph]]

At LHC, heavy higgs mass measurable?

→ it seems possible if they are light.

would like to investigate what can be done at HL-LHC (CPV phases)
[basis for the future ILC measurements]

- Correlation with 1st order phase transition, EW Baryon number generation


Backup

CPV phase measurement



The former study : O(1) phases compatible to EDM constraints with heavy Higgses ~ 300GeV

S. Kanemura, M. Kubota, K. Yagyu [JHEP 08 (2020) 026]

Angular distribution



Picture by [Jeans, Wilson, PRD98, 013007 (2018)]



At ILC, ζe phase measurements using azimuthal angle dist. in $H_2 H_3 \rightarrow (bb)(\tau\tau)$

$$\mathcal{M} = \mathcal{M}_{h_1 h_2}^{H \rightarrow \tau^+ \tau^-} \mathcal{M}_{h_1}^{\tau^+} \mathcal{M}_{h_2}^{\tau^-}, \quad \mathcal{M}_h^{\tau^+} \sim e^{ih\phi}$$


assuming the heavy higgs masses measured at LHC

→ At LHC, can we discover the heavy higgses? Current reaches?
 Can we measure the masses?

CPV phase measurement at LHC

Collinear approx. not accessible to azimuthal angle at τ -rest frame

Small τ -mass makes it difficult



TauDecay [Eur.Phys.J.C 73 (2013) 2489, K.Hagiwara,T. Li,K.Mawatari,J.Nakamura]

