



Contribution ID: 124

Type: **Oral Contribution**

Quadrupole excitations in self-conjugate nuclei: pushing the limit.

Wednesday 26 October 2022 16:25 (15 minutes)

In nuclei along the $N = Z$ line, as protons and neutrons occupy the same valence orbitals, proton-neutron correlation properties and quadrupole-quadrupole interactions emerges. In heavy even $N = Z$ nuclei the competition between prolate and oblate quadrupole coherence is hitherto not measured. Well-developed deformation in the upper fpg shell starts from ^{68}Se . In ^{68}Se , the intrinsic deformation of the ground-state band has been interpreted as oblate, while a prolate deformation is assigned to the excited band that soon becomes yrast. The tendency leads to the emergence of shape coexistence, which are predicted in the strongly deformed ^{72}Kr , ^{80}Zr and ^{84}Mo [1].

In this study, exploiting an ^{86}Mo radioactive beam produced at NSCL, we measured the lifetime of the first 2^+ state in ^{84}Mo and ^{86}Mo using the GRETINA array and a plunger setup. The reduced transition probability $B(E2; 2^+ \rightarrow 0^+)$ of the Mo isotopes were deduced, thereby understanding their quadrupole collectivity and deformation.

The experimental results will be presented along with their interpretation with state-of-the-art calculations using ZBM3 effective interaction

[1] A. P. Zuker, A. Poves, F. Nowacki, and S. M. Lenzi, Phys. Rev. C 92, 024320 (2015).

Primary author: RECCHIA, Francesco (Universita e INFN, Padova (IT))

Co-author: HA, Jeongsu (University and INFN Padova)

Presenter: RECCHIA, Francesco (Universita e INFN, Padova (IT))

Session Classification: P2 Nuclear Structure, Spectroscopy, and Dynamics

Track Classification: P2 Nuclear Structure, Spectroscopy, and Dynamics