

# The Dark Side of the Universe DSU2022



## Proposals for searches of scalar field dark matter using cavity resonators and capacitors

Victor V. Flambaum, Ben McAllister, Igor B. Samsonov, and Michael E. Tobar

05 December 2022

*Phys. Rev. D* 106 (2022) 5, 055037

# Introduction: The axion story

VOLUME 51, NUMBER 16

PHYSICAL REVIEW LETTERS

17 OCTOBER 1983

## Experimental Tests of the “Invisible” Axion

P. Sikivie

*Physics Department, University of Florida, Gainesville, Florida 32611*

(Received 13 July 1983)

Experiments are proposed which address the question of the existence of the “invisible” axion for the whole allowed range of the axion decay constant. These experiments exploit the coupling of the axion to the electromagnetic field, axion emission by the sun, and/or the cosmological abundance and presumed clustering of axions in the halo of our galaxy.

PHYSICAL REVIEW D

VOLUME 32, NUMBER 11

1 DECEMBER 1985

## Detection rates for “invisible”-axion searches

P. Sikivie

*Physics Department, University of Florida, Gainesville, Florida 32611*


(Received 13 May 1985)

Experiments are described to search for axions floating about in the halo of our galaxy and for axions emitted by the sun. Expressions are given for the signal strengths in these experiments.

$$\mathcal{L}_{int} = \frac{1}{4} g_{a\gamma\gamma} a F_{\mu\nu} \tilde{F}^{\mu\nu} = -g_{a\gamma\gamma} a \vec{E} \cdot \vec{B}$$



axion-photon coupling constant



# Introduction: Axion detection experiments

ADMX




CAST



Light shining through a wall



ORGAN



And others...

# Introduction: scalar field dark matter

$$\mathcal{L}_{\text{int}} = -\frac{1}{4}g_{\phi\gamma\gamma}\phi F_{\mu\nu}F^{\mu\nu} = \frac{1}{2}g_{\phi\gamma\gamma}\phi(\vec{E}^2 - \vec{B}^2)$$

 scalar-photon coupling constant

- $\phi$  is a dilaton-like scalar field motivated by e.g. string theory
- Are axion-search experiments sensitive to the scalar-photon coupling  $g_{\phi\gamma\gamma}$ ?
- How to maximize the sensitivity of these experiments to  $g_{\phi\gamma\gamma}$ ?
- New experimental techniques for searching scalar dark matter with the dilaton-like interaction?

# Theory: scalar-photon interaction

$$\mathcal{L} = \frac{1}{2}(\epsilon \vec{E}^2 - \frac{1}{\mu} \vec{B}^2) + \frac{1}{2} g_{\phi\gamma\gamma} \phi (\vec{E}^2 - \vec{B}^2)$$

dielectric constant    magnetic permeability    scalar photon interaction

$$\phi = \text{Re}[\phi_0 e^{i(\vec{p}\cdot\vec{x} - \omega t)}]$$

plane-wave  
solution

$$\phi_0 = \frac{\sqrt{2\rho_{\text{DM}}}}{m_\phi}$$

local DM density  
scalar field mass

Modified Maxwell equations

$$\begin{aligned} \nabla \cdot (\epsilon \vec{E} + g_{\phi\gamma\gamma} \phi \vec{E}) &= 0 \\ \nabla \times (\mu^{-1} \vec{B} + g_{\phi\gamma\gamma} \phi \vec{B}) - \partial_t (\epsilon \vec{E} + g_{\phi\gamma\gamma} \phi \vec{E}) &= 0 \\ \nabla \times \vec{E} + \partial_t \vec{B} &= 0 \\ \nabla \cdot \vec{B} &= 0 \end{aligned}$$



$E_0$  and  $B_0$  are static background electric and magnetic fields

$$\begin{aligned} \nabla \cdot (\epsilon \vec{E}) &= \rho_\phi \\ \nabla \times (\mu^{-1} \vec{B}) - \partial_t (\epsilon \vec{E}) &= \vec{j}_\phi \end{aligned}$$

effective charge and current densities

$$\begin{aligned} \rho_\phi &= -g_{\phi\gamma\gamma} \nabla \cdot (\phi \vec{E}_0), \\ \vec{j}_\phi &= g_{\phi\gamma\gamma} \vec{E}_0 \partial_t \phi - g_{\phi\gamma\gamma} \nabla \times (\phi \vec{B}_0) \end{aligned}$$

# Theory: Scalar-photon transformation

Photon signal power

$$P = \frac{g_{\phi\gamma\gamma}^2 \rho_{\text{DM}}}{16\pi^2 \epsilon} \int d^3 k \delta(|\vec{k}| - \omega) \left| \int_V d^3 x e^{i(\vec{p}-\vec{k}) \cdot \vec{x}} \vec{n} \times [\vec{E}_0 + \vec{\beta} \times \vec{B}_0] \right|^2$$

Compare with the power of axion-photon transformation

[P. Sikivie, Phys. Rev. D 32, 2988 (1985)]

$$P_{\text{axion}} = \frac{g_{a\gamma\gamma}^2 \rho_{\text{DM}}}{16\pi^2 \epsilon} \int d^3 k \delta(|\vec{k}| - \omega) \left| \int_V d^3 x e^{i(\vec{p}-\vec{k}) \cdot \vec{x}} \vec{n} \times [\vec{B}_0 - \vec{\beta} \times \vec{E}_0] \right|^2$$

# Theory: Scalar-photon transformation ( $E_0=0$ )


$$P = \frac{g_{\phi\gamma\gamma}^2 \rho_{\text{DM}}}{16\pi^2 \epsilon} \int d^3k \delta(|\vec{k}| - \omega) \left| \int_V d^3x e^{i(\vec{p}-\vec{k}) \cdot \vec{x}} \vec{n} \times \vec{\beta} \times \vec{B}_0 \right|^2$$

- $\beta=10^{-3}$  is the velocity of DM particles

$$|\vec{\beta} \times \vec{B}_0|^2 = \beta^2 B_0^2 \sin^2 \theta$$

- $\beta^2=10^{-6}$  is the suppression;  $\sin^2 \theta$  is responsible for signal modulation
- CAST experiment searches for axions produced in the Sun with  $\beta=1$
- CAST experiment is equally sensitive to both axions and scalars thermally produced in the Sun!


# Results: Limits from CAST Experiment



Derived from Ref. [CAST Collaboration, Nature Physics 13, 584 (2017)]  
with a graphical accuracy

# Theory: Transformation in a resonant cavity

Axion signal  
[Sikivie 1985]



Signal power

$$P_{\text{axion}} = \frac{1}{m_a} g_{a\gamma\gamma}^2 \rho_{\text{DM}} B_0^2 V C_\alpha Q_\alpha$$

Form factor

$$C_\alpha = \frac{\left| \int_V d^3x \vec{B}_0 \cdot \vec{E}_\alpha \right|^2}{B_0^2 V \int_V d^3x \epsilon \vec{E}_\alpha \cdot \vec{E}_\alpha}$$

Scalar field signal power in a cavity with electric  $E_0$  and magnetic  $B_0$  fields

$$P = \frac{1}{m_\phi} g_{\phi\gamma\gamma}^2 \rho_{\text{DM}} (B_0^2 + E_0^2) V C_\alpha Q_\alpha$$

$Q_\alpha$  is quality factor and  $C_\alpha$  is the form factor

$$C_\alpha = \frac{1}{(B_0^2 + E_0^2)V} \frac{\left| \int_V d^3x e^{i\vec{p} \cdot \vec{x}} (\vec{B}_0 \cdot \vec{B}_\alpha + \vec{E}_0 \cdot \vec{E}_\alpha) \right|^2}{\frac{1}{2} \int_V d^3x (\mu^{-1} \vec{B}_\alpha \cdot \vec{B}_\alpha + \epsilon \vec{E}_\alpha \cdot \vec{E}_\alpha)}$$

$E_\alpha$  and  $B_\alpha$  are eigenmodes of electric and magnetic fields in the cavity

# Theory: Transformation in a resonant cavity

Cavity permeated by magnetic field  $B_0$

$$P = \frac{1}{m_\phi} g_{\phi\gamma\gamma}^2 \rho_{\text{DM}} B_0^2 V C_\alpha Q_\alpha ,$$

$$C_\alpha = \frac{1}{B_0^2 V} \frac{\left| \int_V d^3x e^{i\vec{p}\cdot\vec{x}} \vec{B}_0 \cdot \vec{B}_\alpha \right|^2}{\int_V d^3x \mu^{-1} \vec{B}_\alpha \cdot \vec{B}_\alpha}$$

$$P = 1.3 \times 10^8 \text{W} \left( \frac{g_{\phi\gamma\gamma}}{\text{GeV}^{-1}} \right)^2 \left( \frac{3\mu\text{eV}}{m_\phi} \right) \left( \frac{\rho_{\text{DM}}}{0.45 \text{GeV/cm}^3} \right) \\ \times \left( \frac{B_0}{7.6 \text{T}} \right)^2 \left( \frac{V}{136 \text{L}} \right) \left( \frac{C_\alpha}{0.4} \right) \left( \frac{Q_\alpha}{30000} \right) .$$


Cavity permeated by electric field  $E_0$

$$P = \frac{1}{m_\phi} g_{\phi\gamma\gamma}^2 \rho_{\text{DM}} E_0^2 V C_\alpha Q_\alpha ,$$

$$C_\alpha = \frac{1}{E_0^2 V} \frac{\left| \int_V d^3x e^{i\vec{p}\cdot\vec{x}} \vec{E}_0 \cdot \vec{E}_\alpha \right|^2}{\int_V d^3x \epsilon \vec{E}_\alpha \cdot \vec{E}_\alpha}$$

$$P = 38 \text{W} \left( \frac{g_{\phi\gamma\gamma}}{\text{GeV}^{-1}} \right)^2 \left( \frac{3\mu\text{eV}}{m_\phi} \right) \left( \frac{\rho_{\text{DM}}}{0.45 \text{GeV/cm}^3} \right) \\ \times \left( \frac{E_0}{1 \text{MV/m}} \right)^2 \left( \frac{V}{136 \text{L}} \right) \left( \frac{C_\alpha}{0.4} \right) \left( \frac{Q_\alpha}{30000} \right) .$$


# Results: Constraints from ADMX




ADMX cavity scalar field form factor  $C_\alpha \approx 10^{-12}$

ADMX sidecar scalar field form factor  $C_\alpha \approx 10^{-8}$


Upper limits on the scalar-photon coupling constant



# Results: Cavity resonators proposals with maximized form factors to the scalar field DM




$$C_\alpha = 0.69$$



$$C_\alpha = O(1)$$



# Results: Proposal for a broadband detection with a capacitor




Effective polarization  $\vec{P}_\phi = g_{\phi\gamma\gamma} \phi \vec{E}_0$

Signal is AC voltage  $\langle V_\phi \rangle = g_{\phi\gamma\gamma} \epsilon^{-1} V_0 \frac{\sqrt{\rho_{\text{DM}}}}{m_\phi}$

Applied voltage  $V_0 \sim 600 \text{ kV}$

Estimated sensitivity of such experiment



R. Oswald et al. arXiv:2111.06883 [hep-ph].

# Summary

- Calculated the power of photon signal from scalar photon transformation in a resonant cavity
- **New limits** on the scalar-photon coupling constant from re-purposing the results of CAST and ADMX experiments
- Proposals for cavity experiments with **maximized sensitivity** to the scalar field dark matter
- A **capacitor-based broadband experiment** is proposed