

Dark Matter in the time of Primordial Black Holes



Based on:

NB & Óscar Zapata – [arXiv:2010.09725](https://arxiv.org/abs/2010.09725), [2011.02510](https://arxiv.org/abs/2011.02510), [2011.12306](https://arxiv.org/abs/2011.12306)

NB, Fazlollah Hajkarim & Yong Xu – [arXiv:2107.13575](https://arxiv.org/abs/2107.13575)

NB, Yuber Pérez-González, Yong Xu & Óscar Zapata – [arXiv:2110.04312](https://arxiv.org/abs/2110.04312)

Nicolás BERNAL



2022 Chung-Ang University - Beyond the Standard Model (CAU BSM) Workshop
February 6-10, 2022




El conocimiento
es de todos

Minciencias



Evidences for Dark Matter

Several observations indicate the existence of non-luminous Dark Matter (missing *gravitational* force) at very different scales!

- * Galactic rotation curves
- * RC in Clusters of galaxies
- * Clusters of galaxies
- * CMB anisotropies





DISTRIBUTION OF DARK MATTER IN NGC 3198







Dark Matter: WIMP

$$\frac{dn_\chi}{dt} + 3Hn_\chi = -\langle v\sigma_\chi \rangle [n_\chi^2 - (n_\chi^{\text{eq}})^2]$$



Dark Matter: WIMP vs FIMP

$$\frac{dn_\chi}{dt} + 3H n_\chi = -\langle v\sigma_\chi \rangle [n_\chi^2 - (n_\chi^{\text{eq}})^2]$$





**What if DM *only* couples to the SM
via *gravitational interactions*?**



**What if DM *only* couples to the SM
via *gravitational interactions*?**

**DM is *unavoidably* produced
by PBH Hawking evaporation!**



Primordial Black Holes

- * Density fluctuations can collapse into a PBH in the early universe
- * Lose mass by emitting *all* particles via Hawking evaporation
 - PBH have a ~black body spectrum, with temperature $T_{\text{BH}} \sim 1/M_{\text{BH}}$
 - PBHs unavoidable radiate DM!
- * If $M_{\text{in}} < 10^9 \text{ g}$, PBH completely evaporate before BBN
 - poorly constrained

Primordial Black Holes

- * Density fluctuations can collapse into a PBH in the early universe
- * Lose mass by emitting *all* particles via Hawking evaporation
 - PBH have a ~black body spectrum, with temperature $T_{\text{BH}} \sim 1/M_{\text{BH}}$
 - PBHs unavoidable radiate DM!
- * If $M_{\text{in}} < 10^9$ g, PBH completely evaporate before BBN
 - poorly constrained

Effective theory: Two free parameters

- * A single PBH characterized by its mass at formation M_{in}
(or equivalently, by the SM temperature T_{in} at formation)
- * Initial PBH energy density $\beta = \rho_{\text{BH}}/\rho_{\text{SM}}$

DM from PBHs

DM density = PBH density x # DM emitted per PBH

Number of DM particles radiated per PBH

→ Only depends on initial PBH mass!

$$N_j = \frac{15 \zeta(3)}{\pi^4} \frac{g_j C_n}{g_*(T_{\text{BH}})} \begin{cases} \left(\frac{M_{\text{in}}}{M_P}\right)^2 & \text{for } m_j \leq T_{\text{BH}}^{\text{in}} \\ \left(\frac{M_P}{m_j}\right)^2 & \text{for } m_j \geq T_{\text{BH}}^{\text{in}} \end{cases}$$

DM from PBHs

DM density = PBH density x # DM emitted per PBH

Number of DM particles radiated per PBH


→ Only depends on initial PBH mass!

$$N_j = \frac{15 \zeta(3)}{\pi^4} \frac{g_j C_n}{g_*(T_{\text{BH}})} \begin{cases} \left(\frac{M_{\text{in}}}{M_P}\right)^2 & \text{for } m_j \leq T_{\text{BH}}^{\text{in}} \\ \left(\frac{M_P}{m_j}\right)^2 & \text{for } m_j \geq T_{\text{BH}}^{\text{in}} \end{cases}$$


As PBH scale like non-relativistic matter,
they can dominate the total energy density of the universe

→ Nonstandard expansion!


DM from PBHs



DM from PBHs






DM from PBHs



1. Self-interacting DM from PBHs

Self-interacting DM from PBHs

- If DM possess sizable self-interactions:
 - DM thermalizes
 - Number-changing interactions: **2** ↔ **3**, **2** ↔ **4**...



Self-interacting DM from PBHs

- If DM possess sizable self-interactions:
 - DM thermalizes
 - Number-changing interactions: $2 \leftrightarrow 3$, $2 \leftrightarrow 4$...
- * What is the energy transferred from PBHs to DM?
- * What is the DM temperature? (kinetic equilibrium)
- * What is DM equilibrium number density? (chemical equilibrium)


Self-interacting DM from PBHs

- If DM possess sizable self-interactions:
 - DM thermalizes
 - Number-changing interactions: $2 \leftrightarrow 3$, $2 \leftrightarrow 4$...
- * What is the energy transferred from PBHs to DM?
- * What is the DM temperature? (kinetic equilibrium)
- * What is DM equilibrium number density? (chemical equilibrium)

Self-interactions:

- Increase the DM density
- Decrease the mean DM kinetic energy

Self-interacting DM from PBHs



* DM production more efficient





→ smaller β could be explored

* DM cools down

→ keV DM becomes viable

* **Model independent result**

Self-interacting DM from PBHs



* DM production more efficient

→ smaller β could be explored


* DM cools down

→ keV DM becomes viable


* **Model independent result**

2. Gravitational UV freeze-in


DM from PBHs




||



DM from PBHs




||




Gravitational UV Freeze-in

An example of UV FIMP, mediated by massless SM gravitons




Depends on:

- * DM mass and spin
 - * Reheating temperature T_{rh}
- No free couplings: M_P
- $$\Omega h^2 \sim m * (T_{\text{rh}}/M_P)^3$$




Gravitational UV Freeze-in

An example of UV FIMP, mediated by massless SM gravitons




Depends on:

- * DM mass and spin
 - * Reheating temperature T_{rh}
- No free couplings: M_P
- $$\Omega h^2 \sim m * (T_{\text{rh}}/M_P)^3$$




Gravitational DM: PBHs & UV Freeze-in



Gravitational UV freeze-in strongly constrains super heavy DM radiated by PBHs!

Gravitational DM: PBHs & UV Freeze-in




Gravitational UV freeze-in strongly constrains super heavy DM radiated by PBHs!

3. QCD Axion and PBHs

Producing Axion DM: Misalignment


Effective axion potential

$$V(\theta) = \chi(T) (1 - \cos \theta)$$



Evolution of the axion field


$$\ddot{\theta} + 3H(T)\dot{\theta} + m_a^2(T) \sin \theta = 0$$



Producing Axion DM: Misalignment


Effective axion potential

$$V(\theta) = \chi(T) (1 - \cos \theta)$$




Evolution of the axion field

$$\ddot{\theta} + 3H(T)\dot{\theta} + m_a^2(T) \sin \theta = 0$$



Axions from PBHs: Dark Radiation



As these axions are ultra-relativistic:

- can't be the cold DM
- contribute to dark radiation $\Delta N_{\text{eff}} \simeq 0.04$

Misalignment with PBHs

Even if axions radiated by PBHs can't be the DM, PBHs can have a strong impact on the DM genesis via the misalignment mechanism

Non-standard cosmological evolution:


- enhanced Hubble expansion rate
- entropy injection by PBH evaporation

Misalignment with PBHs

Even if axions radiated by PBHs can't be the DM, PBHs can have a strong impact on the DM genesis via the misalignment mechanism

Non-standard cosmological evolution:

- enhanced Hubble expansion rate
- entropy injection by PBH evaporation





Misalignment with PBHs


Even if axions radiated by PBHs can't be the DM, PBHs can have a strong impact on the DM genesis via the misalignment mechanism

Non-standard cosmological evolution:

- enhanced Hubble expansion rate
- entropy injection by PBH evaporation




Misalignment with PBHs




4. ALPs and PBHs

QCD Axion and ALPs with PBHs



Conclusions

- It's possible that DM *only* features *gravitational* interactions 
- PBHs formed in the early universe
- $0.1 \text{ g} < M_{\text{in}} < 10^9 \text{ g}$ evaporate before BBN
- PBHs could Hawking radiate the *whole* DM density
- DM masses: $1 \text{ MeV} < m_{\text{DM}} < 10^{18} \text{ GeV}$
- DM self-interactions:
 - boost DM density
Boost factors of several order of magnitude can be computed in a *model independent* way!
 - cools down DM: keV DM becomes viable
- Gravitational DM production is unavoidable!
- PBHs radiates axions → Dark radiation within the reach of CMB-S4
- Nonstandard cosmology due to PBHs have a strong impact on misalignment
 - preferred axion mass wider: lighter axions allowed
 - ALPs within the read of future ABRACADABRA, KLASH, ADMX, and DM-Radio



감사 해요 !

