




Identical-particle (pion and kaon) femtoscopy in Pb–Pb collisions at 5.02 TeV with Thermanator 2 modeled with (3+1)D viscous hydrodynamics

Pritam Chakraborty¹, Ashutosh Kumar Pandey², and Sadhana Dash¹

¹ Indian Institute of Technology Bombay, India, ² University of Tsukuba, Japan

Introduction


- Femtoscopy is a technique, used to probe the space-time geometry of the system created in heavy-ion collisions using two-particle correlation function



$$C(\mathbf{k}^*) = \frac{\int S(\mathbf{r}^*, \mathbf{k}^*) |\Psi(\mathbf{r}^*, \mathbf{k}^*)|^2}{\int S(\mathbf{r}^*, \mathbf{k}^*)}$$


Results I

- Radii for pions in all 3 directions decrease with increasing m_T in all centralities, the trends are fitted with power-law of m_T
- R in long direction falls with increasing m_T more steeply compared to out and side direction
- Radii for pions increase with multiplicity in all k_T bins, the trends are fitted with linear-function of $\langle dN_{ch}/d\eta \rangle^{1/3}$




Results III

- A common m_T scaling is approximately followed in out and side direction and slightly violated in long direction
- The slopes of 1D radii for pions, kaons and protons as the function of m_T are different.
- But a common m_T scaling is observed for different particles after scaling the radii with Lorentz boost: $f = \sqrt{(\sqrt{\gamma_T} + 2)/3}$



Results II

- Similar to pion femtoscopy, radii for kaons in all 3 directions also decrease with increasing m_T in all centralities and the trends are fitted with power-law of m_T
- R in long direction falls with increasing m_T more steeply compared to out and side direction
- Radii for kaons increase with multiplicity in all k_T bins, the trends are fitted with linear-function of $\langle dN_{ch}/d\eta \rangle^{1/3}$



Summary

- The radii decrease from most central to peripheral events and also with increasing m_T .
- For both pions and kaons, the slope of $\langle dN_{ch}/d\eta \rangle^{1/3}$ scaling of the radii is almost similar in all directions
- The m_T -scaling in out and side directions are less steeper than long direction, corresponding to the larger flow velocity
- Violation of m_T scaling corresponds to the hadronic re-scattering phase, present in the system
- A common effective scaling of 1D radii for all particles is observed after scaling the obtained radii with Lorentz factor

Reference

- P. Chakraborty et. al., Eur. Phys. J. A 57 (2021) 338.
- A. Kisiel et. al., Phys. Rev. C 90, 064914 (2014)
- P. Bozek, Phys. Rev. C 85, 034901 (2012)
- P. Chakraborty [ALICE Collaboration], Quark Matter 2022

Methodology

Pair Interaction

Bosons

Fermions

$$\Psi_{K,\pi} = 1 + \cos(2\mathbf{k}^* \cdot \mathbf{r}^*)$$

$$\Psi_p = 1 - \frac{1}{2} \cos(2\mathbf{k}^* \cdot \mathbf{r}^*)$$

Source Function

$$3D \rightarrow S(\mathbf{r}) \approx N \exp\left(-\frac{r_{out}^2}{4R_{out}^2} - \frac{r_{side}^2}{4R_{side}^2} - \frac{r_{long}^2}{4R_{long}^2}\right)$$

$$1D \rightarrow S(r^*) \approx N \exp\left(-\frac{r^{*2}}{4R_{inv}^2}\right)$$

Correlation Function

$$3D \rightarrow C(\mathbf{q}) = 1 + \lambda \exp(-R_{out}^2 q_{out}^2 - R_{side}^2 q_{side}^2 - R_{long}^2 q_{long}^2)$$

$$1D \rightarrow C(q_{inv}) = 1 + \lambda \exp(-R_{inv}^2 q_{inv}^2)$$

Analysis Details

- This study focuses on the femtoscopic study of identical pions, kaons and protons in Pb-Pb events simulated using (3+1)D viscous hydrodynamics, coupled with THERMINATOR 2 at $\sqrt{s_{NN}} = 5.02$ TeV
- The hadronic rescattering has not been considered in this model.
- The source-sizes in 3 dimensions (R_{out} , R_{side} and R_{long}) and 1 dimension (R_{inv}) are estimated in as the function of centrality (expressed as $\langle dN_{ch}/d\eta \rangle^{1/3}$) and k_T (and m_T) of the pair