



Contribution ID: 595

Type: Theory poster

Constraining 3-3-1 Models at the LHC and Future Hadron Colliders

Tuesday 17 May 2022 19:00 (1 hour)

In this work, we derive lower mass bounds on the Z' gauge boson based on the dilepton data from LHC with 13 TeV of center-of-mass energy, and forecast the sensitivity of the High-Luminosity-LHC with $L = 3000 fb^{-1}$, the High-Energy LHC with $\sqrt{s} = 27$ TeV, and also at the Future Circular Collider with $\sqrt{s} = 100$ TeV. We take into account the presence of exotic and invisible decays of the Z' gauge boson to find a more conservative and robust limit, different from previous studies. We investigate the impact of these new decays channels for several benchmark models in the scope of two different 3-3-1 models. We found that in the most constraining cases, LHC with $139 fb^{-1}$ can impose $m_{Z'} > 4$ TeV. Moreover, we forecast HL-LHC, HE-LHC, and FCC bounds that yield $m_{Z'} > 5.8$ TeV, $m_{Z'} > 9.9$ TeV, and $m_{Z'} > 27$ TeV, respectively. Lastly, put our findings into perspective with dark matter searches to show the region of parameter space where a dark matter candidate with the right relic density is possible.

Authors: Ms SÁNCHEZ VILLAMIZAR, Yoxara (IIP & DFTE - Federal University of Rio Grande do Norte (UFRN)); ALVES, Alexandre (Universidade Federal de São Paulo / UNIFESP); Ms OVIEDO-TORRES, Yohan; QUEIROZ, FARINALDO; KOVALENKO, Sergey (Universidad Andres Bello (CL)); Ms DUARTE, Laura

Presenter: Ms SÁNCHEZ VILLAMIZAR, Yoxara (IIP & DFTE - Federal University of Rio Grande do Norte (UFRN))

Session Classification: Poster Session I

Track Classification: TeV-Scale BSM