

---

# SMEFT status from a theoretical perspective

*Going Beyond the Dim-6 SMEFT in the high luminosity/energy era*

---

*Lepton-Photon 2023,  
Melbourne, Australia*

Rick Sandeepan Gupta  
Tata Institute of Fundamental Research (Mumbai)

---



# PRECISION HIGGS PHYSICS

- Studying the properties of the Higgs and other electroweak states is an obvious goal for particle physicists today.
- **Precision Higgs physics** has matured into a sophisticated field in the last 11 years since Higgs discovery
- The **theoretical framework** that has become **standard** is the **dimension 6 Standard Model Effective Theory (SMEFT)**

$$\mathcal{L} = \mathcal{L}_{SM} + \sum_{i=1}^{59} \frac{c_i}{\Lambda^2} \mathcal{O}_i^6 + \dots$$

*model independent way to parametrise effect of heavy particles*



---

# OVERVIEW

---

- EFTs in the context of BSM studies have seen intense activity recently on both the theory and experimental side.
  - Many new theoretical developments:
    1. HEFT vs SMEFT
    2. Dimension 8
    3. Amplitude approach
    4. Differential/multivariate signatures of EFT operators
    5. Positivity bounds
    6. Many technical breakthroughs in operator counting, matching, RG etc
  - I will cover the first 4 topics, indeed positivity constraints have so far been mostly restricted to dimension 8 operators
-



---

# OVERVIEW

---

- EFTs in the context of BSM studies have seen intense activity recently on both the theory and experimental side.
  - Many new theoretical developments:
    1. HEFT vs SMEFT
    2. Dimension 8
    3. Amplitude approach
    4. Differential/multivariate signatures of EFT operators
    5. Positivity bounds
    6. Many technical breakthroughs in operator counting, matching, RG etc
  - I will **cover the first 4 topics**, indeed positivity constraints have so far been mostly restricted to dimension 8 operators
-



---

# HIGHER LUMINOSITIES/ENERGIES AT FUTURE COLLIDERS

---

- We are now entering the era of **higher luminosities/energies** with many proposed future colliders.
- These have the potential to achieve a **new level of precision** in Higgs physics





- 
- 
- Can we say anything qualitatively new with all this new data ?
  - Or just improve our existing constraints on EFT couplings ?
-

- 
- 
- Can we say anything qualitatively new with all this new data ? ✓
  - Or just improve our existing constraints on EFT couplings ?
-




---

PRESENT STATUS → HIGHER LUMINOSITIES/  
ENERGIES

---

Experimental observables used:


Mostly rates & some  
one dimensional  
distributions



Fully differentiable  
observables, Multivariate  
distributions, Machine  
learning

Theoretical framework used:

Dimension 6 SMEFT




Dimension 8 SMEFT, HEFT,  
Amplitudes

*Go beyond SMEFT assumptions, test them !*

---




# BEYOND SMEFT





# BEYOND SMEFT



*If we do establish that  
BSM deviations lie in the  
SMEFT subspace  
This will be a way to  
**test the SMEFT!***



# BEYOND SMEFT I: HEFT

- SMEFT not always the right choice.

- SMEFT: Observed 125 GeV  $h$  and goldstones eaten by  $W, Z$  make a doublet.



***Essence of Higgs mechanism!***

$$\begin{pmatrix} G^\pm \\ iG_0 + v + h/\sqrt{2} \end{pmatrix}$$

- HEFT: More general, includes SMEFT as a special case. No connection assumed between  $h$  and goldstones/VEV.

$$(G^\pm, v + iG_0) + h$$



$$U = \exp(2iX_i\pi_i/v)$$



---

# BEYOND SMEFT I: HEFT

---

- Lot of recent work:
1. Alonso, Jenkins & Manohar(2016)
  2. Alonso, Jenkins & Manohar(2016)
  3. Falkowski & Rattazzi (2019)
  4. Cohen, Craig, Lu & Sutherland (2020)
  5. Banta, Cohen, Craig, Lu & Sutherland (2021)
  6. Alonso & West (2021)
  7. Alonso & West (2022)
  8. Bertuzzo, Grojean & RSG (in prep)
-



---

# HEFT BUT NOT SMEFT: UV SCENARIOS

---

- Recent work shows that **UV theories** that **map to HEFT and not SMEFT** are ubiquitous. **Whenever we integrate out states that get a majority of their mass from electroweak VEV, theory maps to HEFT. Eg. 4th generation fermions, 2HDMS etc**
- Such particles were dubbed ‘Loryons’ by Cohen et al.
- **Parameter space for many such UV scenarios wide open.**

Falkowski & Rattazzi (2019),  
Cohen, Craig, Lu & Sutherland (2020)

---



---

But **what is the difference** between these 2 expansions ?

How do we **distinguish** these 2 possibilities **experimentally** ?

Answer: The **difference** becomes **clear** at the level of **anomalous couplings**

---




# ANOMALOUS COUPLINGS

Anomalous couplings are **QCD & EM invariant Lagrangian terms**

(1) Higgs observables (20):

$$h W_{\mu\nu}^+ W^{-\mu\nu}$$


$$h Z_\mu \bar{f}_{L,R} \gamma^\mu f_{L,R}$$



(2) Electroweak precision observables (9):

$$Z_\mu \bar{f}_{L,R} \gamma^\mu f_{L,R}$$

$$W_\mu^+ \bar{\nu}_L \gamma^\mu e_L$$




(3) Triple and Quartic Gauge couplings (3+4):

$$g_1^Z c_{\theta_W} Z^\mu \left( W^{+\nu} \hat{W}_{\mu\nu}^- - W^{-\nu} \hat{W}_{\mu\nu}^+ \right)$$

$$\kappa_\gamma s_{\theta_W} \hat{A}^{\mu\nu} W_\mu^+ W_\nu^-$$

$$\lambda_\gamma s_{\theta_W} \hat{A}^{\mu\nu} \hat{W}_\mu^{-\rho} \hat{W}_{\rho\nu}^+$$





---

# ANOMALOUS COUPLINGS: HEFT VS SMEFT

---

- In SMEFT some linear combination of anomalous couplings are suppressed by powers of wrt HEFT.

- Eg. :  $Vll$  couplings ( $l$  is a lepton)

$$SM + \delta g_{e_L}^Z Z_\mu \bar{e}_L \gamma^\mu e_L + \delta g_{e_R}^Z Z_\mu \bar{e}_R \gamma^\mu e_L + \delta g_{\nu_L}^Z Z_\mu \bar{\nu}_L \gamma^\mu \nu_L + \delta g_L^W (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.)$$

4 anomalous couplings

- In HEFT all these arise independently at  $\mathcal{O}(v^2/\Lambda^2)$

- In SMEFT 3 are  $\mathcal{O}(v^2/\Lambda^2)$  and  $\delta g_L^W = \frac{\cos \theta_W (\delta g_{\nu_L}^Z - \delta g_{e_L}^Z)}{\sqrt{2}} = \mathcal{O}(v^4/\Lambda^4)$



# CORRELATIONS BETWEEN W/Z COUPLING DEVIATIONS

- 4 anomalous couplings related to  $Zff$ ,  $Wff$  deviations

$$SM + \delta g_{e_L}^Z Z_\mu \bar{e}_L \gamma^\mu e_L + \delta g_{e_R}^Z Z_\mu \bar{e}_R \gamma^\mu e_L + \delta g_{\nu_L}^Z Z_\mu \bar{\nu}_L \gamma^\mu \nu_L + \delta g_L^W (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.)$$

- At D6 level only 3 operators break these D4 predictions at  $\mathcal{O}(v^2/\Lambda^2)$

$$\mathcal{O}_{e_R} = iH^\dagger \overleftrightarrow{D}H \bar{e}_R \gamma^\mu e_R \quad \mathcal{O}_{L1} = iH^\dagger \overleftrightarrow{D}H \bar{L} \gamma^\mu L \quad \mathcal{O}_{L3} = iH^\dagger \sigma^a \overleftrightarrow{D}H \bar{L} \sigma^a \gamma^\mu L$$

- For leptons four anomalous couplings and only 3 operators so 1 prediction:

$$\delta g_L^W = \frac{\cos \theta_W (\delta g_{\nu_L}^Z - \delta g_{e_L}^Z)}{\sqrt{2}} = 0$$



---

# BREAKING OF D6 CORRELATION AT D8

---

- At **D8 level** another  $SU(2) \times U(1)$  invariant operator **breaks D6 prediction at  $\mathcal{O}(v^4/\Lambda^4)$**

$$\mathcal{O}_{L3'} = iH^\dagger \overleftrightarrow{D}_\mu H (H^\dagger \sigma^a H) \bar{L} \sigma^a \gamma^\mu L$$

- So of the **4 D4 predictions 3 are broken at  $\mathcal{O}(v^2/\Lambda^2)$  and 1 at  $\mathcal{O}(v^4/\Lambda^4)$**

$$\delta g_L^W - \frac{\cos \theta_W (\delta g_{\nu_L}^Z - \delta g_{e_L}^Z)}{\sqrt{2}} = \mathcal{O}(v^4/\Lambda^4)$$

- At D6 level there were 3 independent couplings, **at D8 we unblock a further observable/ open a 4th BSM primary**



# BREAKING OF D6 CORRELATION AT D8

- At **D8 level** another  $SU(2) \times U(1)$  invariant operator **breaks D6 prediction** at  $\mathcal{O}(v^4/\Lambda^4)$

$\mathcal{O}(v^4/\Lambda^4)$  **D6 SMEFT Prediction: Once  $Z$  coupling deviations are measured,  $W$  coupling deviation completely fixed!**

- So of the 4 **D4 predictions** at  $\mathcal{O}(v^4/\Lambda^4)$

$$\delta g_L^W - \frac{\cos \theta_W (\delta g_{\nu_L}^Z - \delta g_{e_L}^Z)}{\sqrt{2}} = \mathcal{O}(v^4/\Lambda^4)$$

- At D6 level there were 3 independent couplings, **at D8 we unblock a further observable/ open a 4th BSM primary**



# SMEFT VS HEFT

HEFT

4 anomalous couplings

SMEFT

All 4  
Couplings  
 $\mathcal{O}(v^2/\Lambda^2)$

3  $\mathcal{O}(v^2/\Lambda^2)$   
linear combinations

1  $\mathcal{O}(v^4/\Lambda^4)$   
linear combinations



# CONSIDER ALL MAJOR HIGGS PROCESSES

We can extend this approach to all the **anomalous couplings** that contribute to these Higgs production/ decay processes

## List of Processes

$$gg \rightarrow h, hh, hhh$$

$$VV \rightarrow h, hh, hV$$

$$ff(ff') \rightarrow Zh(W h)$$

$$h \rightarrow bb, cc, \tau\tau, \mu\mu$$

$$h \rightarrow \gamma\gamma, Z\gamma$$

$$h \rightarrow Wl\nu, Wjj, Zll, Zjj$$



# 52 ANOMALOUS COUPLINGS

We find that these anomalous couplings → that contribute to these processes\*

\*under certain assumptions

List of Anomalous Couplings
<b>Vff couplings (9)</b>
$\Delta\mathcal{L}_{Vff} = \sum_f \delta g_f^Z Z_\mu \bar{f} \gamma^\mu f_{L,R} + \delta g_{e\nu}^W (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.) + \delta g_{ud}^W (W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c.)$
<b>Anomalous TGC (4)</b>
$\Delta\mathcal{L}_{TGC} = igc_{\theta_W} [\delta g_1^Z Z_\mu (W_\nu^+ \mathcal{W}^{-\mu\nu} - W_\nu^- \mathcal{W}^{+\mu\nu}) + \delta \kappa^Z W_\mu^+ W_\nu^- Z^{\mu\nu}]$ $+ ie \delta \kappa^\gamma W_\mu^+ W_\nu^- A^{\mu\nu} + g_5 \epsilon^{\mu\nu\rho\sigma} W_\mu^+ \overleftrightarrow{D}_\rho W_\nu^- Z_\sigma$
<b>Anomalous QGC (5)</b>
$\Delta\mathcal{L}_{QGC} = g^2 c_{\theta_W}^2 [\delta g_{ZZ1}^Q Z^\mu Z^\nu W_\mu^- W_\nu^+ - \delta g_{ZZ2}^Q Z^\mu Z_\mu W^{-\nu} W_\nu^+] + \frac{g^2}{4c_{\theta_W}^4} h_{ZZ}^Q (Z^\mu Z_\mu)^2$ $+ \frac{g^2}{2} [\delta g_{WW1}^Q W^{-\mu} W^{+\nu} W_\mu^- W_\nu^+ - \delta g_{WW2}^Q (W^{-\mu} W_\mu^+)^2]$
<b>Single Higgs (19)</b>
$\Delta\mathcal{L}_h = \delta g_{VV}^h h [W^{+\mu} W_\mu^- + \frac{1}{2c_{\theta_W}^2} Z^\mu Z_\mu] + \delta g_{ff}^h (h \bar{f}_L f_R + h.c.) + \delta g_{ZZ}^h h \frac{Z^\mu Z_\mu}{2c_{\theta_W}^2}$ $\sum_f g_{Zff}^h \frac{h}{v} Z_\mu \bar{f} \gamma^\mu f + g_{WQ}^h \frac{h}{v} (W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c.) + g_{WL}^h \frac{h}{v} (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.)$ $+ \kappa_{ZZ}^h \frac{h}{2v} Z^{\mu\nu} Z_{\mu\nu} + \kappa_{\gamma\gamma}^h \frac{h}{2v} A^{\mu\nu} A_{\mu\nu} + \kappa_{Z\gamma}^h \frac{h}{v} A^{\mu\nu} Z_{\mu\nu} + \kappa_{WW}^h \frac{h}{v} \mathcal{W}^{+\mu\nu} \mathcal{W}_{\mu\nu}^-$ $+ \kappa_{GG}^h \frac{h}{2v} G^{A\mu\nu} G_{\mu\nu}^A$
<b>hV<sup>3</sup> couplings (5)</b>
$\Delta\mathcal{L}^{hV^3} = igc_{\theta_W} \frac{h}{v} [g_{Z1}^{hV^3} Z_\mu (W_\nu^+ \mathcal{W}^{-\mu\nu} - W_\nu^- \mathcal{W}^{+\mu\nu}) + \kappa_Z^{hV^3} W_\mu^+ W_\nu^- Z^{\mu\nu}]$ $+ ie \kappa_\gamma^{hV^3} \frac{h}{v} W_\mu^+ W_\nu^- A^{\mu\nu} + g_5^{hV^3} \frac{h}{v} \epsilon^{\mu\nu\rho\sigma} W_\mu^+ \overleftrightarrow{D}_\rho W_\nu^- Z_\sigma$ $+ ig_{\partial h Z}^{hV^3} \frac{g}{2c_{\theta_W}} \frac{\partial_\mu h}{v} Z_\nu (W_\mu^+ W^{-\nu} - W_\mu^- W^{+\nu})$
<b>h<sup>2</sup>V<sup>2</sup> couplings (8)</b>
$\Delta\mathcal{L}^{hh} = \delta g_{VV}^{hh} \frac{h^2}{2} [W^{+\mu} W_\mu^- + \frac{1}{2c_{\theta_W}^2} Z^\mu Z_\mu] + \delta g_{ZZ}^{hh} \frac{h^2}{2} \frac{Z^\mu Z_\mu}{2c_{\theta_W}^2}$ $+ g_{Z1}^{hh} \frac{(\partial_\nu h)^2}{v^2} + g_{Z2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} \frac{Z^\mu Z^\nu}{c_{\theta_W}^2}$ $+ g_{W1}^{hh} \frac{(\partial_\nu h)^2}{v^2} W^{+\mu} W_\mu^- + g_{W2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} (W^{+\mu} W^{-\nu} + h.c.)$ $+ \kappa_{WW}^{hh} \frac{h^2}{2v^2} \mathcal{W}^{+\mu\nu} \mathcal{W}_{\mu\nu}^- + \kappa_{ZZ}^{hh} \frac{h^2}{4v^2} Z^{\mu\nu} Z_{\mu\nu}$
<b>h<sup>2</sup>G<sup>2</sup> couplings (1)</b>
$\Delta\mathcal{L}_{GG}^{hh} = \kappa_{GG}^{hh} \frac{h^2}{4v^2} G^{A\mu\nu} G_{\mu\nu}^A$
<b>Higgs potential corrections (2)</b>
$\Delta\mathcal{L}^{h^n} = -\delta\lambda_3 v h^3 - \delta\lambda_4 \frac{h^4}{4}$



# SMEFT VS HEFT

HEFT

52 anomalous couplings

SMEFT

All linear  
combinations  
 $\mathcal{O}(v^2/\Lambda^2)$

17  $\mathcal{O}(v^2/\Lambda^2)$   
linear combinations

23  $\mathcal{O}(v^4/\Lambda^4)$   
linear combinations

12  $\mathcal{O}(v^6/\Lambda^6)$   
linear combinations



# 35 LINEAR COMBINATIONS $\leq \mathcal{O}(v^4/\Lambda^4)$

W/Z-decays (2)	
$\delta^8 g_Q^W$	$\delta g_Q^W - \frac{c_{\theta W}}{\sqrt{2}}(\delta g_{u_L}^Z - \delta g_{d_L}^Z)$
$\delta^8 g_L^W$	$\delta g_L^W - \frac{c_{\theta W}}{\sqrt{2}}(\delta g_{\nu_L}^Z - \delta g_{e_L}^Z)$
TGC (2)	
$\delta^8 \kappa^Z$	$\delta \kappa^Z - \delta g_1^Z + t_{\theta W}^2 \delta \kappa^\gamma$
$g_5$	New Structure
QGC (5)	
$\delta^8 g_{WW1}^Q$	$\delta g_{WW1}^Q - 2c_{\theta W}^2 \delta g_1^Z$
$\delta^8 g_{WW2}^Q$	$\delta g_{WW2}^Q - 2c_{\theta W}^2 \delta g_1^Z$
$\delta^8 g_{ZZ1}^Q$	$\delta g_{ZZ1}^Q - 2\delta g_1^Z$
$\delta^8 g_{ZZ2}^Q$	$\delta g_{ZZ2}^Q - 2\delta g_1^Z$
$h^{ZZ}$	New Structure
Higgs Production and decay (12)	
$\delta^8 g_{ZZ}^h$	$\delta g_{ZZ}^h - (\delta g_1^Z s_{\theta W}^2 - \delta \kappa^\gamma t_{\theta W}^2) g^2 v$
$\delta^8 \kappa_{WW}^h$	$\kappa_{WW}^h - \delta \kappa^\gamma - \frac{c_{\theta W}}{s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h$
$\delta^8 \kappa_{ZZ}^h$	$\kappa_{ZZ}^h - \frac{1}{c_{\theta W}^2} \delta \kappa^\gamma - \frac{c_{2\theta W}^2}{c_{\theta W} s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h$
$\delta^8 g_{WL}^h$	$g_{WL}^h - \sqrt{2} c_{\theta W} (\delta g_{\nu_L}^Z - \delta g_{e_L}^Z - g c_{\theta W} \delta g_1^Z) + 2\delta g_L^W \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{WQ}^h$	$g_{WQ}^h - \sqrt{2} c_{\theta W} (\delta g_{u_L}^Z - \delta g_{d_L}^Z - g c_{\theta W} \delta g_1^Z) + 2\delta g_Q^W \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{Zf}^h$	$g_{Zf}^h - \frac{2g}{c_{\theta W}} Y_f t_{\theta W}^2 \delta \kappa^\gamma - 2\delta g_f^Z + \frac{2g}{c_{\theta W}} (T_3^f c_{\theta W}^2 + Y_f s_{\theta W}^2) \delta g_1^Z + 2c_{2\theta W} \delta g_f^Z \delta g_1^Z$
CorrectionNew Structure to Higgs potential (1)	
$\delta^8 \lambda_4$	$\delta \lambda_4 - 6\delta \lambda_3 + \frac{4}{g^2} \left( \frac{\delta g_{VV}^h}{v} + g^2 c_{\theta W}^2 \delta g_1^Z \right) \left( \frac{m_h^2}{3v^2} + 3\delta \lambda_3 \right)$
$h^2 G^2$ coupling (1)	
$\delta^8 \kappa_{GG}^{hh}$	$\kappa_{GG}^{hh} - \kappa_{GG}^h + \frac{\kappa_{GG}^h}{2} \alpha_r$
$h^2 V^2$ couplings (8)	
$\delta^8 \kappa_{WW}^{hh}$	$\kappa_{WW}^{hh} - \delta \kappa^\gamma - \frac{c_{\theta W}}{s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h + \frac{\kappa_{WW}^h}{2} \alpha_r$
$\delta^8 \kappa_{ZZ}^{hh}$	$\kappa_{ZZ}^{hh} - \frac{1}{c_{\theta W}^2} \delta \kappa^\gamma - \frac{c_{2\theta W}^2}{c_{\theta W} s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h + \frac{\kappa_{ZZ}^h}{2} \alpha_r$
$\delta^8 g_{VV}^{hh}$	$\delta g_{VV}^{hh} - \frac{4\delta g_{VV}^h}{v} + g^2 \delta g_1^Z c_{\theta W}^2 + \frac{\delta g_{VV}^h}{2v} \alpha_r + 4 \left( g^2 \kappa_{WW}^h + 2 \frac{\delta g_{VV}^h}{v} \right) \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{ZZ}^{hh}$	$\delta g_{ZZ}^{hh} - 5(\delta g_1^Z s_{\theta W}^2 - \delta \kappa^\gamma t_{\theta W}^2) g^2 + \frac{\delta g_{ZZ}^h}{2v} \alpha_r + 4(\kappa_{Z\gamma}^h s_{2\theta W} + \kappa_{ZZ}^h c_{2\theta W} - \kappa_{WW}^h) g^2 \delta g_1^Z$
$g_{W1}^{hh}$	New Structure
$g_{W2}^{hh}$	New Structure
$g_{Z1}^{hh}$	New Structure
$g_{Z2}^{hh}$	New Structure
$hV^3$ couplings (5)	
$\delta^8 g_{Z1}^{hV^3}$	$g_{Z1}^{hV^3} + \frac{2}{c_{\theta W}^2} \left( \frac{\kappa_{Z\gamma}^h}{t_{\theta W}} + \delta \kappa^\gamma + \kappa_{\gamma\gamma}^h \right) + 4 \left( \frac{c_{2\theta W}^2}{2c_{\theta W}^2} + 1 \right) (\delta g_1^Z)^2 c_{\theta W}^2$
$\delta^8 \kappa_\gamma^{hV^3}$	$\kappa_\gamma^{hV^3} + \frac{2}{t_{\theta W}} \kappa_{Z\gamma}^h + 2\kappa_{\gamma\gamma}^h + 4\delta \kappa^\gamma \delta g_1^Z c_{\theta W}^2$
$\delta^8 \kappa_Z^{hV^3}$	$\kappa_Z^{hV^3} + \frac{2}{c_{\theta W}^2} \delta \kappa^\gamma + \frac{2}{t_{\theta W}} \kappa_{Z\gamma}^h + 2\kappa_{\gamma\gamma}^h + 4 \left( \left( \frac{c_{2\theta W}^2}{2c_{\theta W}^2} + 1 \right) \delta \kappa^Z + t_{\theta W}^2 \delta \kappa^\gamma \right) \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{\partial h Z}^{hV^3}$	$g_{\partial h Z}^{hV^3} + 4(\delta \kappa^Z c_{2\theta W} + 2\delta \kappa^\gamma s_{\theta W}^2 - \delta g_1^Z c_{\theta W}^2) \delta g_1^Z c_{\theta W}^2$
$g_5^{hV^3}$	New Structure



# 35 LINEAR COMBINATIONS=0

at  $\mathcal{O}(v^2/\Lambda^2)$

D6 level SMEFT  
predictions !

W/Z-decays (2)	
$\delta^8 g_Q^W$	$\delta g_Q^W - \frac{c_{\theta W}}{\sqrt{2}}(\delta g_{u_L}^Z - \delta g_{d_L}^Z)$
$\delta^8 g_L^W$	$\delta g_L^W - \frac{c_{\theta W}}{\sqrt{2}}(\delta g_{\nu_L}^Z - \delta g_{e_L}^Z)$
TGC (2)	
$\delta^8 \kappa^Z$	$\delta \kappa^Z - \delta g_1^Z + t_{\theta W}^2 \delta \kappa^\gamma$
$g_5$	New Structure
QGC (5)	
$\delta^8 g_{WW1}^Q$	$\delta g_{WW1}^Q - 2c_{\theta W}^2 \delta g_1^Z$
$\delta^8 g_{WW2}^Q$	$\delta g_{WW2}^Q - 2c_{\theta W}^2 \delta g_1^Z$
$\delta^8 g_{ZZ1}^Q$	$\delta g_{ZZ1}^Q - 2\delta g_1^Z$
$\delta^8 g_{ZZ2}^Q$	$\delta g_{ZZ2}^Q - 2\delta g_1^Z$
$h^{ZZ}$	New Structure
Higgs Production and decay (12)	
$\delta^8 g_{ZZ}^h$	$\delta g_{ZZ}^h - (\delta g_1^Z s_{\theta W}^2 - \delta \kappa^\gamma t_{\theta W}^2) g^2 v$
$\delta^8 \kappa_{WW}^h$	$\kappa_{WW}^h - \delta \kappa^\gamma - \frac{c_{\theta W}}{s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h$
$\delta^8 \kappa_{ZZ}^h$	$\kappa_{ZZ}^h - \frac{1}{c_{\theta W}^2} \delta \kappa^\gamma - \frac{c_{2\theta W}}{c_{\theta W} s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h$
$\delta^8 g_{WL}^h$	$g_{WL}^h - \sqrt{2} c_{\theta W} (\delta g_{\nu_L}^Z - \delta g_{e_L}^Z - g c_{\theta W} \delta g_1^Z) + 2\delta g_L^W \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{WQ}^h$	$g_{WQ}^h - \sqrt{2} c_{\theta W} (\delta g_{u_L}^Z - \delta g_{d_L}^Z - g c_{\theta W} \delta g_1^Z) + 2\delta g_Q^W \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{Zf}^h$	$g_{Zf}^h - \frac{2g}{c_{\theta W}} Y_f t_{\theta W}^2 \delta \kappa^\gamma - 2\delta g_f^Z + \frac{2g}{c_{\theta W}} (T_3^f c_{\theta W}^2 + Y_f s_{\theta W}^2) \delta g_1^Z + 2c_{2\theta W} \delta g_f^Z \delta g_1^Z$
CorrectionNew Structure to Higgs potential (1)	
$\delta^8 \lambda_4$	$\delta \lambda_4 - 6\delta \lambda_3 + \frac{4}{g^2} \left( \frac{\delta g_{VV}^h}{v} + g^2 c_{\theta W}^2 \delta g_1^Z \right) \left( \frac{m_h^2}{3v^2} + 3\delta \lambda_3 \right)$
$h^2 G^2$ coupling (1)	
$\delta^8 \kappa_{GG}^{hh}$	$\kappa_{GG}^{hh} - \kappa_{GG}^h + \frac{\kappa_{GG}^h}{2} \alpha_r$
$h^2 V^2$ couplings (8)	
$\delta^8 \kappa_{WW}^{hh}$	$\kappa_{WW}^{hh} - \delta \kappa^\gamma - \frac{c_{\theta W}}{s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h + \frac{\kappa_{WW}^h}{2} \alpha_r$
$\delta^8 \kappa_{ZZ}^{hh}$	$\kappa_{ZZ}^{hh} - \frac{1}{c_{\theta W}^2} \delta \kappa^\gamma - \frac{c_{2\theta W}}{c_{\theta W} s_{\theta W}} \kappa_{Z\gamma}^h - \kappa_{\gamma\gamma}^h + \frac{\kappa_{ZZ}^h}{2} \alpha_r$
$\delta^8 g_{VV}^{hh}$	$\delta g_{VV}^{hh} - \frac{4\delta g_{VV}^h}{v} + g^2 \delta g_1^Z c_{\theta W}^2 + \frac{\delta g_{VV}^h}{2v} \alpha_r + 4 \left( g^2 \kappa_{WW}^h + 2 \frac{\delta g_{VV}^h}{v} \right) \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{ZZ}^{hh}$	$\delta g_{ZZ}^{hh} - 5(\delta g_1^Z s_{\theta W}^2 - \delta \kappa^\gamma t_{\theta W}^2) g^2 + \frac{\delta g_{ZZ}^h}{2v} \alpha_r + 4(\kappa_{Z\gamma}^h s_{2\theta W} + \kappa_{ZZ}^h c_{2\theta W} - \kappa_{WW}^h) g^2 \delta g_1^Z$
$g_{W1}^{hh}$	New Structure
$g_{W2}^{hh}$	New Structure
$g_{Z1}^{hh}$	New Structure
$g_{Z2}^{hh}$	New Structure
$hV^3$ couplings (5)	
$\delta^8 g_{Z1}^{hV^3}$	$g_{Z1}^{hV^3} + \frac{2}{c_{\theta W}^2} \left( \frac{\kappa_{Z\gamma}^h}{t_{\theta W}} + \delta \kappa^\gamma + \kappa_{\gamma\gamma}^h \right) + 4 \left( \frac{c_{2\theta W}}{2c_{\theta W}^2} + 1 \right) (\delta g_1^Z)^2 c_{\theta W}^2$
$\delta^8 \kappa_\gamma^{hV^3}$	$\kappa_\gamma^{hV^3} + \frac{2}{t_{\theta W}} \kappa_{Z\gamma}^h + 2\kappa_{\gamma\gamma}^h + 4\delta \kappa^\gamma \delta g_1^Z c_{\theta W}^2$
$\delta^8 \kappa_Z^{hV^3}$	$\kappa_Z^{hV^3} + \frac{2}{c_{\theta W}^2} \delta \kappa^\gamma + \frac{2}{t_{\theta W}} \kappa_{Z\gamma}^h + 2\kappa_{\gamma\gamma}^h + 4 \left( \left( \frac{c_{2\theta W}}{2c_{\theta W}^2} + 1 \right) \delta \kappa^Z + t_{\theta W}^2 \delta \kappa^\gamma \right) \delta g_1^Z c_{\theta W}^2$
$\delta^8 g_{\partial h Z}^{hV^3}$	$g_{\partial h Z}^{hV^3} + 4(\delta \kappa^Z c_{2\theta W} + 2\delta \kappa^\gamma s_{\theta W}^2 - \delta g_1^Z c_{\theta W}^2) \delta g_1^Z c_{\theta W}^2$
$g_5^{hV^3}$	New Structure




# 12 LINEAR COMBINATIONS $\leq \mathcal{O}(v^6/\Lambda^6)$

$$\begin{aligned} & \delta^8 \kappa_{WW} - c_{\theta_W}^2 \delta^8 \kappa_{ZZ} - 2c_{\theta_W}^2 \delta^8 \kappa_Z \\ & \delta^8 g_{Wud}^h - \frac{c_{\theta_W} (\delta^8 g_{Zu_l}^h - \delta^8 g_{Zd_l}^h)}{\sqrt{2}} - (4\delta^8 g_{ud}^W - \sqrt{2} g c_{\theta_W}^2 \delta^8 \kappa_Z) \\ & \delta^8 g_{W\nu e}^h - \frac{c_{\theta_W} (\delta^8 g_{Z\nu_l}^h - \delta^8 g_{Ze_l}^h)}{\sqrt{2}} - (4\delta^8 g_{\nu e}^W - \sqrt{2} g c_{\theta_W}^2 \delta^8 \kappa_Z) \end{aligned}$$

$$\begin{aligned} & \delta^8 g_{Q2}^{WW} - \delta^8 g_{Q1}^{WW} - 2c_{\theta_W}^4 (\delta^8 g_{Q2}^{ZZ} - \delta^8 g_{Q1}^{ZZ}) \\ & h_Q^{ZZ} + c_{\theta_W}^4 (\delta^8 g_{Q2}^{ZZ} - \delta^8 g_{Q1}^{ZZ}) \\ & g_{hh2}^Z - 4(\delta^8 g_{Q1}^{WW} - 2c_{\theta_W}^2 \delta^8 \kappa_Z) \\ & g_{hh3}^Z + 4(\delta^8 g_{Q1}^{WW} - 2c_{\theta_W}^2 \delta^8 \kappa_Z + c_{\theta_W}^4 (\delta^8 g_{Q2}^{ZZ} - \delta^8 g_{Q1}^{ZZ})) \\ & g_{hh2}^W - 4c_{\theta_W}^4 \delta^8 g_{Q1}^{ZZ} \\ & g_{hh3}^W + 4c_{\theta_W}^4 \delta^8 g_{Q2}^{ZZ} \\ & \delta^8 \kappa^{hZ} - \frac{1}{3} \left( \frac{9\delta^8 g_{VV}^h/v - \delta^8 g_{ZZ}^{h^2}}{g^2} + 3\delta^8 g_1^{hZ} - 3t_{\theta_W}^2 (2\delta^8 g_{Q1}^{WW} + \delta^8 \kappa_{WW}^h + g^{\partial hZ}) \right. \\ & \quad \left. + 6\delta^8 \kappa_Z + s_{\theta_W}^2 (32\delta^8 \kappa_Z + 15\delta^8 \kappa_{ZZ} + 6\delta^8 g_{ZZ}^{Q1} c_{\theta_W}^2) \right) \\ & \delta^8 \kappa^{h\gamma} + \frac{1}{3s_{\theta_W}^2} \left( \frac{9\delta^8 g_{VV}^h/v - \delta^8 g_{ZZ}^{h^2}}{g^2} c_{\theta_W}^2 + 3\delta^8 g_1^{hZ} - 3s_{\theta_W}^2 (2\delta^8 g_{Q1}^{WW} + \delta^8 \kappa_{WW}^h + g^{\partial hZ}) \right. \\ & \quad \left. - 6\delta^8 \kappa_Z c_{\theta_W}^4 + s_{\theta_W}^2 c_{\theta_W}^2 (26\delta^8 \kappa_Z + 15\delta^8 \kappa_{ZZ} + 6\delta^8 g_{ZZ}^{Q1} c_{\theta_W}^2) \right) \end{aligned}$$



# PROBING SMEFT VS TESTING SMEFT





# PROBING SMEFT

Only 17 of these 52  
anomalous couplings  
need to be measured

All other anomalous couplings  
can be **predicted**  
as a function of these 17

List of Anomalous Couplings
<b>Vff couplings (9)</b>
$\Delta\mathcal{L}_{Vff} = \sum_f \delta g_f^Z Z_\mu \bar{f} \gamma^\mu f_{L,R} + \delta g_{e\nu}^W (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.) + \delta g_{ud}^W (W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c.)$
<b>Anomalous TGC (4)</b>
$\Delta\mathcal{L}_{TGC} = igc_{\theta_W} [\delta g_1^Z Z_\mu (W_\nu^+ \mathcal{W}^{-\mu\nu} - W_\nu^- \mathcal{W}^{+\mu\nu}) + \delta \kappa^Z W_\mu^+ W_\nu^- Z^{\mu\nu}]$ $+ ie \delta \kappa^\gamma W_\mu^+ W_\nu^- A^{\mu\nu} + g_5 \epsilon^{\mu\nu\rho\sigma} W_\mu^+ \overleftrightarrow{D}_\rho W_\nu^- Z_\sigma$
<b>Anomalous QGC (5)</b>
$\Delta\mathcal{L}_{QGC} = g^2 c_{\theta_W}^2 [\delta g_{ZZ1}^Q Z^\mu Z^\nu W_\mu^- W_\nu^+ - \delta g_{ZZ2}^Q Z^\mu Z_\mu W^{-\nu} W_\nu^+] + \frac{g^2}{4c_{\theta_W}^4} h_{ZZ}^Q (Z^\mu Z_\mu)^2$ $+ \frac{g^2}{2} [\delta g_{WW1}^Q W^{-\mu} W^{+\nu} W_\mu^- W_\nu^+ - \delta g_{WW2}^Q (W^{-\mu} W_\mu^+)^2]$
<b>Single Higgs (19)</b>
$\Delta\mathcal{L}_h = \delta g_{VV}^h h [W^{+\mu} W_\mu^- + \frac{1}{2c_{\theta_W}^2} Z^\mu Z_\mu] + \delta g_{ff}^h (h \bar{f}_L f_R + h.c.) + \delta g_{ZZ}^h h \frac{Z^\mu Z_\mu}{2c_{\theta_W}^2}$ $\sum_f g_{Zff}^h \frac{h}{v} Z_\mu \bar{f} \gamma^\mu f + g_{WQ}^h \frac{h}{v} (W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c.) + g_{WL}^h \frac{h}{v} (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.)$ $+ \kappa_{ZZ}^h \frac{h}{2v} Z^{\mu\nu} Z_{\mu\nu} + \kappa_{\gamma\gamma}^h \frac{h}{2v} A^{\mu\nu} A_{\mu\nu} + \kappa_{Z\gamma}^h \frac{h}{v} A^{\mu\nu} Z_{\mu\nu} + \kappa_{WW}^h \frac{h}{v} \mathcal{W}^{+\mu\nu} \mathcal{W}_{\mu\nu}^-$ $+ \kappa_{GG}^h \frac{h}{2v} G^{A\mu\nu} G_{\mu\nu}^A$
<b>hV<sup>3</sup> couplings (5)</b>
$\Delta\mathcal{L}^{hV^3} = igc_{\theta_W} \frac{h}{v} [g_{Z1}^{hV^3} Z_\mu (W_\nu^+ \mathcal{W}^{-\mu\nu} - W_\nu^- \mathcal{W}^{+\mu\nu}) + \kappa_Z^{hV^3} W_\mu^+ W_\nu^- Z^{\mu\nu}]$ $+ ie \kappa_\gamma^{hV^3} \frac{h}{v} W_\mu^+ W_\nu^- A^{\mu\nu} + g_5^{hV^3} \frac{h}{v} \epsilon^{\mu\nu\rho\sigma} W_\mu^+ \overleftrightarrow{D}_\rho W_\nu^- Z_\sigma$ $+ ig_{\partial h Z}^{hV^3} \frac{g}{2c_{\theta_W}} \frac{\partial_\mu h}{v} Z_\nu (W_\mu^+ W^{-\nu} - W_\mu^- W^{+\nu})$
<b>h<sup>2</sup>V<sup>2</sup> couplings (8)</b>
$\Delta\mathcal{L}^{hh} = \delta g_{VV}^{hh} \frac{h^2}{2} [W^{+\mu} W_\mu^- + \frac{1}{2c_{\theta_W}^2} Z^\mu Z_\mu] + \delta g_{ZZ}^{hh} \frac{h^2}{2} \frac{Z^\mu Z_\mu}{2c_{\theta_W}^2}$ $+ g_{Z1}^{hh} \frac{(\partial_\nu h)^2}{v^2} + g_{Z2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} \frac{Z^\mu Z^\nu}{c_{\theta_W}^2}$ $+ g_{W1}^{hh} \frac{(\partial_\nu h)^2}{v^2} W^{+\mu} W_\mu^- + g_{W2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} (W^{+\mu} W^{-\nu} + h.c.)$ $+ \kappa_{WW}^{hh} \frac{h^2}{2v^2} \mathcal{W}^{+\mu\nu} \mathcal{W}_{\mu\nu}^- + \kappa_{ZZ}^{hh} \frac{h^2}{4v^2} Z^{\mu\nu} Z_{\mu\nu}$
<b>h<sup>2</sup>G<sup>2</sup> couplings (1)</b>
$\Delta\mathcal{L}_{GG}^{hh} = \kappa_{GG}^{hh} \frac{h^2}{4v^2} G^{A\mu\nu} G_{\mu\nu}^A$
<b>Higgs potential corrections (2)</b>
$\Delta\mathcal{L}^{h^n} = -\delta\lambda_3 v h^3 - \delta\lambda_4 \frac{h^4}{4}$



# TESTING SMEFT

1. Beyond D6 SMEFT
2. SMEFT at D8,D10.. level/HEFT
3. Testing SMEFT assumptions

All these 52 anomalous couplings need to be probed

List of Anomalous Couplings
<i>Vff</i> couplings (9)
$\Delta\mathcal{L}_{Vff} = \sum_f \delta g_f^Z Z_\mu \bar{f} \gamma^\mu f_{L,R} + \delta g_{e\nu}^W (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.) + \delta g_{ud}^W (W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c.)$
Anomalous TGC (4)
$\Delta\mathcal{L}_{TGC} = igc_{\theta_W} [\delta g_1^Z Z_\mu (W_\nu^+ W^{-\mu\nu} - W_\nu^- W^{+\mu\nu}) + \delta \kappa^Z W_\mu^+ W_\nu^- Z^{\mu\nu}]$ $+ ie \delta \kappa^\gamma W_\mu^+ W_\nu^- A^{\mu\nu} + g_5 \epsilon^{\mu\nu\rho\sigma} W_\mu^+ \overleftrightarrow{D}_\rho W_\nu^- Z_\sigma$
Anomalous QGC (5)
$\Delta\mathcal{L}_{QGC} = g^2 c_{\theta_W}^2 [\delta g_{ZZ1}^Q Z^\mu Z^\nu W_\mu^- W_\nu^+ - \delta g_{ZZ2}^Q Z^\mu Z_\mu W^{-\nu} W_\nu^+] + \frac{g^2}{4c_{\theta_W}^4} h_{ZZ}^Q (Z^\mu Z_\mu)^2$ $+ \frac{g^2}{2} [\delta g_{WW1}^Q W^{-\mu} W^{+\nu} W_\mu^- W_\nu^+ - \delta g_{WW2}^Q (W^{-\mu} W_\mu^+)^2]$
Single Higgs (19)
$\Delta\mathcal{L}_h = \delta g_{VV}^h h [W^{+\mu} W_\mu^- + \frac{1}{2c_{\theta_W}^2} Z^\mu Z_\mu] + \delta g_{ff}^h (h \bar{f}_L f_R + h.c.) + \delta g_{ZZ}^h h \frac{Z^\mu Z_\mu}{2c_{\theta_W}^2}$ $\sum_f g_{Zff}^h \frac{h}{v} Z_\mu \bar{f} \gamma^\mu f + g_{WQ}^h \frac{h}{v} (W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c.) + g_{WL}^h \frac{h}{v} (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.)$ $+ \kappa_{ZZ}^h \frac{h}{2v} Z^{\mu\nu} Z_{\mu\nu} + \kappa_{\gamma\gamma}^h \frac{h}{2v} A^{\mu\nu} A_{\mu\nu} + \kappa_{Z\gamma}^h \frac{h}{v} A^{\mu\nu} Z_{\mu\nu} + \kappa_{WW}^h \frac{h}{v} W^{+\mu\nu} W_{\mu\nu}^-$ $+ \kappa_{GG}^h \frac{h}{2v} G^{A\mu\nu} G_{\mu\nu}^A$
<i>hV<sup>3</sup></i> couplings (5)
$\Delta\mathcal{L}^{hV^3} = igc_{\theta_W} \frac{h}{v} [g_{Z1}^{hV^3} Z_\mu (W_\nu^+ W^{-\mu\nu} - W_\nu^- W^{+\mu\nu}) + \kappa_Z^{hV^3} W_\mu^+ W_\nu^- Z^{\mu\nu}]$ $+ ie \kappa_\gamma^{hV^3} \frac{h}{v} W_\mu^+ W_\nu^- A^{\mu\nu} + g_5^{hV^3} \frac{h}{v} \epsilon^{\mu\nu\rho\sigma} W_\mu^+ \overleftrightarrow{D}_\rho W_\nu^- Z_\sigma$ $+ ig_{\partial h Z}^{hV^3} \frac{g}{2c_{\theta_W}} \frac{\partial_\mu h}{v} Z_\nu (W_\mu^+ W^{-\nu} - W_\mu^- W^{+\nu})$
<i>h<sup>2</sup>V<sup>2</sup></i> couplings (8)
$\Delta\mathcal{L}_{V^2}^{hh} = \delta g_{VV}^{hh} \frac{h^2}{2} [W^{+\mu} W_\mu^- + \frac{1}{2c_{\theta_W}^2} Z^\mu Z_\mu] + \delta g_{ZZ}^{hh} \frac{h^2}{2} \frac{Z^\mu Z_\mu}{2c_{\theta_W}^2}$ $+ g_{Z1}^{hh} \frac{(\partial_\nu h)^2}{v^2} + g_{Z2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} \frac{Z^\mu Z^\nu}{c_{\theta_W}^2}$ $+ g_{W1}^{hh} \frac{(\partial_\nu h)^2}{v^2} W^{+\mu} W_\mu^- + g_{W2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} (W^{+\mu} W^{-\nu} + h.c.)$ $+ \kappa_{WW}^{hh} \frac{h^2}{2v^2} W^{+\mu\nu} W_{\mu\nu}^- + \kappa_{ZZ}^{hh} \frac{h^2}{4v^2} Z^{\mu\nu} Z_{\mu\nu}$
<i>h<sup>2</sup>G<sup>2</sup></i> couplings (1)
$\Delta\mathcal{L}_{GG}^{hh} = \kappa_{GG}^{hh} \frac{h^2}{4v^2} G^{A\mu\nu} G_{\mu\nu}^A$
Higgs potential corrections (2)
$\Delta\mathcal{L}^{h^n} = -\delta\lambda_3 v h^3 - \delta\lambda_4 \frac{h^4}{4}$



# TESTING SMEFT

1. Beyond
2. SMEFT
3. Testing

- Thus to go from probing to testing SMEFT many more measurements are required.
- This motivates the development of sophisticated differential observables, wrt energies, angles/Multivariate distributions/Machine learning
- High energies/high luminosities required for such studies

All these  $h^2$  anomalous couplings need to be probed

List of Anomalous Couplings
$Vff$ couplings (9)
$\Delta\mathcal{L}_{Vff} = \sum_f \delta g_f^Z Z_\mu \bar{f} \gamma^\mu f_{L,R} + \delta g_{e\nu}^W (W_\mu^+ \bar{\nu}_L \gamma^\mu e_L + h.c.) + \delta g_{ud}^W (W_\mu^+ \bar{u}_L \gamma^\mu d_L + h.c.)$
Anomalous TGC (4)
$\Delta\mathcal{L}_{TGC} = igc_{\theta_W} [\delta g_1^Z Z_\mu (W_\nu^+ W^{-\mu\nu} - W_\nu^- W^{+\mu\nu}) + \delta \kappa^Z W_\mu^+ W_\nu^- Z^{\mu\nu}]$ $+ i\epsilon \delta \kappa^\gamma W_\mu^+ W_\nu^- A^{\mu\nu} + g_5 \epsilon^{\mu\nu\rho\sigma} W_\mu^+ \overleftrightarrow{D}_\rho W_\nu^- Z_\sigma$

$\Delta\mathcal{L}_{V^2}^{hh} = \delta g_{VV}^{hh} \frac{h^2}{2} \left[ W^{+\mu} W_\mu^- + \frac{1}{2c_{\theta_W}^2} Z^\mu Z_\mu \right] + \delta g_{ZZ}^{hh} \frac{h^2}{2} \frac{Z^\mu Z_\mu}{2c_{\theta_W}^2}$ $+ g_{Z1}^{hh} \frac{(\partial_\nu h)^2}{v^2} + g_{Z2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} \frac{Z^\mu Z^\nu}{c_{\theta_W}^2}$ $+ g_{W1}^{hh} \frac{(\partial_\nu h)^2}{v^2} W^{+\mu} W_\mu^- + g_{W2}^{hh} \frac{\partial_\mu h \partial_\nu h}{2v^2} (W^{+\mu} W^{-\nu} + h.c.)$ $+ \kappa_{WW}^{hh} \frac{h^2}{2v^2} W^{+\mu\nu} W_{\mu\nu}^- + \kappa_{ZZ}^{hh} \frac{h^2}{4v^2} Z^{\mu\nu} Z_{\mu\nu}$
$h^2 G^2$ couplings (1)
$\Delta\mathcal{L}_{GG}^{hh} = \kappa_{GG}^{hh} \frac{h^2}{4v^2} G^{A\mu\nu} G_{\mu\nu}^A$
Higgs potential corrections (2)
$\Delta\mathcal{L}^{h^n} = -\delta\lambda_3 v h^3 - \delta\lambda_4 \frac{h^4}{4}$



---

# BEYOND SMEFT 2: AMPLITUDES

---

- Basic idea: one can try to find **most general Lorentz invariant parameterisation of an amplitude** for a process.
  - There is a **mapping between EFT Wilson coefficients and the parameters** determining the amplitudes.
  - **No of parameters must equal no of Wilson coefficients.**
  - **Amplitudes much more physical. No redundancies** in amplitude parametrisation unlike Wilson coefficients.
-



---


# BEYOND SMEFT 2: AMPLITUDES

---

- Many recent papers with similar objectives:

1. Shadmi & Weiss (2018)
2. Durieux, Kitahara, Shadmi & Weiss (2019)
3. Durieux, Kitahara, Machado, Shadmi & Weiss (2020)
- 4.
5. Ma, Shu & Xiao (2019)
6. Baratella, Fernandez & Pomarol (2020)
- 7.
8. Jiang, Ma & Shu (2020)
9. Dong, Ma, Shu & Zhou (2022)
10. Chang, Chen, Liu and Luty (2022)

*We will focus on this recent work  
in this talk*





# AMPLITUDES EXAMPLE: HIGGSTRAHLUNG

As an example take the **most general amplitude for Higgstrahlung**:

$$\mathcal{M}(f_1 \bar{f}_2 \rightarrow Z_3 h_4) = \bar{u}_2 \Gamma^\mu u_1 \epsilon_{3\mu}^*$$

$$\begin{aligned} \Gamma^\mu = & c_1 p_1^\mu + c_2 p_2^\mu + c_3 p_1^\mu \gamma_5 + c_4 p_2^\mu \gamma_5 + c_5 \gamma^\mu + c_6 p_1^\mu \not{p}_3 + c_7 p_2^\mu \not{p}_3 \\ & + c_8 \gamma^\mu \gamma_5 + c_9 p_1^\mu \not{p}_3 \gamma_5 + c_{10} p_2^\mu \not{p}_3 \gamma_5 + c_{11} \gamma^{\mu\nu} p_{3\nu} \\ & + \epsilon^{\mu\nu\rho\sigma} p_{1\nu} p_{2\rho} p_{3\sigma} (c_{12} + c_{13} \gamma_5 + c_{14} \not{p}_3) + \epsilon^{\mu\nu\rho\sigma} \gamma_\nu (c_{15} p_{1\rho} p_{2\sigma} + c_{16} p_{1\rho} p_{3\sigma} + c_{17} p_{2\rho} p_{3\sigma}) \\ & + \epsilon^{\mu\nu\rho\sigma} p_{1\nu} p_{2\rho} p_{3\sigma} (c_{18} \not{p}_3 \gamma_5) + \epsilon^{\mu\nu\rho\sigma} \gamma_\nu \gamma_5 (c_{19} p_{1\rho} p_{2\sigma} + c_{20} p_{1\rho} p_{3\sigma} + c_{21} p_{2\rho} p_{3\sigma}) \\ & + c_{22} \epsilon_{\nu\rho\sigma\gamma} \gamma^{\mu\nu} p_1^\rho p_2^\sigma p_3^\gamma + \epsilon^{\mu\nu\rho\sigma} \gamma_\nu \gamma p_3^\gamma (c_{23} p_{1\rho} p_{2\sigma} + c_{24} p_{1\rho} p_{3\sigma} + c_{25} p_{2\rho} p_{3\sigma}) \\ & + \epsilon^{\mu\nu\rho\sigma} \gamma_\nu \rho (c_{26} p_{1\sigma} + c_{27} p_{2\sigma} + c_{28} p_{3\sigma}) \\ & + \epsilon^{\alpha\beta\gamma\delta} \gamma_\alpha p_{1\beta} p_{2\gamma} p_{3\delta} (c_{29} p_1^\mu + c_{30} p_2^\mu + c_{31} p_1^\mu \gamma_5 + c_{32} p_2^\mu \gamma_5). \end{aligned}$$

where  $c_n = f(p_i \cdot p_j)$



# AMPLITUDES EXAMPLE: HIGGSTRAHLUNG

$$\mathcal{M}(f_1 \bar{f}_2 \rightarrow Z_3 h_4) = \frac{1}{v} (\bar{u}_2 \not{\epsilon}_3^* u_1) \left[ \boxed{A} + \boxed{B} \frac{s}{M^2} + C \frac{t}{M^2} + \dots \right] \\ + \frac{1}{v^3} (\bar{u}_2 \not{p}_4 u_1) (p_4 \cdot \epsilon_3^*) \left[ \boxed{A'} + B' \frac{s}{M^2} + C' \frac{t}{M^2} + \dots \right] + \dots$$

Primary:  $h Z_\mu \bar{f} \gamma^\mu f$

Descendant:  $\partial_\rho h \partial^\rho Z_\mu \bar{f} \gamma^\mu f$

Primary:  $ih \tilde{Z}_{\mu\nu} \bar{f} \gamma^\mu \overleftrightarrow{\partial}_\nu f$

- Most general amplitude can be rewritten in above form where there are **primaries** in the amplitude with ‘**Mandelstam descendant**’ contributions ( $B, C, B', C'$  etc) **suppressed by powers of  $s/\Lambda^2, t/\Lambda^2$**  etc
- Each term** in the above expansion corresponds to **an anomalous coupling** (HEFT operator). **Higher order terms are couplings/operators with more derivatives.**



# AMPLITUDES EXAMPLE: HIGGSTRAHLUNG

- While there are an infinite number of independent parameters/anomalous couplings there are **only a finite number of primaries**. These are **all independent**.

- Chang et al list all primary operators (up to arbitrary high dimension) for the important Higgs production and decay processes:

$$\begin{aligned}
 (\bar{f}f, gg, W^+W^-, ZZ) &\rightarrow (h, hh, hZ, h\gamma, hg) \\
 (\bar{f}f', ZW) &\rightarrow hW, \\
 (fg, f\gamma, fZ) &\rightarrow hf, \\
 fW &\rightarrow f'h.
 \end{aligned}$$

- These **can be distinguished in angular measurements**. Measuring these can become a **target for experiments**.

Ex: 12 primaries for Higgstrahlung:

$i$	$\mathcal{O}_i^{hZ\bar{f}f}$
1	$hZ^\mu\bar{\psi}_L\gamma_\mu\psi_L$
2	$hZ^\mu\bar{\psi}_R\gamma_\mu\psi_R$
3	$hZ^{\mu\nu}\bar{\psi}_L\sigma_{\mu\nu}\psi_R + \text{h.c.}$
4	$ih\tilde{Z}_{\mu\nu}\bar{\psi}_L\sigma^{\mu\nu}\psi_R + \text{h.c.}$
5	$ihZ^\mu(\bar{\psi}_L\overset{\leftrightarrow}{\partial}_\mu\psi_R) + \text{h.c.}$
6	$hZ^\mu\partial_\mu(\bar{\psi}_L\psi_R) + \text{h.c.}$
7	$ihZ^\mu\partial_\mu(\bar{\psi}_L\psi_R) + \text{h.c.}$
8	$hZ^\mu(\bar{\psi}_L\overset{\leftrightarrow}{\partial}_\mu\psi_R) + \text{h.c.}$
9	$ih\tilde{Z}_{\mu\nu}(\bar{\psi}_L\gamma^\mu\overset{\leftrightarrow}{\partial}^\nu\psi_L)$
10	$h\tilde{Z}_{\mu\nu}\partial^\mu(\bar{\psi}_L\gamma^\nu\psi_L)$
11	$ih\tilde{Z}_{\mu\nu}(\bar{\psi}_R\gamma^\mu\overset{\leftrightarrow}{\partial}^\nu\psi_R)$
12	$h\tilde{Z}_{\mu\nu}\partial^\mu(\bar{\psi}_R\gamma^\nu\psi_R)$



# AMPLITUDES EXAMPLE: HIGGSTRAHLUNG

- While there are an infinite number of independent parameters/anomalous couplings there are **only a finite number of primaries**. These are **all independent**.

- Chang et al list all primary operators (up to arbitrary high dimension) for the important Higgs production and

This again motivates **new angular observables to pinpoint** these different **anomalous couplings**

$$\begin{aligned} (\bar{f}f', ZW) &\rightarrow hW, \\ (fg, f\gamma, fZ) &\rightarrow hf, \\ fW &\rightarrow f'h. \end{aligned}$$

- These **can be distinguished in angular measurements**. Measuring these can become a **target for experiments**.

Ex: 12 primaries for Higgstrahlung:

$i$	$\mathcal{O}_i^{hZ\bar{f}f}$
1	$hZ^\mu\bar{\psi}_L\gamma_\mu\psi_L$
2	$hZ^\mu\bar{\psi}_R\gamma_\mu\psi_R$
3	$hZ^{\mu\nu}\bar{\psi}_L\sigma_{\mu\nu}\psi_R + \text{h.c.}$
4	$hZ^\mu\bar{\psi}_L\gamma_\mu\psi_R + \text{h.c.}$
5	$hZ^\mu\bar{\psi}_R\gamma_\mu\psi_L + \text{h.c.}$
6	$hZ^\mu\bar{\psi}_L\gamma_\mu\psi_L + \text{h.c.}$
7	$ihZ^\mu\partial_\mu(\bar{\psi}_L\psi_R) + \text{h.c.}$
8	$hZ^\mu(\bar{\psi}_L\overleftrightarrow{\partial}_\mu\psi_R) + \text{h.c.}$
9	$ih\tilde{Z}_{\mu\nu}(\bar{\psi}_L\gamma^\mu\overleftrightarrow{\partial}^\nu\psi_L)$
10	$h\tilde{Z}_{\mu\nu}\partial^\mu(\bar{\psi}_L\gamma^\nu\psi_L)$
11	$ih\tilde{Z}_{\mu\nu}(\bar{\psi}_R\gamma^\mu\overleftrightarrow{\partial}^\nu\psi_R)$
12	$h\tilde{Z}_{\mu\nu}\partial^\mu(\bar{\psi}_R\gamma^\nu\psi_R)$



---


# DIFFERENTIAL STUDIES

---

- To distinguish all these anomalous couplings, we have to move towards **maximally differential studies**
  - Many new studies that try to identify **differential signatures that can pinpoint operators/anomalous couplings**
  - These include **Multivariate methods** to probe such finer effects
1. Franceschini, Pomarol, Panico, Riva & Wulzer (2017)
  2. Panico, Riva and Wulzer (2017)
  3. Azatov, Elias-Miro, Reyimuaji & Venturini (2017)
  4. Englert, Banerjee, RSG & Spannowsky (2018)
  5. Banerjee, RSG, Reines & Spannowsky (2019)
  6. Banerjee, RSG, Reines, Seth & Spannowsky (2019)
  7. Rahaman & Singh (2019)
  8. Chen, Giloti, Panico & Wulzer (2020)
  9. Rao, Rindani & Sarmah (2019, 2021)
-



# HIGGS ANOMALOUS COUPLINGS IN ZH PROD.




All these anomalous couplings can be completely predicted in terms of other more precise measurements, *if we assume D6 SMEFT.*

3  $hZZ$  anomalous couplings



# HIGGS ANOMALOUS COUPLINGS IN ZH PROD.




All these anomalous couplings must be measured, *if we want to test D6 SMEFT*

3  $hZZ$  anomalous couplings




# HIGGS ANOMALOUS COUPLINGS IN ZH PROD.

$$hZ_{\mu}Z^{\mu}$$




Rescales SM  $hZZ$  coupling.  
No differential signature.  
Only changes the rate.

$$hZ_{\mu}\partial_{\nu}Z^{\mu\nu}$$



$$hZ_{\mu\nu}Z^{\mu\nu}, hZ_{\mu\nu}\tilde{Z}^{\mu\nu}$$




3  $hZZ$  anomalous couplings




# HIGGS ANOMALOUS COUPLINGS IN ZH PROD.

$$hZ_\mu Z^\mu$$




$$hZ_\mu \partial_\nu Z^{\mu\nu}$$



Grows with energy wrt SM.  
Dominates at high energies.


$$hZ_{\mu\nu} Z^{\mu\nu}, hZ_{\mu\nu} \tilde{Z}^{\mu\nu}$$



3  $hZZ$  anomalous couplings



# ENERGY GROWING EFFECTS




- We studied  $Z(l\bar{l})H(bb)$  at high energies using **boosted Higgs reconstruction** techniques to obtain **per-mille level bounds** on  $hVff$  couplings that are competitive with LEP:

$$|g_{Zp}^h| < 5 \times 10^{-4}.$$



# HIGGS ANOMALOUS COUPLINGS IN ZH PROD.



Sophisticated angular  
variable required

3  $hZZ$  anomalous couplings

Banerjee, RSG, Reines & Spannowsky (2019)


Banerjee, RSG, Reines, Seth & Spannowsky (2019)



# DIFFERENTIAL INFORMATION IN

$$pp \rightarrow Z(l\bar{l})H(\text{fat jet})$$

- How much differential information in this process?
- Three body phase space so  $3 \times 3 - 4 = 5$  kinematical variables completely define the final state




- Ignoring the boost there are 4:

$$\sqrt{s}, \quad \Theta, \quad \theta, \quad \varphi$$



# A TRIPLE DIFFERENTIAL OBSERVABLE



$$\kappa_{ZZ} \frac{h}{2v} Z^{\mu\nu} Z_{\mu\nu}$$

$$g^5 = C_\varphi S_\Theta S_\theta C_\Theta C_\theta$$

Vanishes when  
integrated over any of  
the 3 angles

Events weighted by sign of  $S_\Theta S_\theta C_\Theta C_\theta$

*Interference resurrection!*


$$\tilde{\kappa}_{ZZ} \frac{h}{2v} Z^{\mu\nu} \tilde{Z}_{\mu\nu}$$

$$g^7 = S_\varphi S_\Theta S_\theta C_\Theta C_\theta$$

Vanishes when  
integrated over any of  
the 3 angles



# A TRIPLE DIFFERENTIAL OBSERVABLE



Banerjee, RSG, Reines & Spannowsky (2019)

Banerjee, RSG, Reines, Seth & Spannowsky (2019)



---

# CONCLUSIONS

---

- SMEFT not the most general EFT for LHC studies. Amplitudes/HEFT provide a more general framework.
  - Many viable UV models, map to HEFT not SMEFT.
  - SMEFT vs HEFT: In SMEFT different linear combinations of anomalous couplings suppressed wrt HEFT by powers of  $v^2/\Lambda^2$ .
  - The 'Amplitudes' approach has identified a set of primary operators that give leading contribution to amplitudes in the EFT derivative expansion.
  - Both approaches require new differential observables that can pinpoint these effects. Many require multivariate studies, high luminosities and high energies.
-



# FULL ANGULAR INFORMATION FOR HIGGSTRAHLUNG

$ff \rightarrow Z(l)h$  matrix element squared

- These 9 coefficients carry full differential information in SM and D6 SMEFT
- Can be extracted using an analog of Fourier analysis called the 'Method of Moments'

$$\begin{aligned} \sum_{L,R} |\mathcal{A}(\hat{s}, \Theta, \theta, \varphi)|^2 = & a_{LL} \sin^2 \Theta \sin^2 \theta + a_{TT}^1 \cos \Theta \cos \theta \\ & + a_{TT}^2 (1 + \cos^2 \Theta)(1 + \cos^2 \theta) + \cos \varphi \sin \Theta \sin \theta \\ & \times (a_{LT}^1 + a_{LT}^2 \cos \theta \cos \Theta) + \sin \varphi \sin \Theta \sin \theta \\ & \times (\tilde{a}_{LT}^1 + \tilde{a}_{LT}^2 \cos \theta \cos \Theta) + a_{TT'} \cos 2\varphi \sin^2 \Theta \sin^2 \theta \\ & + \tilde{a}_{TT'} \sin 2\varphi \sin^2 \Theta \sin^2 \theta. \end{aligned}$$


Consider *these 2 functions*

*Vanish when integrated  
over any of the 3 angles*



# WW, WZ, W $\gamma$ PRODUCTION

$$\begin{aligned} \mathcal{L}_{\text{WWV}}/g_{\text{WWV}} = & ig_1^V (W_{\mu\nu}^\dagger W^\mu V^\nu - W_\mu^\dagger V_\nu W^{\mu\nu}) + i\kappa_V W_\mu^\dagger W_\nu V^{\mu\nu} \\ & + \frac{i\lambda_V}{m_W^2} W_{\lambda\mu}^\dagger W^\mu_\nu V^{\nu\lambda} - g_4^V W_\mu^\dagger W_\nu (\partial^\mu V^\nu + \partial^\nu V^\mu) \\ & + g_5^V \epsilon^{\mu\nu\rho\sigma} (W_\mu^\dagger \vec{\partial}_\rho W_\nu) V_\sigma + i\tilde{\kappa}_V W_\mu^\dagger W_\nu \tilde{V}^{\mu\nu} \\ & + \frac{i\tilde{\lambda}_V}{m_W^2} W_{\lambda\mu}^\dagger W^\mu_\nu \tilde{V}^{\nu\lambda}. \end{aligned}$$



*s-channel contribution*

anomalous triple gauge vertices  
( $\mathbb{I} \rightarrow 6$  CP even + 5 CP odd)

How many of these  $\mathbb{I}$  can we measure if we use all the  
energy/ angular information ?

# EG: PREDICTIONS IN CP EVEN CASE

- 3 D6 SMEFT operators:

$$ig(D^\mu H)^\dagger \sigma^a (D^\nu H) W_{\mu\nu}^a$$

$$\frac{1}{3!} g \epsilon_{abc} W_\mu^a W_{\nu\rho}^b W^{\rho\mu c}$$

$$\frac{ig}{2} \left( H^\dagger \sigma^a \overleftrightarrow{D}^\mu H \right) D^\nu W_{\mu\nu}^a$$

- 6 CP even anomalous couplings. If we measure all of these in differential studies we can verify SMEFT predictions below.

- 3 D6 SMEFT predictions:  $\delta\kappa_Z = \delta g_1^Z - t_{\theta_W}^2 \delta\kappa_\gamma$   $\lambda_Z = \lambda_\gamma$   $g_5 = 0$



# SMEFT OPERATORS

$$\begin{aligned}
 \mathcal{O}_{H\Box} &= |H|^2 \Box |H|^2 \\
 \mathcal{O}_{HD} &= (H^\dagger D_\mu H)^* (H^\dagger D^\mu H) \\
 \mathcal{O}_6 &= |H|^6 \\
 \mathcal{O}_y &= \hat{y}_f |H|^2 \bar{F} H f_R \\
 \mathcal{O}_f &= i H^\dagger \overleftrightarrow{D}_\mu H \bar{f} \gamma^\mu f \\
 \mathcal{O}_F^{(3)} &= i H^\dagger \sigma^a \overleftrightarrow{D}_\mu H \bar{F} \sigma^a \gamma^\mu F \\
 \mathcal{O}_{BB} &= g'^2 |H|^2 B_{\mu\nu} B^{\mu\nu} \\
 \mathcal{O}_{WB} &= gg' H^\dagger \sigma^a H W_{\mu\nu}^a B^{\mu\nu} \\
 \mathcal{O}_{WW} &= g^2 |H|^2 W_{\mu\nu}^a W^{a\mu\nu} \\
 \mathcal{O}_{GG} &= g_s^2 |H|^2 G_{\mu\nu}^A G^{A\mu\nu}
 \end{aligned}$$

Dimension 6

$H^8$
$\mathcal{O}_8 =  H ^8$
$H^6 D^2: 2 \quad 2$
$\mathcal{O}_{H^2 r} =  H ^4  D_\mu H ^2$
$\mathcal{O}_{H^2 T} = \frac{ H ^2}{2} (H^\dagger \overleftrightarrow{D}_\nu H)^2$
$H^4 X^2: 3 \quad 4$
$\mathcal{O}_{H^2 BB} = g'^2  H ^4 B_{\mu\nu} B^{\mu\nu}$
$\mathcal{O}_{H^2 WB} =  H ^2 \mathcal{O}_{WB}$
$\mathcal{O}_{H^2 WW} = g^2  H ^4 W_{\mu\nu}^a W^{a\mu\nu}$
$\mathcal{O}_U = g^2 (H^\dagger \sigma^a H) (H^\dagger \sigma^b H) W_{\mu\nu}^a W^{b\mu\nu}$
$\mathcal{O}_{H^2 GG} = g_s^2  H ^4 G_{\mu\nu}^A G^{A\mu\nu}$
$H^4 D\psi^2: 9 \quad 9$
$\mathcal{O}_{H^2 f} = i  H ^2 H^\dagger \overleftrightarrow{D}_\mu H \bar{f} \gamma^\mu f$
$\mathcal{O}_{H^2 F} = i  H ^2 H^\dagger \overleftrightarrow{D}_\mu H \bar{F} \gamma^\mu F$
$\mathcal{O}_{H^2 F}^{(3)} = i  H ^2 H^\dagger \sigma^a \overleftrightarrow{D}_\mu H \bar{F} \sigma^a \gamma^\mu F$
$\mathcal{O}_{3F}^{(3)} = i H^\dagger \overleftrightarrow{D}_\mu H (H^\dagger \sigma^a H) \bar{F} \sigma^a \gamma^\mu F$
$H^4 D^2 X: 3 \quad 3$
$\mathcal{O}_{HWH} = ig W_{\mu\nu}^a (H^\dagger \sigma^a D_\mu H + h.c.) H^\dagger \overleftrightarrow{D}_\nu H$
$\mathcal{O}_{\partial W} = ig W_{\mu\nu}^a \partial_\mu (H^\dagger H) H^\dagger \sigma^a \overleftrightarrow{D}_\nu H$
$\mathcal{O}_{\partial B} = ig' B_{\mu\nu} \partial_\mu (H^\dagger H) H^\dagger \overleftrightarrow{D}_\nu H$
$H^4 D^4: 3 \quad 3$
$\mathcal{O}_{DH1} =  D_\mu H ^4$
$\mathcal{O}_{DH2} = (D_\mu H^\dagger D_\nu H + D_\nu H^\dagger D_\mu H)^2$
$\mathcal{O}_{DH3} = (D_\mu H^\dagger D_\nu H - D_\nu H^\dagger D_\mu H)^2$

Dimension 8



# PROBING D6 SMEFT

- Only 17 D6 operators contribute to the processes we are considering
- Only 17 measurements sufficient to constrain these

$$\begin{aligned}\mathcal{O}_{H\Box} &= |H|^2 \Box |H|^2 \\ \mathcal{O}_{HD} &= (H^\dagger D_\mu H)^* (H^\dagger D^\mu H) \\ \mathcal{O}_6 &= |H|^6 \\ \mathcal{O}_y &= \hat{y}_f |H|^2 \bar{F} H f_R \\ \mathcal{O}_f &= i H^\dagger \overleftrightarrow{D}_\mu H \bar{f} \gamma^\mu f \\ \mathcal{O}_F^{(3)} &= i H^\dagger \sigma^a \overleftrightarrow{D}_\mu H \bar{F} \sigma^a \gamma^\mu F \\ \mathcal{O}_{BB} &= g'^2 |H|^2 B_{\mu\nu} B^{\mu\nu} \\ \mathcal{O}_{WB} &= g g' H^\dagger \sigma^a H W_{\mu\nu}^a B^{\mu\nu} \\ \mathcal{O}_{WW} &= g^2 |H|^2 W_{\mu\nu}^a W^{a\mu\nu} \\ \mathcal{O}_{GG} &= g_s^2 |H|^2 G_{\mu\nu}^A G^{A\mu\nu}\end{aligned}$$