

Complementarity of μ TRISTAN and Belle II in searches for CLFV.

Gabriela Lima Lichtenstein, in collaboration with
Michael Schmidt, German Valencia, Raymond Volkas



Lepton Photon 2023
July 17, 2023

Charged Lepton Flavour Violation

Charged Lepton Flavour Violation (LFV) was not observed so far. However, neutrino oscillations show that there are no individual lepton numbers L_e , L_μ , and L_τ conservation.

Several BSM models predict cLFV.

Experimental bounds are stringent, especially on electron-muon cLFV.

$$\mu \rightarrow e\gamma,$$

$$\mu \rightarrow eee.$$

cLFV involving τ :

Data are less constraining; Belle II future sensitivity will increase significantly.

Lepton Triality

Avoids $\mu \leftrightarrow e$ conversions
Allows LFV tau decays.

Ma,(2010) 1006.3524

Motivated by flavour structure models

Tribimaximal mixing of neutrino flavours **Altarelli, Feruglio (2006)**

Residual Z_3 symmetry may emerge from breaking of A_4 flavour symmetry Residual Z_3

He, Keum, Volkas (2006)
[hep-ph/0601001](https://arxiv.org/abs/hep-ph/0601001)

Lepton Triality

$L \rightarrow \omega^T L$ and $e_R \rightarrow \omega^T e_R$,
 $\omega = e^{\frac{2\pi i}{3}}$

H, quarks are singlets under triality

$$\mathcal{L}_Y = y_{e_i} L_i e_{R_i} H + h.c.$$

\mathcal{L}_Y is diagonal under Z_3

The leptons e, μ , τ are charged under flavour triality
 $T = 1, 2, 3$

• Triality sums modulo 3

$$\mu^- \rightarrow e^- \gamma \quad \Delta T=0 \times$$

$T=2 \quad T=1$

$$\tau^- \rightarrow \mu^+ e^- e^- \quad \Delta T=0 \checkmark$$

$T=3 \quad T=-2 \quad T=1 \quad T=1$

Simple extensions to the SM using scalar bileptons can mediate these triality-preserving interactions


Lepton Triality

Models T = 1, 2, 3 for the doubly charged singlet k_i

$$\mathcal{L}_{k_1} = \frac{1}{2} \left(2f_1 \overline{(\tau_R)^c} \mu_R + f_2 \overline{(e_R)^c} e_R \right) k_1 + \text{h.c.}$$

$$\mathcal{L}_{k_2} = \frac{1}{2} \left(2g_1 \overline{(\tau_R)^c} e_R + g_2 \overline{(\mu_R)^c} \mu_R \right) k_2 + \text{h.c.}$$

$$\mathcal{L}_{k_3} = \frac{1}{2} \left(2h_1 \overline{(\mu_R)^c} e_R + h_2 \overline{(\tau_R)^c} \tau_R \right) k_3 + \text{h.c..}$$




Tau Decays

Tau LFV decays:

$$\tau \rightarrow e/\mu + \gamma,$$

$$\tau \rightarrow e/\mu + l^+l^- \text{ where } l = e/\mu.$$



Observable	Present constraint	Projected sensitivity
$\text{BR}(\tau^- \rightarrow \mu^- \mu^- e^+)$	$1.7 \times 10^{-8} *$	$2.6 \times 10^{-10} **$
$\text{BR}(\tau^- \rightarrow \mu^+ e^- e^-)$	$1.5 \times 10^{-8} *$	$2.3 \times 10^{-10} **$

* Belle Collaboration (2010) 1001.3221

** Belle II (2022) 2203.14919

$$\mathcal{L}_{6,LFV} = C^{\prime\prime}(\bar{L}\gamma_\mu L)(\bar{L}\gamma^\mu L) + C^{ee}(\bar{e}_R\gamma_\mu e_R)(\bar{e}_R\gamma^\mu e_R) + C^{le}(\bar{L}\gamma_\mu L)(\bar{e}_R\gamma^\mu e_R)$$

$$C_{ee,1312}^{VRR} = \frac{f_1 f_2}{4m_{k_1}^2}$$

$$BR(\tau^\pm \rightarrow \mu^\mp e^\pm e^\pm) = \\ \frac{f_1^2 f_2^2}{64 G_F^2 m_{k_1}^4} BR(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)$$

$$C_{ee,2321}^{VRR} = \frac{g_1 g_2}{4m_{k_2}^2}$$

$$BR(\tau^\pm \rightarrow \mu^\pm \mu^\pm e^\mp) = \\ \frac{g_1^2 g_2^2}{64 G_F^2 m_{k_2}^4} \tilde{J}\left(\frac{m_\mu^2}{m_\tau^2}\right) BR(\tau^- \rightarrow \mu^- \bar{\nu}_\mu \nu_\tau)$$

Present bounds:

$$\sqrt{f_1 \times f_2} < 0.17 \frac{m_{k_1}}{\text{TeV}}$$



$$\sqrt{g_1 \times g_2} < 0.17 \frac{m_{k_2}}{\text{TeV}}$$

Prediction for future sensitivity:

$$\sqrt{f_1 \times f_2} < 0.06 \frac{m_{k_1}}{\text{TeV}}$$

$$\sqrt{g_1 \times g_2} < 0.06 \frac{m_{k_2}}{\text{TeV}}$$

Belle II sensitivity on cLFV tau decays from Triality T=1



Bigaran, He, Schmidt, Valencia, Volkas, (2022) 2212.09760.

μ TRISTAN

Hamada, Kitano, Matsudo, Takaura and Yoshida, (2022) 2201.06664

Ultracold muon technology from g-2 at J-PARC

$\mu^+ \mu^+$ proposal $\sqrt{s} = 2$ TeV;

1 TeV μ^+ beams;

expected luminosity of 12 fb^{-1} per year.

$\mu^+ e^-$ proposal with asymmetric beam energies;

μ^+ beams up to 1 to 3 TeV;



e^- beams from Tristan at 30 to 50 GeV;

expected luminosity of 100 fb^{-1} per year.

Future lepton Colliders

Model	Process	Lepton Collider
T=1	$\mu^+ e^- \rightarrow e^+ \tau^-$	μ TRISTAN
T=1	$e^+ e^- \rightarrow e^+ e^-$	$e^+ e^-$
T=1	$e^- e^- \rightarrow e^- e^-$	-
T=1	$e^- e^- \rightarrow \tau^- \mu^-$	-
T=2	$\mu^+ \mu^+ \rightarrow \tau^+ e^+$	μ TRISTAN
T=2	$\mu^+ \mu^+ \rightarrow \mu^+ \mu^+$	μ TRISTAN
T=2	$\mu^+ e^- \rightarrow \tau^+ \mu^-$	μ TRISTAN
T=2	$\mu^+ \mu^- \rightarrow \mu^+ \mu^-$	$\mu^+ \mu^-$
T=3	$\mu^+ e^- \rightarrow \mu^+ e^-$	μ TRISTAN
T=3	$\mu^+ e^+ \rightarrow \tau^+ \tau^+$	-

CLFV s-channel at $\mu^+\mu^+$




90% C.L. contour
assuming no background
 $N = 2.44;$

$$\sqrt{g_1 g_2} \lesssim 0.15 \left(\frac{N}{L_s} \right)^{\frac{1}{4}} \frac{m_{k_2}}{\text{TeV}}$$

For $\sqrt{s} = 2$ TeV:


$$\sqrt{g_1 g_2} \lesssim 0.17 \frac{m_{k_2}}{\text{TeV}} .$$

CLFV u-channel at $\mu^+ e^-$





$E_e = 30 \text{ GeV}, G_\mu = 1 \text{ TeV}$

$E_e = 50 \text{ GeV}, E_\mu = 3 \text{ TeV}$




$$\sqrt{f_1 f_2} \lesssim 0.13 \frac{m_{k_1}}{\text{TeV}}$$

Resonances in elastic scattering $\mu^+\mu^+ \rightarrow \mu^+\mu^+$



Resonances in elastic scattering $\mu^+\mu^+ \rightarrow \mu^+\mu^+$




90% C.L. contour;
SM contributions as
background;
 $S = 1.64$

$$S = \frac{|\sigma - \sigma_{SM}|}{\sqrt{\sigma_{SM}}} \sqrt{L};$$

$$g_2 \lesssim 0.18 \left(\frac{S^2}{Ls} \right)^{1/4} m_k;$$
$$g_2 \lesssim 0.09 \frac{m_{k_2}}{\text{TeV}}.$$

Elastic scattering $\mu^+ e^- \rightarrow \mu^+ e^-$



$$E_c = 30 \text{ GeV}, \quad E_\mu = 1 \text{ TeV}$$

$$E_c = 50 \text{ GeV}, \quad E_\mu = 3 \text{ TeV}$$

$$S = \frac{|\sigma - \sigma_{SM}|}{\sqrt{\sigma_{SM}}} \sqrt{L};$$

$$h_1 \lesssim 0.17 \frac{m_{k3}}{\text{TeV}}.$$

Summary

Lepton Flavour Triality avoids CLFV bounds from muon decays while allowing tau LFV interactions;

Belle II sensitivity to tau LFV processes will increase significantly.

$$\text{Belle} \quad L = 782 \text{ fb}^{-1} \quad \sqrt{f_1 f_2} < 0.17 \frac{m_\tau}{\text{TeV}}, \quad \sqrt{g_1 g_2} < 0.17 \frac{m_\tau}{\text{TeV}}$$

$$\text{Belle II} \quad L = 50 \text{ ab}^{-1} \quad \sqrt{f_1 f_2} < 0.06 \frac{m_\tau}{\text{TeV}}, \quad \sqrt{g_1 g_2} < 0.06 \frac{m_\tau}{\text{TeV}}$$

μ TRISTAN $\mu^+ \mu^+$ collider

Resonances searches

$$\begin{aligned} \mu^+ \mu^- &\rightarrow \tau^+ e^- & \sqrt{g_1 g_2} &< 0.01 \frac{m_\tau}{\text{TeV}} \\ \mu^+ \mu^- &\rightarrow \mu^+ \mu^- & g_2 &< 0.09 \frac{m_\tau}{\text{TeV}} \end{aligned} \quad \left. \right\} \begin{array}{l} L = 12 \text{ fb}^{-1} \\ \sqrt{s} = 2 \text{ TeV} \end{array}$$

μ TRISTAN $\mu^+ e^-$ collider

$$\begin{aligned} \mu^+ e^- &\rightarrow e^+ \tau^- & \sqrt{f_1 f_2} &< 0.13 \frac{m_\tau}{\text{TeV}} \\ \mu^+ e^- &\rightarrow \mu^- \tau^+ & \sqrt{g_1 g_2} &< 0.13 \frac{m_\tau}{\text{TeV}} \\ \mu^+ e^- &\rightarrow \mu^+ e^- & h_1 &< 0.17 \frac{m_\tau}{\text{TeV}} \end{aligned} \quad \left. \right\} \begin{array}{l} L = 100 \text{ fb}^{-1} \\ \sqrt{s} = 0.346 \text{ GeV} \end{array}$$