

Capture of Dark Matter in Neutron Stars

[2004.14888] + [2010.13257] + [2012.08918] + [2104.14367] +
[2108.02525]

G. Busoni

The Australian National University

Lepton Photon 2023, Melbourne, 18/07/23

Outline

- 1 Why Compact Stars?
 - Searching for DM collisions: Direct Detection
 - Searching for DM collisions: DM Capture in the Sun
 - Searching for DM collisions: DM Capture in Neutron Stars
 - Advantages of NS
 - Detection prospects
- 2 From capture in the Sun to NS and WD
 - Key differences
- 3 Results
 - Leptons: highlights and results
 - Baryons: highlights and results
 - Summary

Outline

1

Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars
- Advantages of NS
- Detection prospects

2

From capture in the Sun to NS and WD

- Key differences

3



Results

- Leptons: highlights and results
- Baryons: highlights and results
- Summary

Searching for DM collisions

Direct Detection

Direct Detection



SI interaction give much stronger bounds than SD ones

Searching for DM collisions

Direct Detection

- Constraints depend strongly on interaction type
- Strong target dependence
- Some operators are suppressed by kinematics (momentum/velocity suppressed)
- Recoil energy is small, nonrelativistic kinematics
- Experimental detectors have recoil energy thresholds

Outline

1 Why Compact Stars?

- Searching for DM collisions: Direct Detection
- **Searching for DM collisions: DM Capture in the Sun**
- Searching for DM collisions: DM Capture in Neutron Stars
- Advantages of NS
- Detection prospects

2 From capture in the Sun to NS and WD

- Key differences

3 Results


- Leptons: highlights and results
- Baryons: highlights and results
- Summary

Searching for DM collisions

DM Capture in the Sun

DM Capture in Stars


- DM can be captured and accumulate in Stars
- Dark matter scatters, loses energy, becomes gravitationally bound to star
- Accumulates in centre of Sun
- Can potentially annihilate at the center
- At equilibrium
Capture=Annihilation
- Probes same observables as DD



Searching for DM collisions

DM Capture in the Sun

- SI: DD wins
- SD: DM in Sun wins
- DM in Sun requires some few more assumptions, like that it annihilates, and the annihilation channel



Some other ways to infer indirectly DM presence in the Sun: modified energy transport (see 1411.6626 , 1703.07784)

Outline

1 Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars**
- Advantages of NS
- Detection prospects

2 From capture in the Sun to NS and WD

- Key differences

3 Results

- Leptons: highlights and results
- Baryons: highlights and results
- Summary

Searching for DM collisions

DM Capture in NS

- Very large density means very efficient capture
- Whole DM flux can be captured already for $\sigma \sim 10^{-45} cm^2$



Searching for DM collisions

DM Capture in NS

Possible observable signals

- NS to BH collapse (more likely for bosonic DM)
- Gravitational waves: DM increases tidal deformability
(1803.03266)
- Kinetic Heating (M. Baryakhtar et al. PRL 119, 131801 (2017)
arXiv:1704.01577)
- Kinetic + Annihilation heating (Bramante, Delgado and Martin
1704.01577)

Searching for DM collisions

DM Capture in NS


NS temperature evolution

- NS have no known large heating sources
- Lose energy by neutrino and photon emission
- Neutrino dominates early stages of NS life, photon the late stages
- In absence of other heating sources, one expects $T \sim 1000K$ after $10Myr$ and $T \sim 100K$ after $1Gyr$
- Kinetic heating: sets a maximum equilibrium temperature of $T_{eq,th} \sim 1700K$ if whole DM flux is captured
- Kin+Ann heating: maximum equilibrium temperature is raised to $T_{eq,th} \sim 2400K$

Searching for DM collisions

DM Capture in NS

NS temperature evolution



Searching for DM collisions

DM Capture in NS

Equilibrium Temperature

$$C_{geom} = \frac{\pi R^2 (1 - B(R))}{v_s B(R)} \frac{\rho_\chi}{m_\chi} \text{Erf} \left(\sqrt{\frac{3}{2}} \frac{v_s}{v_d} \right) \quad (1)$$

$$C \langle \delta E_R \rangle = \sigma_{SB} T_{eq}^4 \pi R^2 \quad (2)$$

T_{eq} independent of star radius for geometric limit, $T_{eq}^4 \propto \rho_\chi$. Can define threshold cross section:

$$C = C_{geom} \frac{\sigma}{\sigma_{th}}, \quad T_{eq} \propto \left(\frac{\sigma}{\sigma_{th}} \right)^{1/4} \quad (3)$$

Outline

1 Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars
- **Advantages of NS**
- Detection prospects

2 From capture in the Sun to NS and WD

- Key differences

3 Results

- Leptons: highlights and results
- Baryons: highlights and results
- Summary

Searching for DM collisions

Advantages of NS

- High capture probability
- DM particles accelerated to $\mathcal{O}(0.5)c$ means no momentum/velocity suppression
- Cross section of $\sigma_{th} = 10^{-45} \text{ cm}^2$ enough to reach maximum capture
- No threshold recoil energy
- Similar sensitivity for SI and SD interaction
- Similar sensitivity for momentum/velocity suppressed interactions comparing to unsuppressed ones
- Observation of old cold NS of temperature $T < T_{eq,th}$:


$$\sigma \leq \sigma_{th} \left(\frac{T}{T_{eq,th}} \right)^4 \quad (4)$$

Searching for DM collisions

Advantages of NS

Example: Inelastic DM

- The high incoming energy allows to probe larger mass splittings
- For large mass, $\delta < 330\text{MeV}$
- For small mass, $\delta < 330\text{MeV} \frac{m_\chi}{\text{GeV}}$
- This is 3-4 orders of magnitude more than for Sun or Earth
- Absence of energy threshold give further advantage comparing to DD




Searching for DM collisions

Advantages of NS

Example:
Momentum-suppressed
Operators

- The high incoming energy makes all types of operators giving similar results at large energies
- This is opposed to DD, where there is a hierarchy due to non-relativistic suppression



Outline

1 Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars
- Advantages of NS
- Detection prospects

2 From capture in the Sun to NS and WD

- Key differences

3 Results


- Leptons: highlights and results
- Baryons: highlights and results
- Summary

Searching for DM collisions

Detection prospects


James Webb Telescope

- Launched 25 December 2021
- Situated at L2
- Primary mirror: $6.5m$
- Operating wavelengths: $0.6 - 28.5\mu m$



Searching for DM collisions


Detection prospects



Searching for DM collisions

Detection prospects

James Webb and Hubble compared




Source: Nasa

BBC

Searching for DM collisions

Detection prospects

Telescopes cover different parts of the electromagnetic spectrum




Searching for DM collisions

Detection prospects

Detection prospects

- Warning: rough estimates
- Cold NS very faint: need large observation time
- Need $t_{obs} \gtrsim 10^5 s \sim 1d$
- $t_{obs} \sim \mathcal{O}(10^6 s), \mathcal{O}(10^7 s)$ should allow detection of some NS of $T \gtrsim 1000 K$
- Coldest NS: PSR J2144–3933, $T < 42000 K$



Outline

1 Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars
- Advantages of NS
- Detection prospects

2 From capture in the Sun to NS and WD

- Key differences

3 Results

- Leptons: highlights and results
- Baryons: highlights and results
- Summary

From capture in the Sun to NS and WD

Key differences

- DM Capture in the Sun formalism developed by Gould in the '80
- We adapted this formalism to NS

Sun/Gould+Extensions	NS/Our
Newtonian gravity	GR
Sun structure from Standard Solar Model	NS structure from EOS
Non-relativistic kinematics	Relativistic kinematics
Atomic Nuclei Targets	Baryon and Lepton targets
Non-relativistic matrix element	Relativistic matrix element
MB distribution for targets	FD distribution for targets
Capture probability $\neq 1$	Capture probability = 1*
Star opacity	Star Opacity
MS requires MC approach	MS can be treated analytically
Targets have FF	Targets have FF
Fixed Target mass	Density-dependent Target mass

* for some masses

Outline

1

Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars
- Advantages of NS
- Detection prospects

2

From capture in the Sun to NS and WD

- Key differences

3


Results

- Leptons: highlights and results
- Baryons: highlights and results
- Summary

From capture in the Sun to NS and WD


Leptons: highlights and results

- Very degenerate and relativistic target due to their low mass
- Relativistic treatment is very important for these targets






From capture in the Sun to NS and WD

Leptons: highlights and results



From capture in the Sun to NS and WD

Leptons: highlights and results



Outline

1

Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars
- Advantages of NS
- Detection prospects

2

From capture in the Sun to NS and WD

- Key differences

3


Results

- Leptons: highlights and results
- **Baryons: highlights and results**
- Summary

From capture in the Sun to NS and WD


Baryons: highlights and results

- Baryons are composite particles
- Strong force mean field effects require treatment beyond free Fermi gas




From capture in the Sun to NS and WD

Baryons: highlights and results






From capture in the Sun to NS and WD

Baryons: highlights and results



From capture in the Sun to NS and WD

Baryons: highlights and results



Outline

1

Why Compact Stars?

- Searching for DM collisions: Direct Detection
- Searching for DM collisions: DM Capture in the Sun
- Searching for DM collisions: DM Capture in Neutron Stars
- Advantages of NS
- Detection prospects

2

From capture in the Sun to NS and WD

- Key differences

3

Results

- Leptons: highlights and results
- Baryons: highlights and results
- Summary

Conclusions

Summary

- Neutron Stars: cosmic laboratory to probe DM scattering interactions
- Completely different kinematic regime to direct detection experiments
- High energy scattering washes away momentum suppression
- Higher reach on inelastic scattering
- Can probe a very large mass range
- Very sensitive for all interactions, including momentum-suppressed and leptons
- Current coolest NS of $\mathcal{O}(10^4)K$
- Prospects for observation in the coming decade