



Contribution ID: 72

Type: **Talk**

【347】 Model-independent measurement of charm-mixing parameters in $D^0 \rightarrow K_S^0 \pi^+ \pi^-$

Wednesday 29 June 2022 18:30 (15 minutes)

We present a measurement of charm mixing and CP -violation parameters using $D^0 \rightarrow K_S^0 \pi^+ \pi^-$ decays reconstructed in pp collisions collected by the LHCb experiment from years 2016 to 2018. In particular, the analysis measures the dimensionless parameter x related to the mass difference between the mass eigenstates of the D^0 meson. This was observed to be non-zero with a significance exceeding seven standard deviations in $D^{*+} \rightarrow (D^0 \rightarrow K_S^0 \pi^+ \pi^-) \pi^+$ decay [PRL 127, 111801 (2021)]. We present here an independent measurement where D^0 candidates are reconstructed from $B^- \rightarrow D^0 \mu^- X$ decays as the sample covers wider D^0 decay time distribution. A combination of the two analyses is performed to maximise the sensitivity.

Author: EK-IN, Surapat (EPFL - Ecole Polytechnique Federale Lausanne (CH))

Presenter: EK-IN, Surapat (EPFL - Ecole Polytechnique Federale Lausanne (CH))

Session Classification: Nuclear, Particle- & Astrophysics

Track Classification: Nuclear, Particle- and Astrophysics (TASK)