

Flavor-Changing Light Bosons with Accidental Longevity

Kun-Feng Lyu

School of Physics and Astronomy, University of Minnesota

In collaboration with Yohei Ema, Zhen Liu and Maxim Pospelov

2211.XXXX

Outline

- Motivation
- Model
- Muon Magnetic Moment
- e-mu case
- mu-tau case

Motivation

- The latest muon $g-2$ measurement shows 3.7 sigma deviation from SM prediction.
- There have been many scenarios to explain.
- We propose a simple scalar flavor off-diagonally coupled to lepton. A new mass window is probed.
- In the mass window, it becomes accidentally long-lived and thus can be probed at the beamdump experiment.

Model

- Consider a complex scalar ϕ carrying +1 mu number and -1 l number

$$\mathcal{L} = |\partial\phi|^2 - m_\phi^2 |\phi|^2 + \phi \bar{\mu} (g_V + g_A \gamma_5) l + \phi^* \bar{l} (g_V^* - g_A^* \gamma_5) \mu,$$

- In the mass range, it can become long-lived accidentally


$$|m_\mu - m_l| < m_\phi < m_\mu + m_l$$

$$m_\phi < m_\mu + m_l$$

ϕ can not decay to $\mu + l$

$$|m_\mu - m_l| < m_\phi$$


Avoid lepton decay branching ratio.



Muon Magnetic Moment

- The latest measurement


$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (251 \pm 59) \times 10^{-11} \quad 2104.03281$$



$$a_\mu^{(\phi)} = \frac{m_\mu}{8\pi^2} \int_0^1 dz \frac{(1-z)^2 \left[(|g_V|^2 + |g_A|^2) z m_\mu + (|g_V|^2 - |g_A|^2) m_l \right]}{-z(1-z)m_\mu^2 + (1-z)m_l^2 + z m_\phi^2}$$

e-mu case

Muon g-2 constraint



Shaded: electron g-2 exclusion region

At this region, the decay length is around 10^{10} km

- Muonium
the quasi-bound state of mu and e, can decay to phi plus a photon.


$$\text{Br}(\text{Mu}^{(S=1)} \rightarrow \text{invisible}) < 5.7 \times 10^{-6}$$

1209.0060


LSND Experiment

- We need intense muon source.
- Originally to measure mu-neutrino to e-neutrino oscillation.
- Mu^+ can be produced by the meson π^+ , it can form muonium when traveling through the water.
- muonium decay $\rightarrow \text{phi}$

Phi decay in the LSND detector.




hep-ex/0104049




The parameter space that can explain muon $g-2$ is excluded by the LSND!

mu-tau case


- Phi is heavier than that in e-mu case.
- We can probe it in the higher energy beamdump experiments.
- There are mainly 3 channels for the production.



“Direct EW”




“ D_s decay”




“ μ on target”

Experiment	E_{beam} (GeV)	POT	D (m)	L (m)	A (m^2)	$\epsilon_{\text{acpt}}^{\text{geo}}$	Major production
CHARM	400	2×10^{18}	480	35	3×3	1.3%	EW, D_s , μOT
NuTeV	800	μOT	850	34	2.54×2.54	$\mathcal{O}(1)\%$	μOT
SHiP	400	2×10^{20}	60	60	5×10	54%	EW, D_s , μOT



The corresponding decay length is around 10 km.

Can be detected in the far detector.



- CHARM and NuTeV cannot exclude the parameter region that explains the muon $g - 2$ anomaly.
- While the future SHiP experiment covers the whole parameter space.

Conclusion

- We study the phenomenology of a complex scalar coupled to mu and either e or tau motivated by muon $g-2$.
- The scalar becomes long-lived in the mass window we considered.
- For e-mu case, phi can be produced by the muonium that is formed in LSND proton beamdump experiment. The excess events shows the the $g-2$ allowed region is excluded.
- For mu-tau case, phi can be produced by the Drell-Yan process, heavy meson decay and muon-target. The future SHiP can cover most parameter space.

Thank you!