



# Analytic two-loop soft and beam functions for leading-jet $p_T$

Samuel Abreu

CERN & The University of Edinburgh

---


Based on 2204.02987 and 2205.xxxxx

Together with J. Gaunt, P. Monni, R. Szafron + L. Rottoli

SCET 2022 — 21st of April 2022, Bern

# Motivation


- ♦ Jet-vetoed cross sections: reduce backgrounds in Higgs analyses
  - ✓ e.g.  $H \rightarrow WW$  vs  $t\bar{t}$



Formalism for resummation

[Banfi, Monni, Salam, Zanderighi <sup>12</sup>;  
 Becher, Neubert <sup>12</sup>;  
 Becher, Neubert, Rothen <sup>13</sup>;  
 Stewart, Tackmann, Walsh, Zuberi <sup>13</sup>]

## Higgs jet-vetoed cross section



State of the art:  $N^3\text{LO+NNLL}$  [Banfi et al <sup>16</sup>]

Goal: improve  
resummation to  $N^3\text{LL}$

# Cross-section factorisation

- ♦ Colour-singlet production with jet veto  $p_T^{\text{jet}} < p_T^{\text{veto}}$

$$\frac{d\sigma(p_T^{\text{veto}})}{d\Phi_{\text{Born}}} \quad \xrightarrow{\hspace{1cm}} \text{logarithms } \log(p_T^{\text{veto}}/Q)$$

- ♦ In the limit  $p_T^{\text{veto}} \ll Q$

[Becher, Neubert 12;  
Becher, Neubert, Rothen 13;  
Stewart, Tackmann, Walsh, Zuberi 13]

$$\frac{d\sigma(p_T^{\text{veto}})}{d\Phi_{\text{Born}}} = |A_{\text{Born}}^F|^2 \mathcal{H}(Q; \mu) \mathcal{B}_n(x_1, Q, p_T^{\text{veto}}, R^2; \mu, \nu) \mathcal{B}_{\bar{n}}(x_2, Q, p_T^{\text{veto}}, R^2; \mu, \nu) \mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu)$$

- ✓ Hard function:  $\mathcal{H}(Q; \mu)$
- ✓ Beam functions:  $\mathcal{B}_{n,\bar{n}}(x, Q, p_T^{\text{veto}}, R^2; \mu, \nu)$
- ✓ Soft function:  $\mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu)$

# Evolution equations

- ♦ Hard function  $\mathcal{H}(Q; \mu) = |\mathcal{C}(Q; \mu)|^2$

[Becher, Neubert, Rothen 13;  
Stewart, Tackmann, Walsh, Zuberi 13]

$$\frac{d}{d \ln \mu} \ln \mathcal{C}(Q; \mu) = \Gamma_{\text{cusp}}(\alpha_s(\mu)) \ln \frac{-Q^2}{\mu^2} + \gamma_H(\alpha_s(\mu))$$

- ♦ Soft function

$$\frac{d}{d \ln \mu} \ln \mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu) = 4 \Gamma_{\text{cusp}}(\alpha_s(\mu)) \ln \frac{\mu}{\nu} + \gamma_S(\alpha_s(\mu))$$

$$\frac{d}{d \ln \nu} \ln \mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu) = -4 \int_{p_T^{\text{veto}}}^{\mu} \frac{d\mu'}{\mu'} \Gamma_{\text{cusp}}(\alpha_s(\mu')) + \gamma_\nu(p_T^{\text{veto}}, R^2)$$

- ♦ Beam functions

$$\frac{d}{d \ln \mu} \ln \mathcal{B}_n(x, Q, p_T^{\text{veto}}, R^2; \mu, \nu) = 2 \Gamma_{\text{cusp}}(\alpha_s(\mu)) \ln \frac{\nu}{Q} + \gamma_B(\alpha_s(\mu))$$

$$\frac{d}{d \ln \nu} \ln \mathcal{B}_n(x, Q, p_T^{\text{veto}}, R^2; \mu, \nu) = 2 \int_{p_T^{\text{veto}}}^{\mu} \frac{d\mu'}{\mu'} \Gamma_{\text{cusp}}(\alpha_s(\mu')) - \frac{1}{2} \gamma_\nu(p_T^{\text{veto}}, R^2)$$

Missing for N<sup>3</sup>LL: two-loop  $\mathcal{S}$  and  $\mathcal{B}_n$ , three-loop  $\gamma_\nu$

# **SOFT FUNCTION**

## **CALCULATION AND RESULTS**

---

# The soft function

- ♦ Operator definition:

$$\mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu) = \frac{1}{d_F} \sum_{X_s} \text{Tr} \left\{ \mathcal{M}(p_T^{\text{veto}}, R^2) \langle 0 | Y_n^\dagger Y_{\bar{n}} | X_s \rangle \langle X_s | Y_{\bar{n}}^\dagger Y_n | 0 \rangle \right\}$$

- ✓ Soft Wilson lines:  $Y_{n,\bar{n}}$
- ✓ Measurement function:  $\mathcal{M}(p_T^{\text{veto}}, R^2)$

$$\mathcal{M}(p_T^{\text{veto}}, R^2) = \Theta(p_T^{\text{veto}} - \max\{p_T^{\text{jet}_i}\}) \Theta_{\text{cluster}}(R^2)$$

- ♦ Regularisation of divergences:

- ✓ UV/IR/Coll. divergences: dimensional regularisation
- ✓ Rapidity divergences: exponential regulator

- ♦ Exponential regulator: modify phase-space integration measure


$$\prod_i d^d k_i \delta(k_i^2) \theta(k_i^0) \rightarrow \prod_i d^d k_i \delta(k_i^2) \theta(k_i^0) \exp \left[ \frac{-e^{-\gamma_E}}{\nu} (n \cdot k_i + \bar{n} \cdot k_i) \right]$$

# Soft function decomposition

- ♦ Reference observable:


$$\mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu) = \mathcal{S}_{\perp}(p_T^{\text{veto}}, \mu, \nu) + \Delta\mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu)$$

[e. g. : Banfi, Salam, Zanderighi 12  
 Gangal, Gaunt, Stahlhofen, Tackmann 16  
 Bauer, Manohar, Monni 20]



- ✓  $\mathcal{S}_{\perp}(p_T^{\text{veto}}, \mu, \nu)$ : soft function for  $p_T$  resummation


[with the same regulator:  
 Li, Neill, Zhu 16; Li, Zhu 16]



- ✓  $\Delta\mathcal{S}(p_T^{\text{veto}}, R^2; \mu, \nu)$ : remainder

$$\Delta\mathcal{M}(p_T^{\text{veto}}, R^2) \equiv \Theta(p_T^{\text{veto}} - \max\{p_T^{\text{jet}_i}\})\Theta_{\text{cluster}}(R^2) - \Theta\left(p_T^{\text{veto}} - \left|\sum_{X_s} p_T^{\text{jet}_i}\right|\right)$$

- ✓ Remainder: only rapidity divergences, work in four dimensions



- ✓  $\Delta\mathcal{S}^{(2)}(p_T^{\text{veto}}, R^2; \mu, \nu)$ : double-real diagrams with two soft gluons or a soft quark-antiquark pair

[Campbell, Glover 98  
 Dokshitzer, Lucenti, Marchesini, Salam 97  
 Catani, Grazzini 99]

- ♦ Two-loop correlated and uncorrelated contributions

$$\Delta\mathcal{S}^{(2)}(p_T, R^2; \mu, \nu) = \Delta S^{\text{corr.}}(p_T, R^2; \mu, \nu) + \Delta S^{\text{uncorr.}}(p_T, R^2; \mu, \nu)$$

- ✓  $\Delta S^{\text{uncorr.}}(p_T, R^2; \mu, \nu)$ : two emissions are widely separated in rapidity
- ✓  $\Delta S^{\text{corr.}}(p_T, R^2; \mu, \nu)$ :  $\rightarrow 0$  when two emissions are widely separated

# Calculation: setup

- ♦ Momenta parametrisation:

$$k_i = k_{i\perp} (\cosh \eta_i, \cos \phi_i, \sin \phi_i, \sinh \eta_i), \quad i = 1, 2$$

$$\{k_{2\perp}, \eta_2, \phi_2\} \rightarrow \{\zeta \equiv k_{2\perp}/k_{1\perp}, \eta \equiv \eta_1 - \eta_2, \phi \equiv \phi_1 - \phi_2\}$$

- ♦ Squared amplitudes:

$$\mathcal{A}^{\text{cor./uncor.}}(k_1, k_2) = \frac{1}{k_{1\perp}^4} \frac{1}{\zeta^2} \mathcal{D}^{\text{cor./uncor.}}(\zeta, \eta, \phi)$$

- ♦ Integrals to compute:

$$\int \frac{dk_{1\perp}}{k_{1\perp}} d\eta_1 \frac{d\zeta}{\zeta} d\eta \frac{d\phi}{2\pi} e^{-2k_{1\perp} \frac{e^{-\gamma_E}}{\nu} [\cosh(\eta_1) + \zeta \cosh(\eta - \eta_1)]} \mathcal{D}(\zeta, \eta, \phi) \Delta\mathcal{M}(p_T^{\text{veto}}, R^2)$$

- ♦ Measurement function, after some manipulation

$$\Delta\mathcal{M}(p_T^{\text{veto}}, R^2) \equiv \left[ \Theta(p_T^{\text{veto}} - k_{1\perp} \max\{1, \zeta\}) - \Theta\left(p_T^{\text{veto}} - k_{1\perp} \sqrt{1 + \zeta^2 + 2\zeta \cos \phi}\right) \right] \Theta(\eta^2 + \phi^2 - R^2)$$

- ♦ Goal:  $\Delta\mathcal{S}^{(2)}(p_T, R^2; \mu, \nu)$  as a series in powers of  $R^2$

# Calculation: correlated contribution

- ◆ Rapidity divergences:

$$\mathcal{D}^{\text{cor.}}(\zeta, \eta, \phi) \rightarrow 0 \text{ for } \eta = \eta_1 - \eta_2 \rightarrow \infty \quad \Rightarrow$$

Only  $\eta_1$  integral needs rapidity regulation

- ◆ Integrate over  $\eta_1$  (easy!), keep terms that survive when  $\nu \rightarrow \infty$

$$I(p_T^{\text{veto}}/\nu, R^2) = \int \frac{dk_{1\perp}}{k_{1\perp}} \frac{d\zeta}{\zeta} d\eta \frac{d\phi}{2\pi} \Omega\left(\frac{k_{1\perp}}{\nu}, \zeta, \eta\right) \mathcal{D}^{\text{cor.}}(\zeta, \eta, \phi) \Delta\mathcal{M}(p_T^{\text{veto}}, R^2)$$

$$\Omega\left(\frac{k_{1\perp}}{\nu}, \zeta, \eta\right) = \eta + 2 \ln \frac{\nu}{k_{1\perp}} - \ln(1 + \zeta e^\eta) - \ln(\zeta + e^\eta)$$

- ◆ Measurement function:  $\Theta(\eta^2 - R^2 + \phi^2) = \underbrace{\Theta(\phi^2 - R^2)}_{\text{part A}} + \underbrace{\Theta(R^2 - \phi^2) \Theta(\eta^2 - R^2 + \phi^2)}_{\text{part B}}$

✓  $I_A(p_T^{\text{veto}}/\nu, R^2)$ : full  $R^2$  dependence, expand in powers of  $R^2$

✓  $I_B(p_T^{\text{veto}}/\nu, R^2)$ : regular at  $R^2=0$

[HypExp, Huber, Maitre 05  
PolyLogTools, Duhr, Dulat 19]

- Compute  $\frac{\partial}{\partial R^2} I_B$  order by order in  $R^2$

- Solve differential equation order by order in  $R^2$ ,  $I(p_T^{\text{veto}}/\nu, 0)$

# Calculation: uncorrelated contribution

10

- ◆ Rapidity divergences: both on  $\eta$  and  $\eta_1$ !

$$\mathcal{D}^{\text{uncor.}}(\zeta, \eta, \phi) = 16C_R^2$$

$$\int \frac{dk_{1\perp}}{k_{1\perp}} d\eta_1 \frac{d\zeta}{\zeta} d\eta \frac{d\phi}{2\pi} e^{-2k_{1\perp} \frac{e^{-\gamma_E}}{\nu} [\cosh(\eta_1) + \zeta \cosh(\eta - \eta_1)]} \Delta\mathcal{M}(p_T^{\text{veto}}, R^2)$$

- ◆ More subtle  $\eta$  and  $\eta_1$  integration of exponential regulator
  - ✓ Set  $w = e^\eta, x = e^{\eta_1}$ , take Laplace transform
  - ✓ In Laplace space, expand exponential regulator in distributions
  - ✓ Take inverse Laplace transform, keep terms that survive when  $\nu \rightarrow \infty$

$$\int \frac{dx}{x w} e^{-k_{1\perp} \frac{e^{-\gamma_E}}{\nu x} [1 + w\zeta + \frac{x^2}{w} (w + \zeta)]} \rightarrow 4\delta(w) \ln\left(\frac{k_{1\perp}}{\nu}\right) \ln\left(\frac{\zeta k_{1\perp}}{\nu}\right) + \left[\frac{1}{w}\right]_+ \ln\left(\frac{\nu^2 w}{k_{1\perp}^2 (w + \zeta)(1 + \zeta w)}\right) + \mathcal{O}\left(\frac{1}{\nu^2}\right)$$

- ◆ Continue as for the correlated contributions for remaining integrals

# Calculation: numerical calculation, full $R$ dependence

---

## ♦ Correlated contributions


- ✓ **Variables:**  $\phi, \eta, \eta_t = \frac{1}{2}(\eta_1 + \eta_2), z = \frac{k_{1\perp}^2}{k_{1\perp}^2 + k_{2\perp}^2}, \mathcal{K}_T^2 = k_{1\perp}^2 + k_{2\perp}^2$
- ✓ **Exponential regulator:** as in analytic calculation
- ✓ **Analytic integrations:**  $\eta_t, \mathcal{K}_T^2$
- ✓ **Numerical integrations:**  $\eta, \phi, z$

## ♦ Uncorrelated corrections


- ✓ **Variables:**  $\phi, \eta_1, \eta_2, z = \frac{k_{1\perp}^2}{k_{1\perp}^2 + k_{2\perp}^2}, \mathcal{K}_T^2 = k_{1\perp}^2 + k_{2\perp}^2$
- ✓ **Exponential regulator:** as in analytic calculation
- ✓ **Analytic integrations:**  $\eta_1, \eta_2, \mathcal{K}_T^2$
- ✓ **Numerical integrations:**  $\phi, z$

## ♦ Retain full $R$ dependence, per mille precision in numerical evaluation

- ♦ Analytic results for  $\Delta\mathcal{S}^{(2)}(p_T^{\text{veto}}, R^2; \mu, \nu)$  to  $\mathcal{O}(R^8)$  (also  $\mathcal{S}_{\perp}^{(2)}(p_T^{\text{veto}}, R^2; \mu, \nu)$ )  
 ✓ see also numerical evaluation in different scheme [Bell, Rahn, Talbert <sup>18,20</sup>]
- ♦ Reproduce known rapidity anomalous dimension [Banfi, Monni, Salam, Zanderighi <sup>12</sup>; Becher, Neubert, Rothen <sup>13</sup>; Stewart, Tackmann, Walsh, Zuberi <sup>13</sup>]
- ♦ Verify suitability of  $R^2$  expansion for  $0 < R < 1$



Comparison with numerics, with full  $R^2$  dependence



Check  $\mathcal{O}(R^8)$  corrections are negligible

$$\delta_{\mathcal{C}}(R) = \left| 1 - \frac{\Delta\mathcal{S}_{\mathcal{C}}^{(2)}(p_T, R^2; \mu, p_T)|_{R^6}}{\Delta\mathcal{S}_{\mathcal{C}}^{(2)}(p_T, R^2; \mu, p_T)|_{R^8}} \right|$$

# **BEAM FUNCTION**

## **CALCULATION AND RESULTS**

---

# The beam functions

[e.g., Becher, Neubert 12;  
 Luo, Wang, Xu, Yang, Yang, Zhu 19;  
 Luo, Yang, Zhu, Zhu, 19]

- ◆ Operator definition:

$$\mathcal{B}_q(x, Q, p_T^{\text{veto}}, R^2; \mu, \nu) = \frac{1}{2\pi} \sum_{X_C} dte^{-ixt\bar{n}\cdot p} \mathcal{M}(p_T^{\text{veto}}, R^2) \langle P(p) | \bar{\chi}_n(t\bar{n}) \frac{\not{t}}{2} | X_C \rangle \langle X_C | \chi_n(0) | P(p) \rangle$$

$$\mathcal{B}_g(x, Q, p_T^{\text{veto}}, R^2; \mu, \nu) = -\frac{x\bar{n} \cdot p}{2\pi} \sum_{X_C} dte^{-ixt\bar{n}\cdot p} \mathcal{M}(p_T^{\text{veto}}, R^2) \langle P(p) | \mathcal{A}_{\perp}^{\mu, a}(t\bar{n}) | X_C \rangle \langle X_C | \mathcal{A}_{\perp, \mu}^a(0) | P(p) \rangle$$

- ✓  $\chi_n, \mathcal{A}_{\perp}^{\mu, a}$ : collinear gauge invariant fields
- ✓ Measurement function: as for Soft function

- ◆ Must be perturbatively matched to PDFs
- ◆ Regularisation of divergences: as for Soft function

- ✓ Exponential regulator:

$$\prod_i d^d k_i \delta(k_i^2) \theta(k_i^0) \rightarrow \prod_i d^d k_i \delta(k_i^2) \theta(k_i^0) \exp \left[ \frac{-e^{-\gamma_E}}{\nu} (n \cdot k_i + \bar{n} \cdot k_i) \right]$$


- ♦ Reference observable:

$$\mathcal{B}(x, p_T^{\text{veto}}, R^2; \mu, \nu) = \mathcal{B}_\perp(x, p_T^{\text{veto}}, \mu, \nu) + \Delta \mathcal{B}(x, p_T^{\text{veto}}, R^2; \mu, \nu)$$

- ♦ Same approach as Soft function: decompose into different contributions
  - ✓ Several **channels** and several **colour factors**

- ♦ Structure of our results

$$\Delta \mathcal{B}^{(2)}(x, R^2) = \delta(1 - x) f_1(R^2) + \left[ \frac{1}{1 - x} \right]_+ \left( f_2(x, R^2) + f_3(x) \right)$$






# Results and checks

- ♦ Analytic results for  $\Delta\mathcal{B}^{(2)}(x, p_T^{\text{veto}}, R^2; \mu, \nu)$  to  $\mathcal{O}(R^8)$

- ♦ Reproduce known rapidity anomalous dimension

(requires 0-bin subtraction and handling of Soft-Collinear mixing)

- ♦ Verify suitability of  $R^2$  expansion for  $0 < R < 1$



Comparison with numerics, with full  $R^2$  dependence

Check  $\mathcal{O}(R^8)$  corrections are negligible

$$\delta_{\mathcal{C}}(R) = \left| 1 - \frac{\Delta\mathcal{B}_{\mathcal{C}}^{(2)}(p_T, R^2; \mu, p_T)|_{R^6}}{\Delta\mathcal{B}_{\mathcal{C}}^{(2)}(p_T, R^2; \mu, p_T)|_{R^8}} \right|$$

# CONCLUSION AND OUTLOOK

---

- ♦ Fully analytic two-loop soft function for jet-vetoed cross sections
- ♦ Analytic two-loop beam functions (up to boundary condition at  $R = 0$ )
- ♦ To do: Validate finite terms of soft/beam functions by reproducing NNLO DY and Higgs cross sections
- ♦ To do: Determine the three-loop  $\gamma_\nu$
- ♦ To do: Setup N<sup>3</sup>LL resummation of jet-vetoed cross section

# **THANK YOU!**