



Synchrotron Radiation Light Sources

Lenny Rivkin

Paul Scherrer Institute (PSI)

and

Swiss Federal Institute of Technology Lausanne (EPFL)



Paul Scherrer Institute

SwissFEL

Swiss Light Source

Instruments development: 400 years of discoveries with "telescopes" and "microscopes"



Galileo Galilei



Zacharias Janssen



The First Compound Microscope (circa 1595)

Accelerator driven applications to meet the needs of society

- Advanced instruments for basic and applied science
- Analysis of physical, chemical and biological materials
- Modification of physical, chemical and biological properties of matter
- Medical: diagnostics, treatment and targeted drug design
- Security: cargo scanning, IT hardware
- Environment
- Energy

A beam of particles is a very useful tool.


A beam of the right particles with the right energy at the right intensity can shrink a tumor, produce cleaner energy, spot suspicious cargo, make a better radial tire, clean up dirty drinking water, map a protein, study a nuclear explosion, design a new drug, make a heat-resistant automotive cable, diagnose a disease, reduce nuclear waste, detect an art forgery, implant ions in a semiconductor, prospect for oil, date an archaeological find, package a Thanksgiving turkey
or
discover the secrets of the universe

From the [Report "Accelerators for America's Future"](#), US Department of Energy, 2010

The role of accelerators in Physical and Life Sciences

“Instruments have a life of their own. They do not merely follow theory; often they determine theory, because instruments determine what is possible, and what is possible determines to a large extent what can be thought. The telescope, the microscope, the chronograph, the photograph: all gave rise to a blossoming of theoretical understanding not possible before their invention”

Hankins & Silverman,
Instruments and the Imagination



24 Nobel Prizes in Physics that had direct contribution from accelerators

Year	Name	Accelerator-Science Contribution to Nobel Prize-Winning Research
1939	Ernest O. Lawrence	Lawrence invented the cyclotron at the University of Californian at Berkeley in 1929 [12].
1951	John D. Cockcroft and Ernest T.S. Walton	Cockcroft and Walton invented their eponymous linear positive-ion accelerator at the Cavendish Laboratory in Cambridge, England, in 1932 [13].
1952	Felix Bloch	Bloch used a cyclotron at the Crocker Radiation Laboratory at the University of California at Berkeley in his discovery of the magnetic moment of the neutron in 1940 [14].
1957	Tsung-Dao Lee and Chen Ning Yang	Lee and Yang analyzed data on K mesons (θ and τ) from Bevatron experiments at the Lawrence Radiation Laboratory in 1955 [15], which supported their idea in 1956 that parity is not conserved in weak interactions [16].
1959	Emilio G. Segrè and Owen Chamberlain	Segrè and Chamberlain discovered the antiproton in 1955 using the Bevatron at the Lawrence Radiation Laboratory [17].
1960	Donald A. Glaser	Glaser tested his first experimental six-inch bubble chamber in 1955 with high-energy protons produced by the Brookhaven Cosmotron [18].
1961	Robert Hofstadter	Hofstadter carried out electron-scattering experiments on carbon-12 and oxygen-16 in 1959 using the SLAC linac and thereby made discoveries on the structure of nucleons [19].
1963	Maria Goeppert Mayer	Goeppert Mayer analyzed experiments using neutron beams produced by the University of Chicago cyclotron in 1947 to measure the nuclear binding energies of krypton and xenon [20], which led to her discoveries on high magic numbers in 1948 [21].
1967	Hans A. Bethe	Bethe analyzed nuclear reactions involving accelerated protons and other nuclei whereby he discovered in 1939 how energy is produced in stars [22].
1968	Luis W. Alvarez	Alvarez discovered a large number of resonance states using his fifteen-inch hydrogen bubble chamber and high-energy proton beams from the Bevatron at the Lawrence Radiation Laboratory [23].
1976	Burton Richter and Samuel C.C. Ting	Richter discovered the J/ψ particle in 1974 using the SPEAR collider at Stanford [24], and Ting discovered the J/ψ particle independently in 1974 using the Brookhaven Alternating Gradient Synchrotron [25].
1979	Sheldon L. Glashow, Abdus Salam, and Steven Weinberg	Glashow, Salam, and Weinberg cited experiments on the bombardment of nuclei with neutrinos at CERN in 1973 [26] as confirmation of their prediction of weak neutral currents [27].

1980	James W. Cronin and Val L. Fitch	Cronin and Fitch concluded in 1964 that CP (charge-parity) symmetry is violated in the decay of neutral K mesons based upon their experiments using the Brookhaven Alternating Gradient Synchrotron [28].
1981	Kai M. Siegbahn	Siegbahn invented a weak-focusing principle for betatrons in 1944 with which he made significant improvements in high-resolution electron spectroscopy [29].
1983	William A. Fowler	Fowler collaborated on and analyzed accelerator-based experiments in 1958 [30], which he used to support his hypothesis on stellar-fusion processes in 1957 [31].
1984	Carlo Rubbia and Simon van der Meer	Rubbia led a team of physicists who observed the intermediate vector bosons W and Z in 1983 using CERN's proton-antiproton collider [32], and van der Meer developed much of the instrumentation needed for these experiments [33].
1986	Ernst Ruska	Ruska built the first electron microscope in 1933 based upon a magnetic optical system that provided large magnification [34].
1988	Leon M. Lederman, Melvin Schwartz, and Jack Steinberger	Lederman, Schwartz, and Steinberger discovered the muon neutrino in 1962 using Brookhaven's Alternating Gradient Synchrotron [35].
1989	Wolfgang Paul	Paul's idea in the early 1950s of building ion traps grew out of accelerator physics [36].
1990	Jerome I. Friedman, Henry W. Kendall, and Richard E. Taylor	Friedman, Kendall, and Taylor's experiments in 1974 on deep inelastic scattering of electrons on protons and bound neutrons used the SLAC linac [37].
1992	Georges Charpak	Charpak's development of multiwire proportional chambers in 1970 were made possible by accelerator-based testing at CERN [38].
1995	Martin L. Perl	Perl discovered the tau lepton in 1975 using Stanford's SPEAR collider [39].
2004	David J. Gross, Frank Wilczek, and H. David Politzer	Gross, Wilczek, and Politzer discovered asymptotic freedom in the theory of strong interactions in 1973 based upon results from the SLAC linac on electron-proton scattering [40].
2008	Makoto Kobayashi and Toshihide Maskawa	Kobayashi and Maskawa's theory of quark mixing in 1973 was confirmed by results from the KEKB accelerator at KEK (High Energy Accelerator Research Organization) in Tsukuba, Ibaraki Prefecture, Japan, and the PEP II (Positron Electron Project II) at SLAC [41], which showed that quark mixing in the six-quark model is the dominant source of broken symmetry [42].



2013: François Englert and Peter W. Higgs "for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles, and which recently was confirmed through the discovery of the predicted fundamental particle, by the ATLAS and CMS experiments at **CERN's Large Hadron Collider**"

20 Nobels with X-rays

Chemistry

1936 Peter Debye

1962 Max Perutz and Sir John Kendrew

1976 William Lipscomb

1985 Herbert Hauptman and Jerome Karle

1988 Johann Deisenhofer, Robert Huber and Hartmut Michel

1997 Paul D. Boyer and John E. Walker

2003 Peter Agre and Roderick Mackinnon

2009 V. Ramakrishnan, Th. A. Steitz, A. E. Yonath

2012 Robert J. Lefkowitz and Brian K. Kobilka

Physics

1901 Wilhelm Rontgen

1914 Max von Laue

1915 Sir William Bragg and son

1917 C. G. Barkla

1924 Manne Siegbahn

1927 A. H. Compton

1981 Kai Siegbahn


Medicine

1946 Hermann Muller

1962 Frances Crick, James Watson and Maurice Wilkins

1979 Alan Cormack and Godfrey Hounsfield

THE ELECTROMAGNETIC SPECTRUM



Wavelength continuously tunable !

Imaging things

on all length and time scales
using accelerators,

e.g. latest X-Ray and
computational technologies
(developed at accelerators)

Spatial Scales


Meters
(10^0)

Centimeters
(10^{-2})

Millimeters
(10^{-3})

Micrometers
(10^{-6})

Nanometers
(10^{-9})



S
T
R
U
C
T
U
R
E

Materials


On a typical day in Europe

Modern day materials

Accelerators had an impact on a wide range of materials



Modern day materials



Synchrotron Light Sources: about 50 storage ring based



60'000 users
world-wide

Established, mature technology



The «brightness» of a light source




$$\text{Brightness} = \text{constant} \times \frac{F}{S \times \Omega}$$

X – Rays Brightness

Average Brightness



Bertha Roentgen's hand (exposure: 20 min)



Easter morning 1900: 5th / the automobile.




Easter morning 1913: 5th Ave, New York City. Spot the horse.



Brightness: disruptive change

- X-ray Tubes
- Storage Rings
- FELs
- ? Compact sources ?



Sources of Synchrotron Radiation


Bright beams of particles: phase space density

Incoherent,
spontaneous
emission of light:




Large phase space

Coherent, stimulated
emission of light



Undulators




$$T_{obs} = T_{emit} (1 - \beta)$$

$$\lambda_{light} \approx \frac{\lambda_u}{2\gamma^2}$$

3 types of storage ring sources:


1. Bending magnets: $B \sim N_e$





3 types of storage ring sources:

2. Wigglers:

$$B \sim N_e N_w \times 10$$




large undulations




3 types of storage ring sources:


3. Undulators:



Anatomy of a light source



Undulator line width



Undulator of infinite length

$$N_u = \infty \Rightarrow \frac{\Delta\lambda}{\lambda} = 0$$

Finite length undulator


- radiation pulse has as many periods as the undulator
- the line width is

$$\frac{\Delta\lambda}{\lambda} \sim \frac{1}{N_u}$$


Due to the electron energy spread

$$\frac{\Delta\lambda}{\lambda} = 2 \frac{\sigma_E}{E}$$



Undulator based beamline



Undulator radiation



$$\lambda = \frac{\lambda_u}{2n\gamma^2} \left(1 + \frac{K^2}{2} + \gamma^2 \theta^2 \right)$$



Particle beam emittance

Source
area, S



Angular
divergence, Ω

$$\text{Emittance} = S \times \Omega$$


X-ray emittance from electron source: a convolution of electron and photon phase space

$$\text{Brightness} = \frac{\Phi}{(2\pi)^2 \Sigma_x \Sigma_{x'} \Sigma_y \Sigma_{y'}}$$



$$\Sigma^2 = \sigma_e^2 + \sigma_\gamma^2$$

$$\sigma_\gamma' = \sqrt{\frac{\lambda}{L}}$$

$$\sigma_\gamma = \frac{\sqrt{\lambda L}}{4\pi}$$




Undulator radiation
from 6 GeV beam with
zero emittance, energy
spread (example ESRF)




Emittance $4 \text{ nm}\cdot\text{rad}$,
1% coupling,
finite energy spread

Diffraction limited storage rings

HITTING THE DIFFRACTION LIMIT




BRIGHTNESS:



Light of wavelength
 λ
focused to spot size
 Δx
will diffract with angle
 $\Delta\psi = \sim \lambda/\Delta x$

Diffraction limited light sources



LEAPS is the largest consortium of analytical facilities world-wide and further expanding its service to an interdisciplinary European user community

- 19** facilities - **16** institutions - **10** countries
- > **300** operating End Stations
- > **1.000.000** h beamtime /year
- > **5.000** publications/year
- > **15** spin off companies
- > **35.000** users from all EU & beyond
researchers from all research area



Construction and Operation (~ 800 M€/year) through national funding

UPGRADES



Storage Rings

Hard Energy X-ray Facilities

- ESRF
- ESRF EBS
- PETRA III
- PETRA IV*

Medium Energy X-ray Facilities

- ALBA
- ALBA II
- ANKA
- Diamond
- Diamond II*
- MAX IV 3 GeV
- MAX IV 3 GeV* ring upgrade
- SOLEIL
- SOLEIL ring upgrade
- SLS
- SLS2.0


Soft Energy X-ray Facilities


- MAX IV 1.5 GeV
- ELETTRA
- ELETTRA II
- BESSY II
- BESSY III*
- SOLARIS
- ASTRID2
- MLS
- DAFNE-Light

2015 2020 2025 2030 2035 2040

*facility planning (subject to approval)

■ Planning Period
 ■ Construction
 ■ User Operation





High Temperature Superconductors (HTS)
example of technology development

HTS superconducting magnet technology: developments

Using non-insulated HTS tapes:


18.2 Tesla field reached recently



HTS superconducting magnet technology undulators

Using bulk HTS material: can reach 2 Tesla for very short period magnets


Put the structure into a solenoid magnet, cool it and trap the field



Transverse coherence

- High brightness gives coherence
- Wave optics methods for X-rays (all chapters in Born & Wolf)
- Holography


The knee of a spider





phase contrast imaging

Phase contrast X-Ray imaging: improved soft tissue contrast

Absorption



Phase contrast




Same dose

X-ray Radiography of a fish

conventional
Absorption

a

(+ details c , e, g)



a

2 mm




b

Phase contrast
Microscopy

b

(+ details d, f, h)




c



d




e



f




g



h


Imaging for Life Science Applications

absorption




17.5 keV,
synchrotron results

phase contrast



Into the hospital ?


Source photon degeneracy parameter



Photons per spatial and temporal mode

$$\delta = B \lambda^3 / 4c = F_c \tau_c$$

Muscles and tracheal network *during* flight



R. Mokso *et al.*, Scientific Reports 5 8727 (2015)

Fake news?



Intel Core Pentium G3260 (3300) Dual Core, 22 nm

**Bloomberg
Businessweek**

October 8, 2018

The Big Hack




How China used
a tiny chip to
infiltrate America's
top companies

Holler et al., Nature 543, 402–406 (16 March 2017)


X-Ray tomography

Architecture of artificial and natural intelligence on all scales

Nature Electronics 2, 464-470 (2019)



Brain of a mouse in 3-D
Miettinen et al.







SLS 2.0: upscaling Swiss Light Source

$$B = \frac{\text{flux}}{\text{emittance}}$$

Brilliance



HUMAN HAIR (0.017 millimeters)

SLS

SLS 2.0

Beam profile: less emittance and more photons



Power economy SLS2.0 vs. SLS

More radiated X-ray power for users
Less electricity consumption

SLS → SLS2.0

E_{e^-} 2.4 GeV → 2.7 GeV

P_{SR} 310 GeV → 365 kW


W_{elec}/y 24 GWh → 17 GWh

Key savings:

Electromagnets → Permanent magnets

Klystrons → Solid state amplifiers


standard pumps → modern pumps for cooling



Free Electron Lasers


COHERENT EMISSION BY THE ELECTRONS

Intensity $\propto N$




INCOHERENT EMISSION

Intensity $\propto N^2$




COHERENT EMISSION

FIRST DEMONSTRATIONS OF COHERENT EMISSION (1989-1990)



180 MeV electrons



T. Nakazato et al., Tohoku University, Japan




30 MeV electrons

J. Ohkuma et al., Osaka University, Japan

MUCH HIGHER BRIGHTNESS CAN BE REACHED WHEN THE ELECTRONS COOPERATE





INCOHERENT EMISSION



COHERENT EMISSION

THE ELECTRON BEAM SHOULD BE $\sim 1 \text{ \AA}$
AS SMALL AS THE X-RAY WAVELENGTH!




Paul Scherrer Institute

SwissFEL




Free Electron Laser Keywords:

- Linac driven, single-pass
(no optics for cavity mirrors)




FEL Principles

Z. Huang





- Electrons **slip** behind EM wave by λ_1 per undulator period (λ_u)




- Due to sustained interaction, some electrons lose energy, while others gain \rightarrow energy modulation at λ_1
- e^- losing energy slow down, and e^- gaining energy catch up \rightarrow density modulation at λ_1 (microbunching)
- Microbunched beam radiates coherently at λ_1 , enhancing the process \rightarrow exponential growth of radiation power


X-Ray Free Electron Lasers



Spin-off product from basic research



Ultrafast Sources and Science



1878: E. Muybridge at Stanford


Tracing motion of animals
by spark photography



E. Muybridge




L. Stanford



Muybridge and Stanford disagree whether all feet leave the ground at one time during the gallop...

E. Muybridge, *Animals in Motion*, ed. by L. S. Brown (Dover Pub. Co., New York 1957).


FELs and ERLs COMPLEMENT the Ring sources




Compact accelerators:
sources of photons, neutrons, electrons etc.

Compact

Compton backscattering sources of hard X-Rays




Compact Light Source




Compact accelerators




50 MHz Ring Cyclotron



12 GHz CLIC




Laser
THz?



The
Economist

Disruptive storage rings technology change:
a much brighter compact sources
e.g. for shorter wavelength lithography

COSAMI design
5 x 12 meter
Footprint



Advanced Accelerator
Technologies

ENGINES OF DISCOVERY



A Century of Particle Accelerators

Andrew Sessler • Edmund Wilson

« Le seul véritable voyage ... ce ne serait pas d'aller vers de nouveaux paysages, mais d'avoir d'autres yeux, de voir l'univers avec les yeux d'un autre, de cent autres, de voir les cent univers que chacun d'eux voit, que chacun d'eux est. »

(Marcel Proust, La Prisonnière, 1923)

“The real voyage of discovery consists not in seeking new landscapes but in having new eyes”

Marcel Proust