


Update on particle tracking studies for asynchronous beam dumps

# TCDQ positioning and losses on TCT's

Brennan & Thomas

# IR6 Schematic Overview



# Tracking results

- Initial results already present in MPP in March 2009
- 7 TeV tracking now made with more 'realistic' setup; collimator positions, TCDQ settings, orbit, beta-beating...
- For some seeds IR7 collimators do not intercept beam.
  - Expected, as beta-beating changes phase advances
- But also...losses seen occasionally on some TCTs during asynchronous dump
  - Was puzzling as retraction is  $0.8 \sigma$ ...
  - Investigated further and is now understood...

# Collimator settings hierarchy

Values assumed in tracking


IR	Type	450 GeV [ $\sigma$ ]	7 TeV [ $\sigma$ ]
IR 1/IR 5	TCT	17.0	8.3
	TCL	10.0	10.0
IR 2/IR 8	TCLI	6.8	15.0
	TDI	6.8	17.0
	TCT	17.0	8.3
IR 3	TCP	8.0	15.0
	TCSG	9.3	18.0
	TCLA	10.0	20.0
IR 6	TCSG	7.0	7.5
	TCDQ	8.0	8.0
IR 7	TCP	5.7	6.0
	TCSG	6.7	7.0
	TCLA	10.0	10.0

# MKD kick waveform used


MKD system kick – pre-trigger case  
calculated from measured waveforms  
with 1000ns retriggering delay plus 15 ns  
/ generator

Worst case generator O (largest  $\beta_x$ )


Nominal





MKD.A pre-trigger



MKD.O pre-trigger







# Measured prefire pulse form – (from pre-trigger last week!)



Rise time in the very beginning slower than assumed in MAD-X model – data extracted for dedicated simulations.... (Thanks to Jan and Etienne)

# 7 TeV prefire case




Average losses – some seeds worse!

# 7 TeV tracking results

'sigma'	With all errors						Clean machine					
	Prefire case			Normal case			Prefire case			Normal case		
	8	10	15	8	10	15	8	10	15	8	10	15
'TCDQ.4R6'	4,66E+12	4,07E+12	3,11E+12	3,08E+12	2,85E+12	2,42E+12	4,81E+12	4,15E+12	3,11E+12	3,12E+12	2,88E+12	2,42E+12
'TCDQM.4R6'	0	0	0	0	0	0	0	0	0	0	0	0
'TCLA.A7R7.B1'	0	2,76E+08	8,18E+10	0	0	3,87E+10	0	0	7,08E+10	0	0	3,40E+10
'TCLA.C6R7.B1'	0	0	1,32E+10	0	0	5,84E+09	0	0	1,01E+10	0	0	5,06E+09
'TCP.C6L7.B1'	2,39E+09	8,83E+09	4,60E+10	1,66E+09	4,19E+09	7,54E+10	0	0	1,01E+10	0	0	5,98E+09
'TCSG.4R6'	2,32E+11	1,84E+11	1,24E+11	9,99E+10	8,11E+10	6,46E+10	1,79E+11	1,35E+11	8,56E+10	6,95E+10	4,83E+10	3,50E+10
'TCSG.6R7.B1'	7,62E+10	2,38E+11	4,53E+11	2,87E+10	8,44E+10	1,64E+11	5,80E+10	2,82E+11	5,79E+11	2,21E+10	8,51E+10	2,01E+11
'TCSG.A4L7.B1'	9,20E+07	1,84E+08	1,84E+08	0	0	4,60E+07	x	x	x	x	x	x
'TCSG.A4R7.B1'	x	x	x	4,60E+07	4,60E+07	4,60E+07	x	x	x	x	x	x
'TCSG.A5L7.B1'	x	x	x	0	0	4,60E+07	x	x	x	x	x	x
'TCSG.A6L7.B1'	8,19E+09	1,56E+10	5,90E+10	3,17E+09	5,43E+09	6,56E+10	5,52E+09	1,75E+10	7,27E+10	3,68E+09	5,06E+09	2,48E+10
'TCSG.B5L7.B1'	x	x	x	0	0	6,90E+08	x	x	x	x	x	x
'TCSG.B5R7.B1'	9,20E+08	1,29E+09	1,66E+09	5,06E+08	5,06E+08	5,52E+08	1,84E+09	1,84E+09	1,84E+09	9,20E+08	9,20E+08	9,20E+08
'TCSG.D4L7.B1'	0	2,76E+08	5,36E+11	0	0	2,85E+11	0	0	5,58E+11	0	0	2,57E+11
'TCTH.4L1.B1'	7,36E+08	6,64E+10	1,04E+11	1,84E+08	2,54E+10	4,21E+10	0	6,53E+10	1,05E+11	0	3,40E+10	5,15E+10
'TCTH.4L2.B1'	2,76E+08	1,53E+11	2,11E+11	1,38E+08	5,77E+10	8,24E+10	0	1,64E+11	2,11E+11	0	6,16E+10	8,37E+10
'TCDSA.4L6.B1'	4,89E+12	5,14E+12	5,14E+12	2,00E+12	2,10E+12	2,10E+12	2,00E+12	2,10E+12	2,10E+12	2,00E+12	2,10E+12	2,10E+12

For first 10 seeds, TCTs were hit in every case!





# Tracking with 3.5 TeV


- Started to implement with latest 3.5 TeV settings from Adriana
  - 7 TeV sequence and strength adapted to 3.5 TeV (same tune)
  - TCDQ (TCSG) 11.6 (10.2), TCTH/V.IR1 12.8, etc.
  - Beta\* in: IP1=2m, IP2=10m, IP5=2m, IP8=3m
  - Crossing bumps off (2mm oscillation if on -> unmatched IP??) , separation on

Optics seems to be ok, but still some issues with MAD-X concerning tracking (awkward behavior) to be solved.

Work ongoing...

Why do we see losses on the TCT??

# MKD kick (pretrigger or asynch.)




# Single pass $\neq$ multiple pass!

For  $\Delta\psi = 90^\circ$ ;  $\theta = 0$ ,  $R > S$


$$R = \sqrt{S^2 + n^2}$$

n	R
0.0	7.50
0.5	7.52
1.0	7.57
1.5	7.65
2.0	7.76
2.5	7.91
3.0	8.08
3.5	8.28
4.0	8.50
4.5	8.75
5.0	9.01
5.5	9.30
6.0	9.60




Phase advances/errors and finite beam sizes make it worse....

$$R = \sqrt{S^2 + (S \tan \theta + n)^2}$$



# Consequences

- Even for  $90^\circ$  phase advance, for a single pass the TCDQ intercepts beam at desired setting **ONLY for PX=0**
- Transmitted particle amplitude (in sigma) depends on its PX and on phase advance MKD to TCDQ



MKD to...  
TCDQ entr  
TCDQ exit  
TCSG entr  
TCSG exit

Nominal  $\Delta\psi$

	Deg B1	Deg B2
TCDQ entr	93.55	93.75
TCDQ exit	94.47	94.65
TCSG entr	94.60	94.78
TCSG exit	94.71	94.89

# “Semi analytical” checks


- Used 3.5 TeV settings from Adriana 08/09/09
  - TCSG.IR6 at  $10.2 \sigma$ , *plus*  $1.0 \sigma$  setting/orbit error (0.8 mm)
- Beam distributions generated in normalised phase space
  - **Gaussian** (nominal emittance,  $2 \times$  nominal emittance)
  - **Parabolic** (uniform phase space density)
  - **Hybrid** Gaussian with parabolic tails (90-10% and 99-1%)
- Worst-case asynchronous dump applied
  - MKD.O5L6, 1000 ns retrigger delay + 15 ns per generator (slightly pessimistic at 3.5 TeV).
- Damage level of TCTs assumed to be  $3 \times 10^9$  at 3.5 TeV
  - $1 \times 10^9$  at 7 TeV, scaled by  $E^{1.7}$
- No help from collimators in cleaning insertions
- Worst-case phase advance taken for losses on TCT

# MKD-TCDQ phase advance errors

- Nominal  $\Delta\psi \sim 94.6^\circ$
- Run Twiss on LHC machine with all errors
  - LHC error table 9901, misalignments etc.
  - Corrected tune and chromaticity back to nominal
- Max. MKD – TCDQ phase advance  $106^\circ$ 
  - Max phase advance error is  $\pm 12^\circ$
  - RMS  $5.4^\circ$


MAD-X Simulation output 50 seeds:

nominal	93.80
avg	94.22
max	105.40
min	81.61
stdv	5.42
max dev. (pos.)	11.60
max dev. (pos.)	12.19









# Beam distributions used – tails included...







# Results

1. with  $1.15 \times 10^{11}$  p+/bunch at 25ns
  - Nominal intensity and spacing

- Nominal phase advance MKD to TCDQ (**94.6°**),  $1.15 \times 10^{11}$ , 25ns







- 12° phase advance error MKD to TCDQ (**106.6°**),  $1.15 \times 10^{11}$ , 25ns







# Results

2. with  $7.6 \times 10^{10}$  p+/bunch at 50ns
  - (assumed worst case for 3.5 TeV)
  - Total I of  $6 \times 10^{13}$  in 792 bunches

- Nominal phase advance MKD to TCDQ (**94.6°**),  $7.6 \times 10^{10}$ , 50ns



- 12° phase advance error MKD to TCDQ (**106.6°**),  $7.6 \times 10^{10}$ , 50ns



# Summary of minimum TCT positions [ $\sigma$ ]

Bunch pattern	1.15×10 <sup>11</sup> p+/bunch at 25ns			7.6×10 <sup>10</sup> p+/bunch at 50ns		
MKD-TCDQ phase	94.6°	100.6°	106.6°	94.6°	100.6°	106.6°
Gaussian nominal $\varepsilon_x$	11.3	11.7	12.2	12.2	11.5	11.9
Gaussian 2× nominal $\varepsilon_x$	11.6	11.9	12.5	11.3	11.7	12.2
Gaussian 2× nominal $\varepsilon_x$ with 10% parabolic tail	11.8	12.2	12.8	11.5	11.8	12.3
Parabolic (uniform X-PX)	12.5	13.1	13.8	12.2	12.8	13.3

- **12.3**  $\sigma$  for minimum TCT *position* would seem just OK.
  - What margin to add (setting-up accuracy, dynamic beta-beat, ...)? **0.5**  $\sigma$ ?
- **12.3 + (0.5)**  $\sigma$  as the minimum setting
  - Assuming 2× nominal  $\varepsilon_x$ , 10% parabolic tails, 12° MKD-TCDQ phase error, 7.6×10<sup>10</sup> p+/bunch at 50 ns, 1.0  $\sigma$  TCDQ setting up error.



# Conclusion

- **Losses on TCTs** seen in MAD-X tracking for asynchronous dumps are 'real'
- Particles transmitted with larger amplitudes than TCDQ/TCSG setting
  - Single-pass cut transmits particles with higher PX at larger sigma
- TCTs are not robust
  - TCT damage limit? 1% of a nominal bunch at 7 TeV???
- **Short-term – keep TCTHs sufficiently retracted wrt TCDQ/TCSG**
  - Suggest to keep at **12.3  $\sigma$**  PLUS setting up margin of **0.5 – 1.0  $\sigma$** 
    - TCDQ/TCSG position wrt beam : tolerance 1.0  $\sigma$  (SW interlock only...final value 0.5  $\sigma$ )
    - Worst-case MKD-TCDQ  $\Delta\psi$  : assume 106.6°
    - TCT damage limit : assume 3E9 at 3.5 TeV
    - Worst case X-PX beam distrn : 2x nominal  $\varepsilon_x$  with 10% in parabolic tail cut at 6  $\sigma$
  - Quantify situation for 5 and 7 TeV (could impact minimum  $\beta^*$ )
    - 3.5 TeV needs **2.6 – 3.1  $\sigma$**  between the settings. For higher energy, margin needed will get bigger as TCT damage limit is reduced.
  - Producing files of TCT impact parameters for FLUKA / thermomechanical study
- **Longer term**
  - Will study auxiliary collimators  $\sim\pm 8^\circ$  from TCDQ, upstream of TCTs...
  - (More) robust TCTs from collimation project...?

# Discussion...

- Losses on TCTs seen in MAD-X tracking for asynchronous dumps are 'real'
- Particles transmitted with larger amplitudes than TCDQ/TCSG setting
  - Single-pass cut transmits particles with higher PX at larger sigma
- TCTs are not robust
  - TCT damage limit? 1% of a nominal bunch at 7 TeV???