

Heavy Neutral Leptons at the Electron Ion Collider

Keping Xie

University of Pittsburgh

11th Workshop of the Long-Lived Particle Community
June 3, 2022

In collaboration with **B. Batell**, **T. Han** (Pitt) and **T. Ghosh** (HRI, India)
Snowmass whitepaper [2203.06705](#), and work to appear soon

Introduction and Motivation

- The Electron-Ion Collider (EIC) is approved by the U.S. DOE with an estimated cost of **\$1.6 to \$2.6 billion**, to be located at Brookhaven National Laboratory.
- The EIC features [\[1212.1701 and Yellow Report, 2103.05419\]](#).
 - Highly polarized (70%) electron and nucleon beams
 - Ions: proton, deuteron to uranium or lead
 - C.o.M energies: 20–100 GeV, upgradable to 140 GeV
 - high luminosity: $10^{33-34} \text{ cm}^{-2}\text{s}^{-1}$ (10-1000 times HERA)
- The EIC goals: (designed as a QCD machine)
 - The proton spin
 - The motion of quarks and gluons in the proton
 - the tomographic images of the proton
 - QCD matter at the extreme gluon density
 - Quark hadronization
- Other physics opportunities: EW and BSM [\[Snowmass whitepaper, 2203.13199\]](#).
We take the **Heavy Neutral Leptons** as a case study.


The Electron-Ion Collider (EIC)

- We want to maximize the machine reachability

$$e(10/18 \text{ GeV}) + p(100/275 \text{ GeV}).$$

- We assume the integrated luminosity to be $\mathcal{L} = 100 \text{ fb}^{-1}$.
- Primary physics goals require a multi-purpose Hermitic detector with excellent tracking resolution and particle ID capabilities over a broad momentum range.
- Detector still under design; see EIC Detector Requirements R&D Handbook

[http://www.eicug.org/web/sites/default/files/EIC_HANDBOOK_v1.2.pdf]



η	Resolution
Tracking (σ_p/p)	
$2.5 < \eta \leq 3.5$	$0.1\% \times p \oplus 2\%$
$1.0 < \eta \leq 2.5$	$0.05\% \times p \oplus 1\%$
$ \eta \leq 1.0$	$0.05\% \times p \oplus 0.5\%$
Electromagnetic calorimeter (σ_E/E)	
$-4.5 \leq \eta < -2.0$	$2\%/\sqrt{E}$
$-2.0 \leq \eta < -1.0$	$7\%/\sqrt{E}$
$-1.0 \leq \eta \leq 4.5$	$12\%/\sqrt{E}$
Hadronic calorimeter (σ_E/E)	
$1.0 < \eta \leq 3.5$	$50\%/\sqrt{E}$
$ \eta \leq 1.0$	$100\%/\sqrt{E}$

The Heavy Neutral Leptons (HNLs) at the EIC



- The HNLs are motivated by the potential connection to the neutrino mass generation, through the Type-I Seesaw Mechanism [Minkowski PLB '77, Gell-Mann et. al. '79, etc.]
- The Lagrangian

$$\mathcal{L} \supset y_V^{iI} L_i H N_I + \text{h.c.}$$

The interactions can be written as



$$\mathcal{L} \supset \frac{g}{\sqrt{2}} U_{iI} W_\mu^- \ell_i^\dagger \bar{\sigma}^\mu N_I + \frac{g}{2 c_W} U_{iI} Z_\mu \nu_i^\dagger \bar{\sigma}^\mu N_I + \text{h.c.}$$

- The HNL production and decays (lifetime)



The Prompt HNL Searches

- The HNLs decay promptly for larger masses and mixing angle
- Three channels are considered
 - Majorana: $e^+ 3j$
 - Majorana: $e^+ \mu^- j + \cancel{E}_T$
 - Dirac: $\ell^+ \ell^- j + \cancel{E}_T$




Majorana signature: e^+3j

- The strongest signal comes the lepton number violating e^+3j :
 $e^-p \rightarrow (N \rightarrow e^+jj)j$.
- The hadronic mode gives largest rate, and allows for full final state reconstruction.
- the main SM background comes from charge mis-identification: e^- fakes as e^+ .
- the e^+jj invariant mass window cut very efficiently

Cut selection	Signal		e^-jjj [pb]
	$m_N = 10$ GeV [pb]	$m_N = 50$ GeV [pb]	
Production	5.53	0.95	449
Exactly 1ℓ : $p_{T_\ell} > 2$ GeV, $0 < \eta_\ell < 3.5$	2.43	0.74	36.7
Exactly $3j$: $p_{T_{j_1}} > 20$ GeV, $p_{T_{j_{2,3}}} > 5$ GeV, $ \eta_{j_{1,2,3}} < 3.5$	0.84	0.43	1.30
Isolation: $\Delta R(\ell, j_{1,2,3}) > 0.4$	0.52	0.41	1.30
$\min(M(\ell j_\alpha j_\beta) - m_N) < 5$ GeV ($\alpha, \beta = 1, 2, 3$)	0.52 ×	×	4.31×10^{-2} 0.59
Require one e^+ [$f^{\text{MID}} = 0.1\%$]	0.52 ×	×	4.31×10^{-5} 5.93×10^{-4}
Require one e^+ [$f^{\text{MID}} = 0.01\%$]	0.52 ×	×	4.31×10^{-6} 5.93×10^{-5}


Other channels

- Majorana signature: $\mu^- e^+ j + \cancel{E}_T$, BK: $\gamma p \rightarrow (\tau^- \rightarrow \mu^- j \nu)(\tau^+ \rightarrow e^+ j \nu) j$
- Dirac search: $\ell^+ \ell^- j + \cancel{E}_T$





With the designed energy and integrated luminosity, EIC can probe HNL prompt decay in the mass range 1–100 GeV and the mixing able of the order 10^{-3} .

Long-Lived signature



- At a small mass/mixing angle, the HNLs are long lived
- The signature of displaced lepton with large transverse impact parameter.
- The SM background arises from the heavy-flavor decay $ep \rightarrow \nu(c \rightarrow D)$ and $eg \rightarrow e(c/b \rightarrow D/B)(\bar{c}/\bar{b} \rightarrow \bar{D}/\bar{B})$.
 - At large impact parameter $d_T = 20$ mm, no SM background.
 - At small impact parameter $d_T = 2$ mm, we can perform cuts, such as $\Delta R_{j\ell} > 0.4$, to largely suppress the SM backgrounds [see backup slides].




Displaced vertex searches

- Event selection


$$p_T^j > 5 \text{ GeV}, p_T^\ell > 1 \text{ GeV}, |\eta_{j,\ell}| < 3.5.$$

- Transverse impact factor $d_T = 2(20) \text{ mm}$
- Cylinder detector configuration $r = 0.4 \text{ m}$, $d = 4 \text{ m}$.
- We show the EIC sensitivity to 5 displaced events



The EIC sensitivity

- the EIC can explore new parameter beyond the current bounds
- At low mass around 5 GeV, we can improve the current bound down to a lower mass
- Other experimental limits [\[1901.09966,2203.08039\]](#).




Invisible decay: $N \rightarrow \text{DMs}$

- Signal: $ep \rightarrow j + \cancel{E}_T$
- The mono-jet event

$$\begin{aligned}\sigma(p + e \rightarrow j + \cancel{E}_T) &= \sigma(p + e \rightarrow j + \nu_e) + \sigma(p + e \rightarrow j + N) \\ &= \sigma_{\text{SM}}(p + e \rightarrow j + \nu_e) [(1 - |U_e|^2) + |U_e|^2 \Phi(m_N)],\end{aligned}$$

- Sensitivity

$$\mathcal{S} = \frac{\Delta N}{\sqrt{N_{\text{SM}} + (\epsilon N_{\text{SM}})^2}} = 2, \quad \Delta N = |N - N_{\text{SM}}|.$$





[Existing bounds: 1709.07001]

Conclusions and Outlooks

- The EIC will open up a new QCD frontier. It is also interesting to ask the opportunity to search the BSM physics.
- We find the EIC has the potential to search HNLs, especially in the few GeV mass range, through the displaced vertex.
- EIC is able to probe HNL prompt decay in the mass range 1–100 GeV and mixing angle of the order 10^{-3} .
- Such studies can inform the EIC detector design, (e.g. tracking system for displaced particle searches)
- Other BSM physics exploration [\[Snowmass whitepaper, 2203.13199\]](#)
 - new light particle in 1–100 GeV mass range
 - SMEFT interactions [\[Boughezal, Petriello, Wiegand, 2004.00748\]](#)
 - lepton flavor violation [\[Gonderinger, Ramsey-Musolf, 1006.5063\]](#)
 - precision EW physics [\[Kumar et al. 1302.6263\]](#)
- It is very early days for the EIC. There is much more room for exploration.

The SM backgrounds for the displaced searches

SM background, $D(B) \rightarrow lX$, can be suppressed with isolation cut.



cuts	c	$c\bar{c}$	$b\bar{b}$
no cut	0.427		
$p_{T,l} > 2\text{GeV}, \eta_l < 3.5, \Delta R_{ll} > 0.3$	0.135	13.0	0.151
$p_{T,j} > 5\text{GeV}, \eta_j < 3.5, \Delta R_{jj} > 0.4$	0.0529	0.0855	$5.02e-3$
$\Delta R_{jl} > 0.4$	-	-	$1.73e-3$