

Phase retrieval by angular streaking of XUV atomic ionization

A.S. Kheifets^a, V.V. Serov^b, I.A. Ivanov^c, A. Li Wang^d and J.P. Cryan^d

^a*Research School of Physics, The Australian National University, Canberra ACT 2601, Australia.*

^b*General, Theoretical and Computer Physics, Saratov State University, Saratov 410012, Russia.*

^c*Center for Relativistic Laser Science, Institute for Basic Science (IBS), Gwangju 61005, South Korea.*

^d*Stanford PULSE Institute, SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA.*

Angular streaking of extreme ultraviolet (XUV) atomic ionization with a circularly polarized IR laser radiation has become a powerful tool for a shot-to-shot characterization of isolated attosecond pulses from free-electron laser sources [?, ?]. This technique is analogous to the attosecond streak camera [?] in that XUV pulses are the primary source of ionization while it is common to the attoclock [?] in that the ionized electrons interact with a circularly polarized laser field which alters the photoelectron momentum distribution. Both the streak camera and the attoclock are sensitive to the ionization phase that can be converted to the atomic time delay and the tunneling time, respectively. Here [?] we demonstrate, for the first time, an accurate phase retrieval of XUV atomic ionization by angular streaking.

In our numerical simulations, we solve the time-dependent Schrödinger equation for an atom ionized by a short XUV pulse in the presence of circularly polarized laser radiation. Our qualitative analysis is based on the strong field approximation [?] and the lowest order perturbation theory [?]. The two alternative techniques suggest the phase retrieval methods that can be applied both to the simulated and experimental data. In simulations, the XUV/IR delay is systematically varied. In an experiment, a second ionization feature in the same target or a mixed target could be used as a reference.

We test the proposed technique by comparing the XUV ionization phase of the hydrogen atom delivered by angular streaking with the analogous set of data returned by a well established RABBITT (reconstruction of attosecond beating by interference of two-photon transitions) technique [?]. The two sets of data agree very well in a wide range of XUV photon energies.

[1] N. Hartmann *et al.*, *Nature Photonics* **12**, 215 (2018)

[2] J. Duris *et al.*, *Nature Photonics* **14**, 30 (2020).

[3] J. Itatani *et al.*, *Phys. Rev. Lett.* **88**, 173903 (2002).

[4] P. Eckle *et al.*, *Nature Physics* **4**, 565 (2008).

[5] A S Kheifets *et al.*, *arXiv* 2202.06147 (2022).

Kazansky2016 A K Kazansky *et al.*, *Phys. Rev. A* **93**, 013407 (2016).

The abstract should be saved as a pdf file (removing this sentence) and uploaded via the abstract submission tab of the conference website at

<https://aip-congress.org.au/submissions.html>