

Higgs Boson properties and couplings

Snowmass EF01 report


Caterina Vernieri, Sally Dawson

April 20, 2022


ECFA WG1 Workshop

Higgs and the exploration of the EF

- The Standard Model is not a complete theory
 - Gravity, neutrino mass, dark matter ...
- The Higgs boson is a **potential window to probe physics Beyond the Standard Model**
 - Searches for additional scalars
 - (Interplay with) Precision measurements of the **Higgs boson properties**
 - Higgs Global fits




Higgs physics at the HL-LHC



The High Luminosity era of LHC will dramatically expand the physics reach for Higgs physics:

- **2-5% precision for most of Higgs couplings**
- **Larger uncertainties on $Z\gamma$ and charm**
- **<50% on the self-coupling**
- **Higgs width 5%**


Higgs physics at the HL-LHC



The High Luminosity era of LHC will dramatically expand the physics reach for Higgs physics:

- **2-5% precision for most of Higgs couplings**
- **Larger uncertainties on $Z\gamma$ and charm**
- **<50% on the self-coupling**
- **Higgs width 5%**

Higgs as a guide



H couplings to:

$O(5-15)\%$

$O(0.1-1)\%$

$O(1)\text{‰}$

H self-coupling to


$<O(50)\%$

$O(20)\%$

$O(1)\%$

Wish list beyond HL-LHC:

1. Establish Yukawa couplings to light flavor \Rightarrow precision & lumi
2. Establish self-coupling \Rightarrow high energy



H couplings to:

$O(5-15)\%$

$O(0.1-1)\%$

$O(1)\text{‰}$

H self-coupling to



$<O(50)\%$

$O(20)\%$


$O(1)\%$

Why leptons?

- Initial state well defined (& polarization) \Rightarrow High-precision measurements
- Higgs bosons appear in 1 in 100 events \Rightarrow Clean experimental environment and trigger-less readout



Higgs at e^+e^-



- ZH is dominant at **250 GeV**
- Above **500 GeV**
 - H $\nu\nu$ dominates
 - ttH opens up
 - HH production accessible with ZHH


Which collider?

LEPTON COLLIDERS

- **Circular e+e-** (CEPC, FCC-ee)
 - **90-350 GeV**
 - *strongly limited by synchrotron radiation above 350– 400 GeV*
- **Linear e+e-** (ILC, CLIC, C³)
 - **250 GeV — > 1 TeV**
 - *Reach higher energies, and can use polarized beams*
- **μ+μ-**
 - **3-30 TeV**

HADRON COLLIDERS

- **75-200 TeV** (FCC-hh)



Exploring the complementarity between e^+e^- and LHC will lead to the most precise measurements

- ***What do we want to learn on top of what HL-LHC will deliver?***
- ***Timelines matter - ideally the next machine will minimize the gap with HL-LHC, also:***
 - As we gain knowledge, how do we prioritize the Higgs measurements?
 - Which energy to target ?
 - top-Yukawa, HH, extended Higgs sector with more generic Yukawa coupling scenarios will require > 500 GeV
 - LC approach is to start at 250 GeV and then ~ 500 GeV - is it enough?
 - ***Which measurement are to be prioritized after the 250 GeV run?***
 - How relevant is to have polarized beams?

EF012-whitereport

Sally Dawson Adrey Korytov Patrick Meade
Isobel Ojalvo Caterina Vernieri

March 2022

1 Introduction

2 Why Higgs is great

2.1 Centrality of Higgs in SM

2.2 Questions the Higgs leaves us

2.3 Status report of what we know for sure and what's missing experimentally (self couplings, light yukawas, "unitarization")

2.4 New types of measurements vs increased precision (e.g. differential)

2.5 What BSM Higgs physics can influence

2.6 Models of BSM Higgs

Models addressing physics questions vs exploring Higgs like possibilities/property modifications

- EFT
- Higgs portal
- Singlet
- Doublet
- Higgs importance in larger frameworks?
- Naturalness, Compositeness

DRAFT!

3 Mass, width, spin measurements

4 Update on theory calculations of Higgs rates

5 Impact of differential measurements

6 Higgs coupling measurements with emphasis on light Yukawas

7 CP violating Higgs coupling measurements

8 Update on prospects for observing HH

9 Update on new analysis techniques for Higgs physics and what they mean

10 How coupling measurements and HH impact understanding of EFTs

11 Why go beyond the SM?

12 The big motivations for BSM Higgs

13 Some models showing complementarity of direct searches and precision measurements: Singlet, 2HDM, ?

14 Targets for precision based on models

15 Detector/accelerator requirements to observe new physics


Expanding 1:

- - How do we go further, energy, luminosity, clean environment (snowmass options, connect to AF, IF)

Higgs couplings at future colliders




- Future colliders under consideration will improve with respect to the HL-LHC the understanding of the Higgs boson couplings - 1-5%
- **Coupling to charm** quark could be measured with an accuracy of $\sim 1\%$ in future e+e- machines
- **Couplings to $\mu/\gamma/Z\gamma$** benefit the most from the large dataset available at HL-LHC
- At low energy top-Higgs coupling is not accessible at future lepton colliders



Higgs couplings at future colliders




- Future colliders under consideration will improve with respect to the HL-LHC the understanding of the Higgs boson couplings - 1-5%
 - **Coupling to charm** quark could be measured with an accuracy of $\sim 1\%$ in future e+e- machines
 - **Couplings to $\mu/\gamma/Z\gamma$** benefit the most from the large dataset available at HL-LHC
 - At low energy top-Higgs coupling is not accessible at future lepton colliders



To be updated: new studies for Snowmass



New from HL-LHC



YR projections based on analyses of partial Run 2 dataset
Full Run 2 measurements have drastically improved previous results
We need to update our HL-LHC projections

Updated Higgs couplings for ILC


coupling	ILC250		ILC500		ILC1000	
	full	no BSM	full	no BSM	full	no BSM
hZZ	0.49	0.38	0.35	0.20	0.34	0.16
hWW	0.48	0.38	0.35	0.20	0.34	0.16
hbb	0.99	0.80	0.58	0.43	0.47	0.31
$h\tau\tau$	1.1	0.95	0.75	0.63	0.63	0.52
hgg	1.6	1.6	0.96	0.91	0.67	0.59
hcc	1.8	1.7	1.2	1.1	0.79	0.72
$h\gamma\gamma$	1.1	1.0	1.0	0.96	0.94	0.89
$h\gamma Z$	8.9	8.9	6.5	6.5	6.4	6.4
$h\mu\mu$	4.0	4.0	3.8	3.7	3.4	3.4
htt	—	—	6.3	6.3	1.0	1.0
hhh	—	—	20	20	10	10
Γ_{tot}	2.3	1.3	1.6	0.70	1.4	0.50
Γ_{inv}	0.36	—	0.32	—	0.32	—



Note C³ would run at 550 GeV, a factor 2 improvement to the top-Yukawa coupling ()*


Updated Higgs couplings for ILC

coupling	ILC250		ILC500		ILC1000	
	full	no BSM	full	no BSM	full	no BSM
hZZ	0.49	0.38	0.35	0.20	0.34	0.16
hWW	0.48	0.38	0.35	0.20	0.34	0.16
hbb	0.99	0.80	0.58	0.43	0.47	0.31
$h\tau\tau$	1.1	0.95	0.75	0.63	0.63	0.52
hgg	1.6	1.6	0.96	0.91	0.67	0.59
hcc	1.8	1.7	1.2	1.1	0.79	0.72
$h\gamma\gamma$	1.1	1.0	1.0	0.96	0.94	0.89
$h\gamma Z$	8.9	8.9	6.5	6.5	6.4	6.4
$h\mu\mu$	4.0	4.0	3.8	3.7	3.4	3.4
htt	—	—	6.3	6.3	1.0	1.0
hhh	—	—	20	20	10	10
Γ_{tot}	2.3	1.3	1.6	0.70	1.4	0.50
Γ_{inv}	0.36	—	0.32	—	0.32	—



Note C^3 would run at 550 GeV, a factor 2 improvement to the top-Yukawa coupling (*)




Combined limit of $\kappa_s < 6.74$ at 95% CL with 900/fb at 250 GeV



- There are extensive comparisons between the FCC-ee plan and the C³/ILC runs that show they are rather **compatible to study the Higgs Boson**
- When analyzing Higgs couplings with SMEFT, 2 ab⁻¹ of polarized running is essentially equivalent to 5 ab⁻¹ of unpolarized running.
 - **Electron polarization is essential** for this
 - There is almost no difference in the expectation with and without **positron polarization**.
 - more cross-checks of systematic errors.
 - relevant at high energy (> TeV) where the most important cross sections are initiated from e⁻L e⁺R

coupling	2/ab-250 +4/ab-500		5/ab-250 + 1.5/ab-350	
	pol.	pol.	unpol.	unpol
HZZ	0.50	0.35	0.41	0.34
HWW	0.50	0.35	0.42	0.35
Hbb	0.99	0.59	0.72	0.62
$H\tau\tau$	1.1	0.75	0.81	0.71
Hgg	1.6	0.96	1.1	0.96
Hcc	1.8	1.2	1.2	1.1
$H\gamma\gamma$	1.1	1.0	1.0	1.0
$H\gamma Z$	9.1	6.6	9.5	8.1
$H\mu\mu$	4.0	3.8	3.8	3.7
Htt	-	6.3	-	-
HHH	-	27	-	-
Γ_{tot}	2.3	1.6	1.6	1.4
Γ_{inv}	0.36	0.32	0.34	0.30
Γ_{other}	1.6	1.2	1.1	0.94

Higgs couplings at the muon collider





Higgs-electron Yukawa



- **Electron** Yukawa at FCC-ee with a dedicated 4 years run at the Higgs mass
 - $\kappa_e < 1.6$ at 95% CL

NEW!



- Goal is to sharpen theoretical expectations / models (see summary from Andrei Gritsan)
 - Connect to broader EFT and distinguish between linear and quadratic effects in the observables
- CP measurement using the **τ Yukawa coupling** at e+e- at 250 GeV with a precision of 75 mrad
 - Higher energy stages, at a TeV or higher, can be used to measure CP effects in the HZZ coupling by studying the ZZ-fusion Higgs production process (to be followed up)
- **Higgs-top CP-structure** via the tth production at the HL-LHC, FCC-hh and muon colliders.

Bounds on α at 95% CL ($\kappa_t = 1$)	Channel	Collider	Luminosity
$ \alpha \lesssim 36^\circ$ [1]	dileptonic $t\bar{t}(h \rightarrow b\bar{b})$	HL-LHC	3 ab^{-1}
$ \alpha \lesssim 25^\circ$ [2]	$t\bar{t}(h \rightarrow \gamma\gamma)$ combination	HL-LHC	3 ab^{-1}
$ \alpha \lesssim 3^\circ$ [1]	dileptonic $t\bar{t}(h \rightarrow b\bar{b})$	100 TeV FCC	30 ab^{-1}
$ \alpha \lesssim 9^\circ$ [3]	semileptonic $t\bar{t}(h \rightarrow b\bar{b})$	10 TeV $\mu^+\mu^-$	10 ab^{-1}
$ \alpha \lesssim 3^\circ$ [3]	semileptonic $t\bar{t}(h \rightarrow b\bar{b})$	30 TeV $\mu^+\mu^-$	10 ab^{-1}

The Higgs self-coupling at future colliders

- CMS & ATLAS projections have been updated
 - New combination can be done ?
- New since the YR, **FCC-hh** : 2.9-5.5% depending on the systematic assumptions (arXiv:2004.03505)
- **Muon collider** 25% (6%) at 3 (10) TeV

	3 TeV μ -coll. $L \approx 1 \text{ ab}^{-1}$	10 TeV μ -coll. $L = 10 \text{ ab}^{-1}$	14 TeV μ -coll. $L \approx 20 \text{ ab}^{-1}$	30 TeV μ -coll. $L = 90 \text{ ab}^{-1}$
		68% prob. interval		
$\delta\kappa_\lambda$	$[-0.27, 0.35] \cup [0.85, 0.94]$	$[-0.035, 0.037]$	$[-0.024, 0.025]$	$[-0.011, 0.012]$
	$\rightarrow [-0.15, 0.16] (2 \times L)$			


collider	single- H	HH	combined
HL-LHC	100-200%	50%	50%
CEPC ₂₄₀	49%	–	49%
C ³ ILC ₂₅₀	49%	–	49%
C ³ ILC ₅₀₀	38%	27%	22%
ILC ₁₀₀₀	36%	10%	10%
CLIC ₃₈₀	50%	–	50%
CLIC ₁₅₀₀	49%	36%	29%
CLIC ₃₀₀₀	49%	9%	9%
FCC-ee	33%	–	33%
FCC-ee (4 IPs)	24%	–	24%
HE-LHC	-	15%	15%
* FCC-hh	-	5%	5%

These values are combined with an independent determination of the self-coupling with uncertainty 50% from the HL-LHC.

Higgs production at large p_T



- European Strategy Studies focused on inclusive measurements : **new opportunities for measurements of the Higgs couplings at large Q^2**
 - BSM effects often grow with energy
 - Clear impact on the extraction of EFT constraints via correlations among different processes and kinematical regimes
 - Also this helps mitigating systematic uncertainties and maximizes the robustness of the results
 - i.e. pile-up rejection and trigger capabilities
- Few **examples**:
 - VH at large invariant mass (double differential distributions sometime needed to restore BSM/SM interference)
 - Probing the HWW coupling at high Q^2 in $pp \rightarrow WH$ at large mass or in VBF is complementary to measure $BR(H \rightarrow WW)$
 - off-shell $gg \rightarrow H^* \rightarrow ZZ \rightarrow 4l$
 - Higgs + high- p_T jet




New result in VH to bb and H to $\tau\tau$



Challenges and Opportunities for Higgs Physics



- Why is exploring the Higgs important?
 1. It is a fundamental part of the SM
 2. In the SM, most everything except the Higgs mass is predicted, so we can make precise comparisons.
- Are existing theory calculations sufficient for the comparisons? How accurately do we need to measure?
- What do we learn from precision measurements of Higgs properties?
 - Precision is window to high scales. How to make the connection? How important are specific models? in SMEFT framework, $\delta\kappa \sim v^2/M^2$
- Higgs is window to high scale physics
 - Future e+e- colliders give increasingly precise measurements of Higgs couplings
 - Can we quantify the complementarity with direct searches at high energy machines?
 - This should be studied by exploring the complementarity between HL-LHC and future colliders (accounting for their different timelines).
- Does EWSB work the way we think it does?
 - How to best explain the importance of measuring triple Higgs coupling?
 - We can quantify how well different machines can do....but we need to explain what various target measurements imply
 - How to connect precision of measurement with reach for new particles in models that predict deviations on the self-coupling?

Unitarity Criteria



- Higgs couplings:
 - Include updated list of machines (muon collider, C³ are recent developments) and their parameters (including timelines)
 - Re-visit some of the assumptions (i.e. flavor..) since the ESG
 - Summary of latest theory cross sections (distributions too if available)
 - New Global fits
- Some example maps of new physics phase space to constraints on EFT operators
 - Plots that demonstrate in creative ways the BSM sensitivities of various measurements
- New physics benchmarks for resonant and non-resonant HH that we could use for interpretations as the precision on the self-coupling improves






Extra

Higgs physics at the HL-LHC




NEW!



First extrapolation of κ_c

NEW!





From 3.3 σ to 4.6 σ on the expected SM σ_{HH}


H to strange coupling



- Exploring ZH with Z going to leptons or neutrinos
- Combined limit of $\kappa_s < 6.74$ at 95% CL with 900/fb at 250 GeV (i.e. half dataset)

NEW!








Extending the sensitivity beyond LHC to 2HDM

A spontaneous flavour violating (SFV) 2HDM allows for large couplings of additional Higgs to strange/light quarks while suppressing flavour-changing neutral currents

- Progress on how to map BSM models to SMEFT constraints
 - Include complete 1-loop matching for other models, more NLO effects in fits, and more distributions





- Progress on how to map BSM models to SMEFT constraints
 - Include complete 1-loop matching for other models, more NLO effects in fits, and more distributions



Model independent bounds on new physics

Coupling (2σ)	HL-LHC	LHeC	HE-LHC	ILC	CLIC	CEPC	FCC-ee	FCC
Unitarity Bound			S2 S2'	250 500 1000	380 1500 3000		240 365	ee/eh/hh
$2\delta\kappa_V$ [%]	3.0	1.5	2.6 1.8	0.58 0.46 0.44	1.0 0.32 0.22	0.28	0.40 0.34	0.24
Λ_V (TeV)	6.0	9	6.4 7.7	14 15 16	10 18 22	20	16 18	21
$2\kappa_g$ [%]	4.6	7.2	3.8 2.4	4.6 1.94 1.32	5.0 2.6 1.8	3.0	3.4 2.0	0.98
Λ_g (TeV)	51	41	56 70	51 78 95	49 68 81	63	59 77	110
$2\kappa_\gamma$ [%]	3.8	15.2	3.2 2.4	13.4 6.8 3.8	196 10 4.4	7.4	9.4 7.8	0.58
Λ_γ (TeV)	120	61	130 150	65 92 120	17 76 110	88	78 86	310
$2\kappa_{Z\gamma}$ [%]	20	—	11.4 7.6	198 172 170	240 30 13.8	16.4	162 150	1.38
$\Lambda_{Z\gamma}$ (TeV)	34	—	45 55	11 12 12	10 28 41	37	12 12	130
$2\delta\kappa_t$ [%]	6.6	—	5.6 3.4	— 13.8 3.2	— — 5.4	—	— —	2.0
Λ_t (TeV)	13	—	14 18	— 9 19	— — 14	—	— —	24
$2\delta\kappa_b$ [%]	7.2	4.2	6.4 4.6	3.6 1.16 0.96	3.8 0.92 0.74	2.4	2.6 1.34	0.86
Λ_b (TeV)	80	100	85 100	110 200 220	110 220 250	140	130 180	230
$2\delta\kappa_\mu$ [%]	9.2	—	5.0 3.4	30 18.8 12.4	640 26 11.6	17.8	20 17.8	0.82
Λ_μ (TeV)	590	—	800 970	320 410 510	70 350 520	420	400 420	2000
$2\delta\kappa_\tau$ [%]	3.8	6.6	3.0 2.2	3.8 1.40 1.14	6.0 2.6 1.76	2.6	2.8 1.46	0.88
Λ_τ (TeV)	220	170	250 290	220 370 410	180 270 330	270	260 360	460
$2\delta\kappa_h$ [%]	94	—	40 40	58 54 20	92 72 22	34	38 38	10
Λ_h (TeV)	15	—	23 23	19 19 32	15 17 30	25	23 23	45



- **We will be working closely together with EF04 within the SMEFT framework:**
 - Estimate EFT uncertainties (NLO, dim-8 effects, linear vs quadratic...), new physics in backgrounds, theoretical constraints (positivity, analyticity)
 - More combined Higgs and top analysis
 1. effects of top dipoles or 4 fermion ops. with tops
 2. constraints on top EW couplings from their NLO effects in Higgs and diboson processes (particularly relevant for low-energy colliders below ttH threshold)
 - Include differential observables
 - Explore more flavor scenarios (and make connection with flavor data)
- SMEFT is a baseline, how we account for specific assumptions and model-dependency?
- Complementarity with new physics searches

Wish list for the global fit

Inputs: Higgs @ HL-LHC



in unit of %

EF04 link

HL-LHC	3 ab⁻¹ @ 14 TeV ATLAS+CMS (S2)				
prod.	ggH	VBF	WH	ZH	ttH
σ	-	-	-	-	-
$\sigma \times BR_{bb}$	19.1	-	8.3	4.6	10.2
$\sigma \times BR_{cc}$	-	-	-	-	-
$\sigma \times BR_{gg}$	-	-	-	-	-
$\sigma \times BR_{zz}$	2.5	9.5	32.1	58.3	15.2
$\sigma \times BR_{ww}$	2.5	5.5	9.9	12.8	6.6
$\sigma \times BR_{\tau\tau}$	4.5	3.9	-	-	10.2
$\sigma \times BR_{\gamma\gamma}$	2.5	7.9	9.9	13.2	5.9
$\sigma \times BR_{\gamma Z}$	24.4	51.2	-	-	-
$\sigma \times BR_{\mu\mu}$	11.1	30.7	-	-	-
$\sigma \times BR_{inv.}$	-	2.5	-	-	-
m_H	10-20MeV				

wishlist: correlation matrix; differential x-section is not included now, but can be considered if available

Accuracy vs. Luminosity



Scenario	Stage \sqrt{s} [GeV]	500			500 LumiUP		
		500	350	250	500	350	250
G-20	$\int \mathcal{L} dt$ [fb^{-1}]	1000	200	500	4000	-	-
	time [years]	5.5	1.3	3.1	8.3	-	-
H-20	$\int \mathcal{L} dt$ [fb^{-1}]	500	200	500	3500	-	1500
	time [years]	3.7	1.3	3.1	7.5	-	3.1
I-20	$\int \mathcal{L} dt$ [fb^{-1}]	500	200	500	3500	1500	-
	time [years]	3.7	1.3	3.1	7.5	3.4	-