

# *Imprint of quark flavor violating SUSY in $h(125)$ decays at future lepton colliders*

*K. Hidaka*

*Tokyo Gakugei University*

*Collaboration with*

*H. Eberl, E. Ginina (HEPHY Vienna)*

*References:*

*Phys. Rev. D 91 (2015) 015007 [arXiv:1411.2840 [hep-ph] ]*

*JHEP 1606 (2016) 143 [arXiv:1604.02366 [hep-ph]]*

*IJMP A34 (2019) 1950120 [arXiv:1812.08010 [hep-ph]]*

*PoS(EPS-HEP2021)594, 2021 [arXiv:2111.02713 [hep-ph]]*

*ILC White Paper for Snowmass 2021 [arXiv:2203.07622]*

*1st Workshop of the ECFA Higgs/Top/EW factory WG1 - HTE group,  
20 Apr. 2022, CREN*

# 1. Introduction

- *What is the SM-like Higgs boson discovered at LHC?*
- *It can be the SM Higgs boson.*
- *It can be a Higgs boson of New Physics.*
- *This is one of the most important issues in the present particle physics field!*
- *Here we study a possibility that it is the lightest Higgs boson  $h^0$  of the Minimal Supersymmetric Standard Model (MSSM), focusing on the decays  $h^0(125) \rightarrow c \bar{c}, b \bar{b}, b \bar{s}, \gamma \gamma, g g$ .*

## 2. MSSM with QFV

*Key parameters in this study are:*

\* *QFV parameters:  $\tilde{c}_{L/R} - \tilde{t}_{L/R}$  &  $\tilde{s}_{L/R} - \tilde{b}_{L/R}$  mixing parameters*

\* *QFC parameter:  $\tilde{t}_L - \tilde{t}_R$  &  $\tilde{b}_L - \tilde{b}_R$  mixing parameters*

*$M^2_{Q23} = (\tilde{c}_L - \tilde{t}_L$  mixing parameter)*

*$M^2_{U23} = (\tilde{c}_R - \tilde{t}_R$  mixing parameter)*

*$M^2_{D23} = (\tilde{s}_R - \tilde{b}_R$  mixing parameter)*

*$T_{U23} = (\tilde{c}_R - \tilde{t}_L$  mixing parameter)*

*$T_{U32} = (\tilde{c}_L - \tilde{t}_R$  mixing parameter)*

*$T_{U33} = (\tilde{t}_L - \tilde{t}_R$  mixing parameter)*

*$T_{D23} = (\tilde{s}_R - \tilde{b}_L$  mixing parameter)*

*$T_{D32} = (\tilde{s}_L - \tilde{b}_R$  mixing parameter)*

*$T_{D33} = (\tilde{b}_L - \tilde{b}_R$  mixing parameter)*

# 3. Constraints on the MSSM

*We respect the following experimental and theoretical constraints:*

*(1) The LHC limits on the masses of squarks, sleptons, gluino, charginos and neutralinos.*

*(2) The constraint on  $(m_{A/H^+}, \tan\beta)$  from MSSM Higgs boson search at LHC.*

*(3) The constraints on the QFV parameters from the B & K meson data.*

$$B(b \rightarrow s \gamma) \quad \Delta M_{B_s} \quad B(B_s \rightarrow \mu^+ \mu^-) \quad B(B_u^+ \rightarrow \tau^+ \nu) \quad \text{etc.}$$

*(4) The constraints from the observed Higgs boson mass and couplings at LHC ; e.g.*

$$121.6 \text{ GeV} < m_{h^0} < 128.6 \text{ GeV} \text{ (allowing for theoretical uncertainty) ,}$$
$$0.71 < \kappa_b < 1.43 \text{ (ATLAS), } 0.56 < \kappa_b < 1.70 \text{ (CMS)}$$

*(5) The experimental limit on SUSY contributions to the electroweak  $\rho$  parameter*

$$\Delta \rho (\text{SUSY}) < 0.0012.$$

*(6) Theoretical constraints from the vacuum stability conditions for the trilinear couplings  $T_{U\alpha\beta}$  and  $T_{D\alpha\beta}$ .*

## 4. Parameter scan

- We compute the  $h^0(125)$  decay widths in the *MSSM with QFV*.
- We take parameter scan ranges as follows:

$$1 \text{ TeV} < M_{\text{SUSY}} < 5 \text{ TeV}$$

$$10 < \tan\beta < 80$$

$$2500 < M_3 < 5000 \text{ GeV}$$

$$100 < M_2 < 2500 \text{ GeV}$$

$$100 < M_1 < 2500 \text{ GeV}$$

$$100 < \mu < 2500 \text{ GeV}$$

$$1350 < m_A(\text{pole}) < 6000 \text{ GeV}$$

*etc. etc.*

- *In the parameter scan, all of the relevant experimental and theoretical constraints are imposed.*
- *377180 parameter points are generated and 3208 points survive the constraints.*

## 5. $h^0 \rightarrow c \bar{c}, b \bar{b}, b \bar{s}$ in the MSSM

- We compute the decay widths  $\Gamma(h^0 \rightarrow c \bar{c})$ ,  $\Gamma(h^0 \rightarrow b \bar{b})$ , and  $\Gamma(h^0 \rightarrow b \bar{s})$  at full 1-loop level in the DRbar renormalization scheme in the **MSSM with QFV**.

- Main 1-loop correction to  $h^0 \rightarrow c \bar{c}$ :

**gluino - su loops [ su = ( $\tilde{t}$  -  $\tilde{c}$  mixture)]**

can be enhanced by large trilinear couplings  $T_{U23}, T_{U32}, T_{U33}$

- Main 1-loop corrections to  $h^0 \rightarrow b \bar{b}$  &  $b \bar{s}$ :

**gluino - sd loops [ sd = ( $\tilde{b}$  -  $\tilde{s}$  mixture)]**

can be enhanced by large trilinear couplings  $T_{D23}, T_{D32}, T_{D33}$


**chargino - su loops [ su = ( $\tilde{t}$  -  $\tilde{c}$  mixture)]**

can be enhanced by large trilinear couplings  $T_{U23}, T_{U32}, T_{U33}$

**In large  $\tilde{c}_{R/L} - \tilde{t}_{R/L}$  &  $\tilde{t}_L - \tilde{t}_R$  mixing scenario;**

$$h^0 \sim H_2^0$$

$$\tilde{u}_{1,2} \sim \tilde{c}_{R/L} + \tilde{t}_{R/L}$$



**In our scenario, “trilinear couplings” ( $\tilde{c}_R - \tilde{t}_L - H_2^0$ ,  $\tilde{c}_L - \tilde{t}_R - H_2^0$ ,  $\tilde{t}_L - \tilde{t}_R - H_2^0$  couplings) =  $(T_{U23} T_{U32}, T_{U33})$  are large!**

**$\tilde{u}_{1,2} - \tilde{u}_{1,2} - h^0$  couplings are large!**


**Gluino loop contributions can be large!**

**Deviation of  $\Gamma(h^0 \rightarrow c \bar{c})$  from SM width can be large!**

*In large  $\tilde{s}_{R/L} - \tilde{b}_{R/L}$  &  $\tilde{b}_L - \tilde{b}_R$  mixing scenario;*

$$h^0 \sim -s\alpha H_1^0 + c\alpha H_2^0$$

$$\tilde{d}_{1,2} \sim \tilde{s}_{R/L} + \tilde{b}_{R/L}$$



*In our scenario, “trilinear couplings“ ( $T_{D23} T_{D32}, T_{D33}$ ) =  $(\tilde{s}_R - \tilde{b}_L - H_1^0, \tilde{s}_L - \tilde{b}_R - H_1^0, \tilde{b}_L - \tilde{b}_R - H_1^0$  couplings) *are large!**

*$\tilde{d}_{1,2} - \tilde{d}_{1,2} - h^0$  couplings are large!*

*Glino loop contributions can be large!*

*Deviation of  $\Gamma(h^0 \rightarrow b \bar{b}/\bar{s})$  from SM width can be large!*




*In large  $\tilde{c}_{R/L} - \tilde{t}_{R/L}$  &  $\tilde{t}_L - \tilde{t}_R$  mixing scenario;*

$$h^0 \sim H_2^0$$

$$\tilde{u}_{1,2} \sim \tilde{c}_{R/L} + \tilde{t}_{R/L}$$

$$\tilde{\chi}^\pm \sim \tilde{W}^\pm + \tilde{H}^\pm$$



*In our scenario, “trilinear couplings” ( $\tilde{c}_R - \tilde{t}_L - H_2^0$ ,  $\tilde{c}_L - \tilde{t}_R - H_2^0$ ,  $\tilde{t}_L - \tilde{t}_R - H_2^0$  couplings) =  $(T_{U23} T_{U32}, T_{U33})$  are large!*

*$\tilde{u}_{1,2} - \tilde{u}_{1,2} - h^0$  couplings are large!*

*Chargino loop contributions can be large!*

*Deviation of  $\Gamma(h^0 \rightarrow b \bar{b}/\bar{s})$  from SM width can be large!*


## 5.1 Deviation of the width from the SM prediction

- *The deviation of the width from the SM prediction:*

$$DEV(h^0 \rightarrow X \bar{X}) = \Gamma(h^0 \rightarrow X \bar{X})_{MSSM} / \Gamma(h^0 \rightarrow X \bar{X})_{SM} - 1$$

$$X = c, b$$

# Scatter plot in $DEV(h^0 \rightarrow c \bar{c}) - DEV(h^0 \rightarrow b \bar{b})$ plane



-  $DEV(h^0 \rightarrow c \bar{c})$  and  $DEV(h^0 \rightarrow b \bar{b})$  can be very large simultaneously!:

$DEV(h^0 \rightarrow c \bar{c})$  can be as large as  $\sim \pm 60\%$ .


$DEV(h^0 \rightarrow b \bar{b})$  can be as large as  $\sim \pm 20\%$ .

- **ILC can observe such large deviations from SM at high significance (arXiv:1908.11299)!:**

$\Delta DEV(h^0 \rightarrow c \bar{c}) = (3.60\%, 2.40\%, 1.58\%)$  at (ILC250, ILC500, ILC1000)

$\Delta DEV(h^0 \rightarrow b \bar{b}) = (1.98\%, 1.16\%, 0.94\%)$  at (ILC250, ILC500, ILC1000)

# Scatter plot in $DEV(h^0 \rightarrow c \bar{c}) - DEV(h^0 \rightarrow b \bar{b})$ plane



## - Recent LHC data:

$$DEV(h^0 \rightarrow b \bar{b}) = 0.12 + 0.92 / -0.62 = [-0.50, 1.04] \text{ (ATLAS)} \text{ (arXiv:1909.02845)}$$

$$DEV(h^0 \rightarrow b \bar{b}) = 0.37 + 1.52 / -1.06 = [-0.69, 1.89] \text{ (CMS)} \text{ (arXiv:1809.10733)}$$

- **Both SM and MSSM are consistent with the recent ATLAS/CMS data!**  
**The errors of the recent ATLAS/CMS data are too large!**

## 5.2 $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$

$BR(h^0 \rightarrow b \bar{s} / s \bar{b}) \cong 0$  (SM)

$BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  can be as large as  $\sim 0.2\%$  (MSSM with QFV)!


(See also Gomez-Heinemeyer-Rehman, PR D93 (2016) 095021 [arXiv:1511.04342].)

ILC(250+500+1000) sensitivity could be  $\sim 0.1\%$  (at  $4 \sigma$  significance)!

Private communication with Junping Tian;

See also Barducci et al., JHEP 12 (2017) 105 [arXiv:1710.06657].

# Scatter plot in $T_{D23} - BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ plane



- There is a strong correlation between  $T_{D23} - BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ !
- $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  can be as large as **0.2%** for large  $T_{D23}$ !
- **ILC(250 + 500 + 1000) sensitivity could be  $\sim 0.1\%$  at 4 sigma significance!**

Private communication with Junping Tian;  
See also Barducci et al., JHEP 12 (2017) 105 [arXiv:1710.06657].

- LHC & HL-LHC sensitivity should not be so good due to huge QCD BG!

# 6. $h^0 \rightarrow \gamma\gamma, gg$ in the MSSM

- As the  $h^0$  decays to photon photon and gluon gluon are *loop-induced decays*, these decays are very *sensitive to New Physics!*

- We compute the widths  $\Gamma(h^0 \rightarrow \gamma\gamma)$  and  $\Gamma(h^0 \rightarrow gg)$  at NLO QCD level in the *MSSM with QFV*.

- Main 1-loop contributions to  $h^0 \rightarrow \gamma\gamma$ :

*[ $W^+$ / top-quark /  $su$ ] - loops [  $su = (\tilde{t} - \tilde{c}$  mixture)]*


*The  $su$ -loops can be enhanced by large trilinear couplings  $T_{U23}, T_{U32}, T_{U33}$ , resulting in sizable deviation of  $\Gamma(h^0 \rightarrow \gamma\gamma)$  from the SM width!*

- Main 1-loop contributions to  $h^0 \rightarrow gg$ :


*[top-quark /  $su$ ] - loops [  $su = (\tilde{t} - \tilde{c}$  mixture)]*

*The  $su$ -loops can be enhanced by large trilinear couplings  $T_{U23}, T_{U32}, T_{U33}$ , resulting in sizable deviation of  $\Gamma(h^0 \rightarrow gg)$  from the SM width!*

$$X = W^+ / t / \tilde{u}_{1,2}$$



$$X = t / \tilde{u}_{1,2}$$





- *We perform a MSSM parameter scan respecting all the relevant theoretical and experimental constraints.*

- *From the parameter scan, we find the followings:*

**(1)  $\text{DEV}(h^0 \rightarrow \gamma\gamma)$  and  $\text{DEV}(h^0 \rightarrow gg)$  can be sizable simultaneously:**


**$\text{DEV}(h^0 \rightarrow \gamma\gamma)$  can be as large as  $\sim \pm 1\%$ ,**

**$\text{DEV}(h^0 \rightarrow gg)$  can be as large as  $\sim \pm 4\%$ .**

**(2) There is a very strong correlation between  $\text{DEV}(h^0 \rightarrow \gamma\gamma)$  and  $\text{DEV}(h^0 \rightarrow gg)$ . This correlation is due to the fact that the stop-loop (stop-scharm mixture loop) contributions dominate the two DEVs.**

**(3) The deviation of the width ratio  $\Gamma(h^0 \rightarrow \gamma\gamma) / \Gamma(h^0 \rightarrow gg)$  in the MSSM from the SM value can be as large as  $\sim \pm 5\%$ .**

# Scatter plot in $\text{DEV}(h^0 \rightarrow \gamma\gamma) - \text{DEV}(h^0 \rightarrow gg)$ plane



- $\text{DEV}(h^0 \rightarrow \gamma\gamma)$  and  $\text{DEV}(h^0 \rightarrow gg)$  can be sizable simultaneously!
- There is a strong correlation between  $\text{DEV}(h^0 \rightarrow \gamma\gamma)$  and  $\text{DEV}(h^0 \rightarrow gg)$ !
- Future lepton colliders such as ILC can observe such sizable deviations from SM!  
(See arXiv:1908.11299 and Backup slides)

# 7. Conclusion

- *We have studied the decays*

*$h^0 (125\text{GeV}) \rightarrow c \bar{c}, b \bar{b}, b \bar{s}, \gamma\gamma, gg$  in the **MSSM with QFV**.*

- *Performing a systematic MSSM parameter scan respecting all of the relevant theoretical and experimental constraints, we have found the followings:*

*\*  $DEV(h^0 \rightarrow c \bar{c})$  and  $DEV(h^0 \rightarrow b \bar{b})$  can be very large simultaneously! :*

*$DEV(h^0 \rightarrow c \bar{c})$  can be as large as  $\sim \pm 60\%$ ,*

*$DEV(h^0 \rightarrow b \bar{b})$  can be as large as  $\sim \pm 20\%$ .*

*\* The deviation of the width ratio  $\Gamma(h^0 \rightarrow b \bar{b}) / \Gamma(h^0 \rightarrow c \bar{c})$  from the SM value can exceed  $\sim +100\%$ .*

*\*  $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  can be as large as  $\sim 0.2\%$ !*

*ILC(250 + 500 + 1000) sensitivity could be  $\sim 0.1\%$  at 4 sigma signal significance!*

\*  $DEV(h^0 \rightarrow \gamma\gamma)$  and  $DEV(h^0 \rightarrow gg)$  can be sizable simultaneously! :

$DEV(h^0 \rightarrow \gamma\gamma)$  can be as large as  $\sim \pm 1\%$ ,

$DEV(h^0 \rightarrow gg)$  can be as large as  $\sim \pm 4\%$ .

\* The deviation of the width ratio  $\Gamma(h^0 \rightarrow \gamma\gamma)/\Gamma(h^0 \rightarrow gg)$  from the SM value can be as large as  $\sim \pm 5\%$ .

\* There is a very strong correlation between  $DEV(h^0 \rightarrow \gamma\gamma)$  and  $DEV(h^0 \rightarrow gg)$ . This correlation is due to the fact that the stop-loop (stop-scharm mixture loop) contributions dominate the two DEVs.

- All of these large deviations in the  $h^0$  (125) decays are due to

large  $\tilde{c} - \tilde{t}$  mixing & large  $\tilde{c}/\tilde{t}$  involved trilinear couplings  $T_{U23}$ ,  $T_{U32}$ ,  $T_{U33}$  and large  $\tilde{s} - \tilde{b}$  mixing & large  $\tilde{s}/\tilde{b}$  involved trilinear couplings  $T_{D23}$ ,  $T_{D32}$ ,  $T_{D33}$ .

- Future lepton colliders such as ILC, CLIC, CEPC, FCC-ee can observe such large deviations from SM at high significance!

- In case the deviation pattern shown here is really observed at the future lepton colliders, then it would strongly suggest the discovery of QFV SUSY (MSSM with QFV)!

- See next slide also.

- *Our analysis suggests the following:*

*PETRA/TRISTAN  $e^- e^+$  collider discovered virtual  $Z^0$  effect for the first time.*

*Later, CERN  $p \bar{p}$  collider discovered the  $Z^0$  boson.*

*Similarly, lepton colliders, such as ILC, could discover virtual Sparticle effects for the first time in  $h^0(125)$  decays!*

*Later, FCC-hh  $p p$  collider could discover the Sparticles!*

*END*

*Thank you!*

# *Backup Slides*

## 2. MSSM with QFV

*The basic parameters of the MSSM with QFV:*

$$\{ \tan\beta, m_A, M_1, M_2, M_3, \mu, M^2_{Q,\alpha\beta}, M^2_{U,\alpha\beta}, M^2_{D,\alpha\beta}, T_{U\alpha\beta}, T_{D\alpha\beta} \}$$

(at  $Q = 1 \text{ TeV}$  scale)      ( $\alpha, \beta = 1, 2, 3 = u, c, t$  or  $d, s, b$ )

$\tan\beta$ :      ratio of VEV of the two Higgs doublets  $\langle H^0_2 \rangle / \langle H^0_1 \rangle$

$m_A$ :      CP odd Higgs boson mass (pole mass)

$M_1, M_2, M_3$ :       $U(1), SU(2), SU(3)$  gaugino masses

$\mu$ :      higgsino mass parameter

$M^2_{Q,\alpha\beta}$ :      left squark soft mass matrix

$M^2_{U\alpha\beta}$ :      right up-type squark soft mass matrix

$M^2_{D\alpha\beta}$ :      right down-type squark soft mass matrix

$T_{U\alpha\beta}$ :      trilinear coupling matrix of up-type squark and Higgs boson

$T_{D\alpha\beta}$ :      trilinear coupling matrix of down-type squark and Higgs boson



## 2. Key parameters of MSSM

*Key parameters in this study are:*

\* *QFV parameters:*  $M^2_{Q23}, M^2_{U23}, M^2_{D23}, T_{U23}, T_{U32}, T_{D23}, T_{D32}$

\* *QFC parameter:*  $T_{U33}, T_{D33}$

$M^2_{Q23} = (\tilde{c}_L - \tilde{t}_L \text{ mixing parameter})$

$M^2_{U23} = (\tilde{c}_R - \tilde{t}_R \text{ mixing parameter})$

$M^2_{D23} = (\tilde{s}_R - \tilde{b}_R \text{ mixing parameter})$

$T_{U23} = (\tilde{c}_R - \tilde{t}_L \text{ mixing parameter})$

$T_{U32} = (\tilde{c}_L - \tilde{t}_R \text{ mixing parameter})$

$T_{U33} = (\tilde{t}_L - \tilde{t}_R \text{ mixing parameter})$

$T_{D23} = (\tilde{s}_R - \tilde{b}_L \text{ mixing parameter})$

$T_{D32} = (\tilde{s}_L - \tilde{b}_R \text{ mixing parameter})$

$T_{D33} = (\tilde{b}_L - \tilde{b}_R \text{ mixing parameter})$



# Constraints on the MSSM parameters from K & B meson and $h^0$ data:

Table 5: Constraints on the MSSM parameters from the  $K$ - and  $B$ -meson data relevant mainly for the mixing between the second and the third generations of squarks and from the data on the  $h^0$  mass and couplings  $\kappa_b$ ,  $\kappa_g$ ,  $\kappa_\gamma$ . The fourth column shows constraints at 95% CL obtained by combining the experimental error quadratically with the theoretical uncertainty, except for  $B(K_L^0 \rightarrow \pi^0 \nu \bar{\nu})$ ,  $m_{h^0}$  and  $\kappa_{b,g,\gamma}$ .

Observable	Exp. data	Theor. uncertainty	Constr. (95%CL)
$10^3 \times  \epsilon_K $	$2.228 \pm 0.011$ (68% CL) [21]	$\pm 0.28$ (68% CL) [40]	$2.228 \pm 0.549$
$10^{15} \times \Delta M_K$ [GeV]	$3.484 \pm 0.006$ (68% CL) [21]	$\pm 1.2$ (68% CL) [40]	$3.484 \pm 2.352$
$10^9 \times B(K_L^0 \rightarrow \pi^0 \nu \bar{\nu})$	$< 3.0$ (90% CL) [21]	$\pm 0.002$ (68% CL) [21]	$< 3.0$ (90% CL)
$10^{10} \times B(K^+ \rightarrow \pi^+ \nu \bar{\nu})$	$1.7 \pm 1.1$ (68% CL) [21]	$\pm 0.04$ (68% CL) [21]	$1.7_{-1.70}^{+2.16}$
$\Delta M_{B_s}$ [ $\text{ps}^{-1}$ ]	$17.757 \pm 0.021$ (68% CL) [21, 41]	$\pm 2.7$ (68% CL) [42]	$17.757 \pm 5.29$
$10^4 \times B(b \rightarrow s \gamma)$	$3.32 \pm 0.15$ (68% CL) [21, 41]	$\pm 0.23$ (68% CL) [11]	$3.32 \pm 0.54$
$10^6 \times B(b \rightarrow s l^+ l^-)$ ( $l = e$ or $\mu$ )	$1.60_{-0.45}^{+0.48}$ (68% CL) [43]	$\pm 0.11$ (68% CL) [44]	$1.60_{-0.91}^{+0.97}$
$10^9 \times B(B_s \rightarrow \mu^+ \mu^-)$	$2.69_{-0.35}^{+0.37}$ (68% CL) [45]	$\pm 0.23$ (68% CL) [46]	$2.69_{-0.82}^{+0.85}$
$10^4 \times B(B^+ \rightarrow \tau^+ \nu)$	$1.06 \pm 0.19$ (68% CL) [41]	$\pm 0.29$ (68% CL) [47]	$1.06 \pm 0.69$
$m_{h^0}$ [GeV]	$125.09 \pm 0.24$ (68% CL) [48]	$\pm 3$ [49]	$125.09 \pm 3.48$
$\kappa_b$	$1.06_{-0.35}^{+0.37}$ (95% CL) [50]		$1.06_{-0.35}^{+0.37}$ (ATLAS)
	$1.17_{-0.61}^{+0.53}$ (95% CL) [51]		$1.17_{-0.61}^{+0.53}$ (CMS)
$\kappa_g$	$1.03_{-0.12}^{+0.14}$ (95% CL) [50]		$1.03_{-0.12}^{+0.14}$ (ATLAS)
	$1.18_{-0.27}^{+0.31}$ (95% CL) [51]		$1.18_{-0.27}^{+0.31}$ (CMS)
$\kappa_\gamma$	$1.00 \pm 0.12$ (95% CL) [50]		$1.00 \pm 0.12$ (ATLAS)
	$1.07_{-0.29}^{+0.27}$ (95% CL) [51]		$1.07_{-0.29}^{+0.27}$ (CMS)

# Main SUSY one-loop contributions to $h^0 \rightarrow c \bar{c}$




Figure 2: The main one-loop contributions with SUSY particles in  $h^0 \rightarrow c \bar{c}$ . The corresponding diagram to (e) with the self-energy contribution to the other charm quark is not shown explicitly.


## 5.2 Deviation of width ratio from the SM prediction

- *The deviation of the width ratio from the SM prediction:*

$$DEV(b/c) = [\Gamma(b) / \Gamma(c)]_{MSSM} / [\Gamma(b) / \Gamma(c)]_{SM} - 1$$

$$\Gamma(X) = \Gamma(h^0 \rightarrow X \bar{X})$$

## Scatter plot in $T_{U32} - DEV(b/c)$ plane



$T_{U32}$  (GeV)

$\tilde{c}_L - \tilde{t}_R$  mixing parameter

- There is a strong correlation between  $T_{U32} - DEV(b/c)$ !
- $DEV(b/c)$  can exceed  $\sim +100\%$  for large  $T_{U32}$  !

## 5.2 $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$

$BR(h^0 \rightarrow b \bar{s} / s \bar{b}) \cong 0$  (SM)

$BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  can be as large as  $\sim 0.2\%$  (MSSM with QFV)!


(See also Gomez-Heinemeyer-Rehman, PR D93 (2016) 095021 [arXiv:1511.04342].)

ILC(250+500+1000) sensitivity could be  $\sim 0.1\%$  (at  $4 \sigma$  significance)!

Private communication with Junping Tian;

See also Barducci et al., JHEP 12 (2017) 105 [arXiv:1710.06657]

# Scatter plot in $T_{D23} - BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ plane




- There is a strong correlation between  $T_{D23} - BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ !
- $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  can be as large as **0.2%** for large  $T_{D23}$ !
- **ILC(250 + 500 + 1000) sensitivity could be  $\sim 0.1\%$  at 4 sigma significance!**

Private communication with Junping Tian;  
 See also Barducci et al., JHEP 12 (2017) 105 [arXiv:1710.06657].

- LHC & HL-LHC sensitivity should not be so good due to huge QCD BG!




# Scatter plot in $T_{D32} - BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ plane




- There is also a strong correlation between  $T_{D32} - BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ !
- $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  can be as large as 0.2% for large  $T_{D32}$ !

# Scatter plot in $BR(h^0 \rightarrow b \bar{s} / s \bar{b}) - DEV(h^0 \rightarrow b \bar{b})$ plane




- There is a strong correlation between  $DEV(h^0 \rightarrow b \bar{b})$  &  $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ !
- This is due to the fact that  $DEV(h^0 \rightarrow b \bar{b})$  &  $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  have a common origin of enhancement effect, i.e. large trilinear couplings  $T_{D23,32,33}$  &  $T_{U23,32,33}$ .

## Scatter plot in $BR(h^0 \rightarrow b \bar{s} / s \bar{b}) - \tan\beta$ plane




- There is a strong correlation between  $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  &  $\tan\beta$ !
- $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$  can be as large as 0.2% for  $\tan\beta \sim 30$ !

*Caveat for very large  $DEV(h^0 \rightarrow c \bar{c})$*



# Caveat for very large $DEV(h^0 \rightarrow c \bar{c})$

***Gluino loop contribution to  $h^0 \rightarrow c \bar{c}$  can be very large (positive and negative) for large  $T_{U32} * M^2_{U23}$ !***




***The interference term between the tree diagram and the gluino one-loop diagram can be very large (positive and negative) for large  $T_{U32} * M^2_{U23}$ , which can lead to even **NEGATIVE** width  $\Gamma(h^0 \rightarrow c \bar{c})$  at one-loop level!***

***In this case perturbation theory breaks down!***




***A large deviation of  $\Gamma(h^0 \rightarrow c \bar{c})$  from the SM value is in principle possible due to large values of the product  $T_{U32} * M^2_{U23}$ .***

***Since there is no significant physical constraint on this product, the deviation  $DEV(h^0 \rightarrow c \bar{c})$  can be unnaturally large. So, we show only the results with a deviation from the SM up to  $\sim \pm 60\%$ .***

Contours of  $DEV(h^0 \rightarrow c \bar{c})$  in  $T_{U32} - M^2_{U23}$  plane



# Correlations among $DEV(h^0 \rightarrow b \bar{b})$ , $BR(h^0 \rightarrow b \bar{s} / s \bar{b})$ , $\tan\beta$



## *Effect of Resummation of the bottom Yukawa coupling at large $\tan\beta$*

*As for  $\Gamma(h^0 \rightarrow b \bar{b})$  &  $\Gamma(h^0 \rightarrow b \bar{s} / s \bar{b})$ , we have considered the large  $\tan\beta$  enhancement and the resummation of the bottom Yukawa coupling [1]. It turns out, however, that in our case with large  $m_A$  close to the decoupling Higgs limit, the **resummation effect ( $\Delta_b$  effect) is very small ( $< 0.1\%$ ) [2].***


[1] *M. Carena et al., Nucl. Phys. B 577 (2000) 88 [hep-ph/9912516].*

[2] *H. Eberl, E. Ginina, A. Bartl, K. Hidaka and W. Majerotto, JHEP 06 (2016) 143 [arXiv:1604.02366 [hep-ph]];*

*E. Ginina, A. Bartl, H. Eberl, K. Hidaka and W. Majerotto, PoS(EPS-HEP2015)146 [arXiv:1510.03714 [hep-ph]].*




## Scatter plot in $T_{U32} - DEV(\gamma/g)$ plane



- There is a strong correlation between  $T_{U32} - DEV(\gamma/g)$  !
- $DEV(\gamma/g)$  can be as large as  $\sim +4\%$  for large  $T_{U32}$  !

## Scatter plot in $T_{U33} - DEV(\gamma/g)$ plane



$T_{U33}$  (GeV)

$\tilde{t}_L - \tilde{t}_R$  mixing parameter

- There is a strong correlation between  $T_{U33} - DEV(\gamma/g)$  !
- $DEV(\gamma/g)$  can be as large as  $\sim +4\%$  for large  $T_{U33}$  !

# Higgs couplings at future colliders


## Higgs coupling precision in % at future colliders

arXiv:1910.11775, arXiv:1905.03764

CERN-LPCC-2018-04



- Future colliders under consideration will improve with respect to the HL-LHC the understanding of the Higgs boson couplings - 1-5%
  - **Coupling to charm** quark could be measured with an accuracy of  $\sim 1\%$  in future e+e- machines
  - **Couplings to  $\mu/\gamma/Z\gamma$**  benefit the most from the large dataset available at HL-LHC
  - At low energy top-Higgs coupling is not accessible at future lepton colliders



# Higgs couplings at future colliders



## Higgs coupling precision in % for ILC

arXiv:2203.07622  
 arXiv:1908.11299  
 arXiv:1506.07830



coupling	ILC250		ILC500		ILC1000	
	full	no BSM	full	no BSM	full	no BSM
$hZZ$	0.49	0.38	0.35	0.20	0.34	0.16
$hWW$	0.48	0.38	0.35	0.20	0.34	0.16
$hbb$	0.99	0.80	0.58	0.43	0.47	0.31
$h\tau\tau$	1.1	0.95	0.75	0.63	0.63	0.52
$hgg$	1.6	1.6	0.96	0.91	0.67	0.59
$hcc$	1.8	1.7	1.2	1.1	0.79	0.72
$h\gamma\gamma$	1.1	1.0	1.0	0.96	0.94	0.89
$h\gamma Z$	8.9	8.9	6.5	6.5	6.4	6.4
$h\mu\mu$	4.0	4.0	3.8	3.7	3.4	3.4
$htt$	—	—	6.3	6.3	1.0	1.0
$hhh$	—	—	20	20	10	10
$\Gamma_{tot}$	2.3	1.3	1.6	0.70	1.4	0.50
$\Gamma_{inv}$	0.36	—	0.32	—	0.32	—

Note  $C^3$  would run at 550 GeV, a factor 2 improvement to the top-Yukawa coupling (\*)



# *DEV error - coupling error relation*

$$\Delta DEV(h \rightarrow X X) = 2 \delta g(hXX)$$

$$\delta g(hXX) = [\text{Expected } \textit{relative} \text{ error of coupling } g(hXX)]$$

$$\Delta DEV(h \rightarrow X X) = [\text{Expected } \textit{absolute} \text{ error of deviation } \\ DEV(h \rightarrow X X)]$$