




# A brief word about Trigger (See Emilio's lectures for a complete description)

Paul Laycock



July 4th 2022




# Data's journey



Data Preparation



# The Atlas Trigger and DAQ



*Here be dragons... and muons*

Muon Spectrometer

Hadronic Calorimeter

Electromagnetic Calorimeter

Solenoid magnet

Tracking

Transition Radiation Tracker

Pixel/SCT detector



Proton

Neutron

Muon

Electron

Photon


Neutrino

The dashed tracks are invisible to the detector


**Physics object reconstruction =**  
**Tracks +**  
**Jets**  
*+ Muons*





# The Atlas Trigger and DAQ



# Calorimeter reconstruction



# Calo reconstruction



**Do simple reconstruction**  
**Simple because time is of the essence**

**Define thresholds**  
**At least X amount of energy is interesting**

**Watch out for how much data your interesting trigger takes (it may not be interesting enough !)**

# ATLAS Trigger Menu - 2018

| Trigger                         | Typical offline selection  | Trigger Selection            |                            | L1 Peak Rate [kHz]                                      | HLT Peak Rate [Hz] |
|---------------------------------|--|------------------------------|----------------------------|---|--------------------|
|                                 |  | L1 [GeV]                     | HLT [GeV]                  | L=2.0×10 <sup>34</sup> cm <sup>-2</sup> s <sup>-1</sup> |                    |
| Single leptons                  | Single isolated $\mu$ , $p_T > 27$ GeV                           | 20                           | 26 (i)                     | 16  | 218                |
|                                 | Single isolated tight $e$ , $p_T > 27$ GeV                       | 22 (i)                       | 26 (i)                     | 31  | 195                |
|                                 | Single $\mu$ , $p_T > 52$ GeV                                    | 20                           | 50                         | 16  | 70                 |
|                                 | Single $e$ , $p_T > 61$ GeV                                      | 22 (i)                       | 60                         | 28  | 20                 |
|                                 | Single $\tau$ , $p_T > 170$ GeV                                  | 100                          | 160                        | 1.4   | 42                 |
| Two leptons                     | Two $\mu$ , each $p_T > 15$ GeV                                  | 2 × 10                       | 2 × 14                     | 2.2   | 30                 |
|                                 | Two $\mu$ , $p_T > 23, 9$ GeV                                    | 20                           | 22, 8                      | 16  | 47                 |
|                                 | Two very loose $e$ , each $p_T > 18$ GeV                         | 2 × 15 (i)                   | 2 × 17                     | 2.0   | 13                 |
|                                 | One $e$ & one $\mu$ , $p_T > 8, 25$ GeV                          | 20 ( $\mu$ )                 | 7, 24                      | 16  | 6                  |
|                                 | One loose $e$ & one $\mu$ , $p_T > 18, 15$ GeV                   | 15, 10                       | 17, 14                     | 2.6   | 5                  |
|                                 | One $e$ & one $\mu$ , $p_T > 27, 9$ GeV                          | 22 (e, i)                    | 26, 8                      | 21  | 4                  |
|                                 | Two $\tau$ , $p_T > 40, 30$ GeV                                  | 20 (i), 12 (i) (+jets, topo) | 35, 25                     | 5.7   | 93                 |
|                                 | One $\tau$ & one isolated $\mu$ , $p_T > 30, 15$ GeV             | 12 (i), 10 (+jets)           | 25, 14 (i)                 | 2.4   | 17                 |
|                                 | One $\tau$ & one isolated $e$ , $p_T > 30, 18$ GeV               | 12 (i), 15 (i) (+jets)       | 25, 17 (i)                 | 4.6   | 19                 |
| Three leptons                   | Three very loose $e$ , $p_T > 25, 13, 13$ GeV                    | 20, 2 × 10                   | 24, 2 × 12                 | 1.6   | 0.1                |
|                                 | Three $\mu$ , each $p_T > 7$ GeV                                 | 3 × 6                        | 3 × 6                      | 0.2   | 7                  |
|                                 | Three $\mu$ , $p_T > 21, 2 × 5$ GeV                              | 20                           | 20, 2 × 4                  | 16  | 9                  |
|                                 | Two $\mu$ & one loose $e$ , $p_T > 2 × 11, 13$ GeV               | 2 × 10 ( $\mu$ )             | 2 × 10, 12                 | 2.2   | 0.5                |
|                                 | Two loose $e$ & one $\mu$ , $p_T > 2 × 13, 11$ GeV               | 2 × 8, 10                    | 2 × 12, 10                 | 2.3   | 0.1                |
| Single photon                   | One loose $\gamma$ , $p_T > 145$ GeV                             | 24 (i)                       | 140                        | 24  | 47                 |
| Two photons                     | Two loose $\gamma$ , each $p_T > 55$ GeV                         | 2 × 20                       | 2 × 50                     | 3.0   | 7                  |
|                                 | Two $\gamma$ , $p_T > 40, 30$ GeV                                | 2 × 20                       | 35, 25                     | 3.0   | 21                 |
|                                 | Two isolated tight $\gamma$ , each $p_T > 25$ GeV                | 2 × 15 (i)                   | 2 × 20 (i)                 | 2.0   | 15                 |
| Single jet                      | Jet ( $R = 0.4$ ), $p_T > 435$ GeV                               | 100                          | 420                        | 3.7   | 35                 |
|                                 | Jet ( $R = 1.0$ ), $p_T > 480$ GeV                               | 111 (topo: $R = 1.0$ )       | 460                        | 2.6   | 42                 |
|                                 | Jet ( $R = 1.0$ ), $p_T > 450$ GeV, $m_{\text{jet}} > 45$ GeV    | 111 (topo: $R = 1.0$ )       | 420, $m_{\text{jet}} > 35$ | 2.6   | 36                 |
| $b$ -jets                       | One $b$ ( $\epsilon = 60\%$ ), $p_T > 285$ GeV                   | 100                          | 275                        | 3.6   | 15                 |
|                                 | Two $b$ ( $\epsilon = 60\%$ ), $p_T > 185, 70$ GeV               | 100                          | 175, 60                    | 3.6   | 11                 |
|                                 | One $b$ ( $\epsilon = 40\%$ ) & three jets, each $p_T > 85$ GeV  | 4 × 15                       | 4 × 75                     | 1.5   | 14                 |
|                                 | Two $b$ ( $\epsilon = 70\%$ ) & one jet, $p_T > 65, 65, 160$ GeV | 2 × 30, 85                   | 2 × 55, 150                | 1.3   | 17                 |
|                                 | Two $b$ ( $\epsilon = 60\%$ ) & two jets, each $p_T > 65$ GeV    | 4 × 15, $ \eta  < 2.5$       | 4 × 55                     | 3.2   | 15                 |
| Multijets                       | Four jets, each $p_T > 125$ GeV                                  | 3 × 50                       | 4 × 115                    | 0.5   | 16                 |
|                                 | Five jets, each $p_T > 95$ GeV                                   | 4 × 15                       | 5 × 85                     | 4.8   | 10                 |
|                                 | Six jets, each $p_T > 80$ GeV                                    | 4 × 15                       | 6 × 70                     | 4.8   | 4                  |
|                                 | Six jets, each $p_T > 60$ GeV, $ \eta  < 2.0$                    | 4 × 15                       | 6 × 55, $ \eta  < 2.4$     | 4.8   | 15                 |
| $E_T^{\text{miss}}$             | $E_T^{\text{miss}} > 200$ GeV                                    | 50                           | 110                        | 5.1   | 94                 |
| $B$ -physics                    | Two $\mu$ , $p_T > 11, 6$ GeV, $0.1 < m(\mu, \mu) < 14$ GeV      | 11, 6                        | 11, 6 (di- $\mu$ )         | 2.9   | 55                 |
|                                 | Two $\mu$ , $p_T > 6, 6$ GeV, $2.5 < m(\mu, \mu) < 4.0$ GeV      | 2 × 6 ( $J/\psi$ , topo)     | 2 × 6 ( $J/\psi$ )         | 1.4   | 55                 |
|                                 | Two $\mu$ , $p_T > 6, 6$ GeV, $4.7 < m(\mu, \mu) < 5.9$ GeV      | 2 × 6 ( $B$ , topo)          | 2 × 6 ( $B$ )              | 1.4   | 6                  |
|                                 | Two $\mu$ , $p_T > 6, 6$ GeV, $7 < m(\mu, \mu) < 12$ GeV         | 2 × 6 ( $Y$ , topo)          | 2 × 6 ( $Y$ )              | 1.2   | 12                 |
| Main Rate                       |  |                              |                            | 86  | 1750               |
| B-physics and Light States Rate |  |                              |                            |   | 200                |

Share rate between different physics final states

According to physics goals and priorities