

Search for light dark matter with cosmic rays and neutrinos

Yongsoo Jho (Yonsei U.)

Based on

Phys.Lett.B 811 (2020) 135863 ([arXiv:2006.13910 \[hep-ph\]](#)),
[arXiv:2101.11262 \[hep-ph\]](#), and work in preparation,

Collaboration with


Jong-Chul Park (Chungnam Natl. U.), Seong Chan Park (IPAP, Seoul and Yonsei U.),
and Po-Yan Tseng (IPAP, Seoul and Yonsei U. and Natl. Tsing Hua U.)

June 8, 2022

CERN-CKC 2022 Workshop

DM candidates


in a point of view of particle physics



In a wide range of Dark Matter mass 10^{-22} eV – $10^{-12} M_{\text{Sun}}$,
Various DM candidates has been suggested.

DM candidates

in a point of view of particle physics





In a wide range of Dark Matter mass 10^{-22} eV – $10^{-12} M_{\text{Sun}}$,
Various DM candidates has been suggested.

In the ranges of keV-GeV masses, DM can be actively upscattered by energetic **cosmic rays** and **neutrinos** in our universe.

The limits on light DM direct detection



<https://supercdms.slac.stanford.edu/dark-matter-limit-plotter>



Conventional searches on halo DM using nuclear/electron recoils usually have the cliffs around 10-100 MeV, due to tiny kinetic energies which are lower than E thresholds.

The limits on light DM direct detection

<https://supercdms.slac.stanford.edu/dark-matter-limit-plotter>





Conventional searches on halo DM using nuclear/electron recoils usually have the cliffs around 10-100 MeV, due to tiny kinetic energies which are lower than E thresholds.

One way to probe light DM ($< \text{MeV-GeV}$, depending on interaction strength) \implies is to find the boosted DM?

The limits on light DM direct detection

<https://supercdms.slac.stanford.edu/dark-matter-limit-plotter>




Conventional searches on halo DM using nuclear/electron recoils usually have the cliffs around 10-100 MeV, due to tiny kinetic energies which are lower than E thresholds.

One way to probe light DM ($< \text{MeV-GeV}$, depending on interaction strength) \implies is to find the boosted DM?


by energetic **cosmic rays** and **neutrinos**

How many electrons?: Observed spectrum of electron CR




A caveat: All of these observation is in the local region.

How many neutrinos?: Spectrum of neutrinos in our universe




[Raffelt, Tamborra, Vitagliano et al. 19']

How many neutrinos?: Spectrum of neutrinos in our universe




[Raffelt, Tamborra, Vitagliano et al. 19']

How many neutrinos?: Spectrum of neutrinos in our universe




[Raffelt, Tamborra, Vitagliano et al. 19']

How many neutrinos?: Spectrum of neutrinos in our universe



- The solar neutrino spectrum has maximum around 300 keV
- & Includes several bumps/peaks from each nuclear channels.
- Dominant contributions to fluxes are very well measured by experiments (e.g. Borexino, Gemma)

How many neutrinos?: Spectrum of neutrinos in our universe




- Thermally produced from neutronization in proto-neutron stars
- Usually has Boltzmann peaks around 5-10 MeV (depending on flavor)
- The amount of flux is determined by star-formation-rate (SFR) including high redshift.
- Extragalactic origin & almost isotropic

A direct detection of DSNB suggested by [Beacom. 10']



Boosting keV-GeV DM with

- Electron Cosmic rays (based on observed data)






Boosting keV-GeV DM with

- Electron Cosmic rays (based on observed data)
- DSNB (extragalactic origin)



Boosting keV-GeV DM with

- Electron Cosmic rays (based on observed data)
- DSNB (extragalactic origin)



- Stellar neutrinos (Galactic/Extragalactic origin, This work)




Boosting keV-GeV DM with

first suggested by Y. Ema et al. (18')

- Electron Cosmic rays (based on observed data)
Electron CR (2 MeV - 90 GeV for the observed data)




- DSNB (extragalactic origin)
Neutrino (O(1)-O(100) MeV) & normalization predicted by SFR

first suggested by A. Das et al. (21')



- Stellar neutrinos (Galactic/Extragalactic origin, Our work)
Neutrino (O(100) keV - 20 MeV) & precisely measured by solar neutrino detection exp.

eCR-Boosted Dark Matter (BDM) & Observation of BDM at DM Direct detection




YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
PLB 811 (2020) 135863 [2006.13910] [hep-ph]

Charged Cosmic ray (electron) BDM provides an interesting possibility in DM direct detection/neutrino observatories.

Caveat) Large e-DM interaction can be suffered from the constraints by cosmology and SN.

$$\mathcal{L} \supset -X_\mu (g_e J_e^\mu + g_\chi J_\chi^\mu) + \dots$$



eCR-Boosted Dark Matter (BDM) & Observation of BDM at DM Direct detection

- The flux of DM, boosted by CR electron

$$\frac{d\Phi_{\text{DM}}}{d\Omega}(K_{\text{DM}}, b, l) = \frac{J(b, l)}{m_{\text{DM}}} \int dK_e \frac{d\Phi_e}{d\Omega} \frac{d\sigma_{\text{DM}e \rightarrow \text{DM}e}}{dK_{\text{DM}}}$$

electron CR flux
e_{CR}⁻ + χ_{halo} → e⁻ + χ_{Boosted}

$$J(b, l) = \int_{l.o.s} d\ell \rho_{\text{DM}}$$





- Boosted DM-electron (in the detector) cross section with light mediator X

$$\frac{d\sigma_X(\text{DM}e \rightarrow \text{DM}e)}{dK_e} = \frac{(g_e g_X)^2}{4\pi} \frac{2m_e(m_{\text{DM}} + K_{\text{DM}})^2 - K_e((m_e + m_{\text{DM}})^2 + 2m_e K_{\text{DM}}) + m_e K_e^2}{(2m_{\text{DM}} K_{\text{DM}} + K_{\text{DM}}^2)(2m_e K_e + m_X^2)^2}$$

eCR-Boosted Dark Matter (BDM) & Observation of BDM at DM Direct detection

YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
 PLB 811 (2020) 135863 [2006.13910] [hep-ph]


- Favored parameters



eCR-Boosted Dark Matter (BDM) & Observation of BDM at DM Direct detection

- Constraints (mediator mass/coupling)

YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
PLB 811 (2020) 135863 [2006.13910] [hep-ph]



Then, How about neutrinos?

**Q1: Can Cosmic "Neutrinos"
boost light Dark Matter in the halo?**


**Q2: Cosmic-Neutrino-Boosted Dark Matter
can be probed at
various ground experiments/observatories?**

Then, How about neutrinos?

Q1: Can Cosmic "Neutrinos" boost light Dark Matter in the halo?


Q2: Cosmic-Neutrino-Boosted Dark Matter can be probed at various ground experiments/observatories?

Q1: Can Cosmic "Neutrinos" boost light Dark Matter in the halo?



Galactic Neutrino-Boosted Dark Matter

Dark Matter boosted by neutrinos emitted from the Sun



$$K_\nu(K_{\text{DM}}, \bar{\theta}) = \frac{K_{\text{DM}}^2 - p'^2}{2(K_{\text{DM}} - p' \cos \bar{\theta})},$$


$$p'(K_{\text{DM}}) = \sqrt{2m_{\text{DM}}K_{\text{DM}} + K_{\text{DM}}^2}.$$

$$\frac{d\Phi_{\text{DM}}}{dK_{\text{DM}}} = \int dV \frac{\rho_{\text{DM}}}{m_{\text{DM}}} \frac{1}{r^2} \int_{K_\nu^{\text{min}}}^{K_\nu^{\text{max}}} \frac{d\sigma_{\nu\text{-DM}}(K_{\text{DM}}, \bar{\theta})}{dK_{\text{DM}}} \cdot \left(\frac{r_\odot}{4r' \sin \bar{\theta}} \right) \cdot \underbrace{\left(\frac{d^2\Phi^{\text{Solar } \nu}}{d\Omega dK_\nu} \right) \cdot \left(\frac{d_\odot}{r'} \right)^2}_{\frac{\dot{N}_\nu}{4\pi r_\odot^2}}$$

(It is valid in the point source limit, $r_\odot \rightarrow 0$)

Galactic Neutrino-Boosted Dark Matter

Dark Matter boosted by neutrinos emitted **from the Sun**



$$K_{\nu}(K_{\text{DM}}, \bar{\theta}) = \frac{K_{\text{DM}}^2 - p'^2}{2(K_{\text{DM}} - p' \cos \bar{\theta})},$$


$$p'(K_{\text{DM}}) = \sqrt{2m_{\text{DM}}K_{\text{DM}} + K_{\text{DM}}^2}.$$

$$\frac{d\Phi_{\text{DM}}}{dK_{\text{DM}}} \approx \frac{\dot{N}_{\nu}}{8\pi} \frac{\rho_{\text{DM}}}{m_{\text{DM}}} \int dV \frac{1}{r^2} \times \left(\left. \frac{dK_{\nu}}{d\bar{\theta}} \right|_{\bar{\theta}=\bar{\theta}_0} \right) \times \left(\left. \frac{d\sigma_{\nu-\text{DM}}}{dK_{\text{DM}}} \right|_{\bar{\theta}=\bar{\theta}_0} \right) \times \frac{1}{r'^2 \sin \bar{\theta}_0}$$

For a single neutrino source (here, our Sun) contribution!
(The volume integration for DM coordinates)

Galactic Neutrino-Boosted Dark Matter

The expectation of **total** Galactic Star neutrino BDM



-Assumption in the evaluation-

1. Symmetric population of Stars&DM
2. All stars have same luminosity as the Sun

Individual star contribution

$$\begin{aligned} \frac{d\Phi_{\text{DM}}^{(1)}(\vec{y})}{dK_{\text{DM}}} &\simeq \frac{1}{8\pi^2} \left(\tilde{f}_1 \frac{d\dot{N}_{\nu}^{\text{Sun}}}{dK_{\nu}} \right) \int d^3\vec{z} \frac{\rho_{\text{DM}}(|\vec{z}|)}{m_{\text{DM}}} \frac{1}{|\vec{x} - \vec{z}|^2} \\ &\times \left(\frac{dK_{\nu}}{d\bar{\theta}} \Big|_{\bar{\theta}=\bar{\theta}_0} \right) \left(\frac{d\sigma_{\nu\text{DM}}}{dK_{\text{DM}}} \Big|_{\bar{\theta}=\bar{\theta}_0} \right) \\ &\times \frac{1}{\sin \bar{\theta}_0} \frac{1}{|\vec{z} - \vec{y}|^2} \times \exp \left(-\frac{|\vec{z} - \vec{y}|}{d_{\nu}} \right), \end{aligned}$$

Total Galaxy contribution


$$\frac{d\Phi_{\text{DM}}}{dK_{\text{DM}}} = \int d^3\vec{y} n_{\text{star}}(\vec{y}) \frac{d\Phi_{\text{DM}}^{(1)}(\vec{y})}{dK_{\text{DM}}}$$

In a realistic estimation,
Production of BDM is highly
anisotropic, and depends on
spectrum of injected neutrinos.

Galactic Neutrino-Boosted Dark Matter


The expectation of **total** Galactic Star neutrino BDM

At the Earth, one can expect $\sim 10^{-15} - 10^{-16} \cdot \frac{d\Phi_{Solar\nu}}{dE_\nu}$




Galactic Neutrino-Boosted Dark Matter

The expectation of **total** Galactic Star neutrino BDM




Galactic Neutrino-Boosted Dark Matter

The expectation of **total** Galactic Star neutrino BDM





Galactic Neutrino-Boosted Dark Matter



The expectation of **total** Galactic Star neutrino BDM



Expectation of Galactic neutrino BDM distribution



Galactic Disk distribution





Arrival direction distribution of BDM

Non-trivial distribution of BDM arrival direction could help to probe nuBDM scenario.

YSJ, J.-C. Park, S.-C. Park, P.-Y. Tseng
[\[2101.11262\]](#) [hep-ph]

Arrival direction distribution of Galactic Neutrino-BDM

- $K_{\text{DM}} \gg m_{\text{DM}}$: **Forward** scattering is dominant (small effective volume)
- $K_{\text{DM}} \ll m_{\text{DM}}$: **Large-angle** scatterings are allowed (large effective volume)



Extragalactic contribution to Neutrino-BDM

- Main contribution to EG-nuBDM


YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
(work in preparation)

Dominant contributions coming from the region
in which both **neutrino** and **DM** are populated.

"Extragalactic" contribution to Cosmic-Neutrino-BDM

YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
(work in preparation)


- Schematic pictures for main contribution to EG-nuBDM



"Extragalactic" contribution to Cosmic-Neutrino-BDM


YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
(work in preparation)

- Schematic pictures for main contribution to EG-nuBDM





Recipes to evaluate Extragalactic contribution

YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
(work in preparation)



The estimation of Halo/Stellar Mass ratio, Size of Halo/Disk are mostly based on observation data.







Extragalactic contribution of neutrino-BDM flux

$$\mathcal{L}_{\text{eff.}} \supset -g_e \bar{e} \gamma^\mu e X_\mu - g_\nu \bar{\nu} \gamma^\mu P_L \nu X_\mu - g_{\text{DM}} \bar{\chi} \gamma^\mu \chi X_\mu$$

$$g_e g_{\text{DM}} = g_\nu g_{\text{DM}} = 10^{-6}$$

YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
(work in preparation)




**Q1: Can Cosmic "Neutrinos"
boost light Dark Matter in the halo?**

**Q2: Cosmic-Neutrino-Boosted Dark Matter
can be probed at
various ground experiments/observatories?**

Sensitivity limits on DM/ neutrino experiments

YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,

(work in preparation)




Sensitivity limits on DM/ neutrino experiments


$$\mathcal{L}_{\text{eff.}} \supset -g_e \bar{e} \gamma^\mu e X_\mu - g_\nu \bar{\nu} \gamma^\mu P_L \nu X_\mu - g_{\text{DM}} \bar{\chi} \gamma^\mu \chi X_\mu$$

YJ, J.-C. Park, S.-C. Park, P.-Y. Tseng,
(work in preparation)


$$g_e = g_\nu = g_X$$



Sensitivity limits on DM/ neutrino experiments





Sensitivity limits on DM/ neutrino experiments




Conclusion

- A noble mechanism to boost light DM by neutrinos emitted from stars in our/distant galaxies is proposed.
- Future neutrino exp (JUNO) and Direct detection (XENON/LUX) & distribution of arrival direction will help to probe neutrino-BDM scenario in near future.
- Extragalactic contribution to neutrino-BDM also has interesting features and depends on DM/mediator masses.




Thank you for your attention

Boosted DM flux



Ly-alpha constraints on neutrino-DM interactions


[0812.0010] [hep-ph]



- $> \sim 10\%$ fraction of light DM component can be constrained by Ly-alpha.

Constraints on light mediator

R. Harnik et al. [JCAP 07 (2012) 026] [1202.6073] [hep-ph]



Constraints on light mediator

R. Harnik et al. [JCAP 07 (2012) 026] [1202.6073] [hep-ph]

