

Cosmic antiparticles

Is there room for dark matter?

Yoann Génolini


Rencontres de Blois – Mai 2022

Collaborators :


M. Boudaud, P.-I. Batista, E. F. Bueno,
F. Calore, S. Caroff, M. Cirelli, L. Derome,
J. Lavalle, A. Marcowith, D. Maurin,
V. Poireau, V. Poulin, S. Rosier, P. Salati,
P. D. Serpico, M. Vecchi and N. Weinrich




Introduction



Introduction




Galactic cosmic-ray sources



SNRs



Star clusters



Pulsar Wind
Nebulae



Colliding wind
binaries




Protostellar jets
microquasars

... and others !

Two reasons to focus on **cosmic-ray antiparticles**:


1 - Very low fluxes:

- ~ 1 antiproton/10⁴ protons
- ~ 1 positron/150 electrons




→ Sensitive to small couplings


2 - Believed to be of secondary origin:



→ Astrophysical bkg can be easily estimated





Dark matter indirect search



Introduction


Dark matter indirect search



Game changer: **high-quality data!**

Numerous precision experiments in a large energy range

Precision level % from GeV to TeV



Outline

- 1 - Few words on positrons
- 2 - More words on antiprotons
- 3 - Conclusion and prospects

Few words on positrons


Outline

1 - Few words on positrons

2 - More words on antiprotons


3 - Conclusion and prospects

Few words on positrons



Also measured by:
 -HEAT (1997)
 -PAMELA (2009)
 -FERMI (2010)


Astrophysical bckg from secondary production:



(Boudaud, ..., Y.G.+ 2015)


Few words on positrons

High-energy cosmic positrons: the need for a primary source




Also measured by:
-HEAT (1997)
-PAMELA (2009)
-FERMI (2010)

Astrophysical bckg from secondary production:




(Boudaud, ..., Y.G.+ 2015)

The WIMP explanation of the positron excess




(Boudaud, ..., Y.G.+ 2015)

The WIMP explanation of the positron excess



(Boudaud, ..., Y.G.+ 2015)

The WIMP explanation of the positron excess



(Boudaud, ..., Y.G.+ 2015)

One DM candidate

→ Very few channels giving a good fit

→ Huge boost factors 10^3 - 10^5

Hadronic channel

Ruled out by $p\bar{p}$ constraints
(See second part)

Leptonic channel

Tensions with CMB+DS constraints
(Lopez, A.+ 2015, Planck Col.XIII+2015)

Cannot come from DM clumpiness
(Lavalle, J.+ 2006, Brun, P.+ 2009)


→ Analysis extended to low-E
No good fit found

(Boudaud, ..., Y.G.+ 2016)

Origin of this excess?

Introduction

Galactic cosmic-ray sources



SNRs



Star clusters



Pulsar Wind
Nebulae



Colliding wind
binaries



Protostellar jets
microquasars


... and others !

→ Could also be SNRs
(see e.g. Mertsch, P.+ 2020)

A natural astrophysical candidate
→ Currently investigated with
multimessenger studies:
 γ -ray/radio signal and e^+ anisotropies

(see e.g. Manconi, S.+ 2019)

Dark matter constraints with low-energy positrons



→ Voyager-1 crossed the heliopause in 2012


→ Direct measurement of the IS $e^+ + e^-$ flux

→ Low-energy cosmic positrons provide a stringent constraint on leptophilique DM models


(Cummings, A. + 2016)

Few words on positrons

Dark matter constraints with low-energy positrons



(Cummings, A. + 2016)



(Boudaud, M.+ 2016, 2018)

→ Stringent constraints on S and P wave annihilation

Outline


1 - Few words on positrons

2 - More words on antiprotons


3 - Conclusion and prospects

Is the case of antiprotons more exciting?


→ Preliminary AMS02 antiproton data from 2015



(Giesen,..., Y.G.+ 2015)



(Evoli, C.+ 2015)



(Kappl, R.+ 2015)

- Secondary predictions very close to the data
- Small deviations may indicate typical WIMP DM

→ Some claimed excesses

(Cui, M-Y.,+2017, Cuoco, A.+2017, Cholis, I.+2017)

Uncertainties data + prediction from different origins...


... A refined treatment of uncertainties is needed!

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
- Comparison with data = discrepancy ~ few 10GV*



(Boudaud, Y.G.+ 2019)


Courtesy M. Boudaud

More words on antiprotons

Secondary


A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
- Comparison with data = discrepancy ~ few 10GV*
- Errors on the data



*Small total error / Different correlation lengths
Dominated by acceptance around the excess*

→ Covariance matrix estimated from detector info.



(Boudaud, Y.G.+ 2019)


Courtesy M. Boudaud

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
Comparison with data = discrepancy ~ few 10GV
- Errors on the data
*Small total error / Different correlation lengths
Dominated by acceptance around the excess*
- Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections



(Boudaud, Y.G.+ 2019)


R [GV]

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
Comparison with data = discrepancy ~ few 10GV
- Errors on the data
Small total error / Different correlation lengths
Dominated by acceptance around the excess
→ Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
New data from NA61/SHINE (p+p)
NA49: (p+C) & LHCb: (p+He)
(Aduszkiewicz+2017, Anticic+ 2010, Aaij+2018)
- Updated parameterisation and uncertainties
(Winkler, M. 2016, Korsmeier+ 2018)



(Boudaud, Y.G.+ 2019)


R [GV]

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
 - Comparison with data = discrepancy ~ few 10GV
- Errors on the data
 - Small total error / Different correlation lengths
 - Dominated by acceptance around the excess
- Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
 - Updated parameterisation and uncertainties
 - Transport



(Boudaud, Y.G.+ 2019)


R [GV]

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
 - Comparison with data = discrepancy ~ few 10GV
- Errors on the data
 - Small total error / Different correlation lengths
 - Dominated by acceptance around the excess
- Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
 - Updated parameterisation and uncertainties
 - Transport
 - New Li, Be, B/C data from AMS02
 - Updated transport models and uncertainties (Y.G.+ 2017, 2019, 2021 Derome+ 2019, Weinrich, Y.G.+ 2020)



(Boudaud, Y.G.+ 2019)


R [GV]

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
 - Comparison with data = discrepancy ~ few 10GV
- Errors on the data
 - Small total error / Different correlation lengths
 - Dominated by acceptance around the excess
- Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
 - Updated parameterisation and uncertainties
 - Transport
 - Updated transport models and uncertainties
 - Parents



(Boudaud, Y.G.+ 2019)


R [GV]

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
Comparison with data = discrepancy ~ few 10GV
- Errors on the data
*Small total error / Different correlation lengths
Dominated by acceptance around the excess*
→ Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
→ Updated parameterisation and uncertainties
 - Transport
→ Updated transport models and uncertainties
 - Parents*New H, He, C,N,O... data from AMS02*
(AMS02 Collab. 2017, 2019)
→ Updated fit and contribution of high-Z elements



(Boudaud, Y.G.+ 2019)


R [GV]

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
 - Comparison with data = discrepancy ~ few 10GV
- Errors on the data
 - Small total error / Different correlation lengths
 - Dominated by acceptance around the excess
- Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
 - Updated parameterisation and uncertainties
 - Transport
 - Updated transport models and uncertainties
 - Parents
 - Updated fit and contribution of high-Z elements



(Boudaud, Y.G.+ 2019)


R [GV]

More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
Comparison with data = discrepancy ~ few 10GV
- Errors on the data
*Small total error / Different correlation lengths
Dominated by acceptance around the excess*
- Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
- Updated parameterisation and uncertainties
 - Transport
- Updated transport models and uncertainties
 - Parents
- Updated fit and contribution of high-Z elements
- Refined covariance matrix for the model



(Boudaud, Y.G.+ 2019)

R [GV]


More words on antiprotons

Secondary

A refined treatment of uncertainties

- Data: AMS02 antiproton from 2016
- Model: semi-analytical (USINE) (Maurin 2020)
- Comparison with data = discrepancy ~ few 10GV*
- Errors on the data
 - Small total error / Different correlation lengths*
 - Dominated by acceptance around the excess*
- Covariance matrix estimated from detector info.
- Errors on the model
 - Pbar production cross-sections
 - Updated parameterisation and uncertainties
 - Transport
 - Updated transport models and uncertainties
 - Parents
 - Updated fit and contribution of high-Z elements
 - Refined covariance matrix for the model
- Chi2 test with:

$$\chi^2 = (\text{data} - \text{model})^T (\mathcal{C}^{\text{model}} + \mathcal{C}^{\text{data}})^{-1} (\text{data} - \text{model})$$



(Boudaud, Y.G. + 2019)

More words on antiprotons

Secondary

Statistical tests (Boudaud, Y.G.+ 2019)

→ Chi2 definition:

$$\chi^2 = (\text{data} - \text{model})^T (\mathcal{C}^{\text{model}} + \mathcal{C}^{\text{data}})^{-1} (\text{data} - \text{model})$$

→ Chi2-test:

$$\chi^2 / \text{dof} = 0.77$$

$$p_{\text{value}} = 0.90$$


→ KS-test:

$$p_{\text{value}} = 0.27$$

→ AMS-02 antiprotons are consistent with a secondary astrophysical origin

Other studies confirmed (Heisig+ 2020)

Does that mean there cannot be statistical evidence for DM?




(Boudaud, Y.G.+ 2019)

R [GV]

More words on antiprotons

Secondary
+Dark Matter

Dark Matter antiproton component



Dark Matter antiproton component

→ Typical DM annihilation channels

$$b\bar{b}, W^+W^-, \mu^+\mu^-, q\bar{q}, hh$$

→ Inputs spectra from **PPPC4MID**

(Cirelli+ 202X)

→ DM profile considered

Generalized NFW profile (Navarro+ 1996)


$$\rho_{DM}(r) = \frac{\rho_s}{(r/r_s)^\gamma (1 + r/r_s)^{3-\gamma}}$$

| Profile | γ | r_s [kpc] | $\rho_s [M_\odot/\text{pc}^3]$ |
|----------------|----------|-------------|--------------------------------|
| benchmark NFW | 1.0 | 19.6 | 0.00854 |
| cored | 0.0 | 7.7 | 0.08931 |
| contracted NFW | 1.25 | 27.2 | 0.00361 |

(McMillan+ 2016 → but renormalized)

→ We use NFW as benchmark

→ Depends on the magnetic halo size H



Above GeV, at first order $\phi_{\bar{p}}^{DM} \propto L$

New AMS02 data on Be/B + e⁺ sensitive to L

→ Reevaluation of the halo size $L \approx 5 \pm 2$ kpc

(Weinrich,..., Y.G. + 2020)

More words on antiprotons

Secondary
+ Dark Matter

Calore, Cirelli, Derome, Genolini, Maurin, Salati, Serpico
SciPost Phys. 12, 163 (2022)

Exploring the nul hypothesis

→ No significant excess found

$$LR = -2 \ln \frac{\sup_{\lambda \in \Lambda} \mathcal{L}(\lambda)}{\sup_{\{\lambda, \mu\} \in \Lambda \cup M} \mathcal{L}(\lambda, \mu)}$$

Chernoff's theorem used, $\langle \sigma v \rangle = 0$
= pure secondary antiprotons

| Final state | Model | m^* [GeV] | $\langle \sigma v \rangle^*$ [cm ³ /s] | LR (denom) | LR (num) | LR | local signif. [σ] |
|--------------|--------|----------------|--|---------------|-------------|------|-------------------------------|
| $b\bar{b}$ | BIG | 109.3 | 1.71e-26 | 48.37 | 51.65 | 3.28 | 1.8 |
| $b\bar{b}$ | SLIM | 109.1 | 1.48e-26 | 48.77 | 51.70 | 2.93 | 1.7 |
| $b\bar{b}$ | QUAINT | 106.7 | 4.28e-27 | 45.32 | 45.53 | 0.22 | 0.5 |
| $q\bar{q}$ | BIG | 88.5 | 4.41e-27 | 50.31 | 51.65 | 1.35 | 1.2 |
| $\mu^+\mu^-$ | BIG | 155.7 | 2.65e-23 | 49.76 | 51.65 | 1.90 | 1.4 |
| W^+W^- | BIG | 106.8 | 2.20e-26 | 49.24 | 51.65 | 2.41 | 1.6 |
| hh | BIG | 166.7 | 3.62e-26 | 49.28 | 51.65 | 2.38 | 1.5 |


→ Major impact of uncertainty choice

| Err. data / model | local signif. [σ] | m^* [GeV] | $\langle \sigma v \rangle^*$ [cm ³ /s] |
|-------------------|-------------------------------|----------------|--|
| cov/cov | 1.81 | 109.3 | 1.71e-26 |
| cov/none | 2.39 | 10.5 | 5.07e-26 |
| diag/cov | 3.33 | 98.8 | 2.14e-26 |
| diag/none | 2.75 | 8.5 | 1.70e-25 |
| stat/cov | 5.19 | 89.7 | 1.48e-26 |
| stat/none | 4.49 | 8.0 | 2.98e-25 |


Some studies confirmed (Heisig+ 2020)

Some less cautious studies find excesses

(Cholis+ 2019, Cuoco+ 2017, Cui+2017)




Upper limits on the DM annihilation xs: **our results**




- Bounds for 5 representative annihilation channels
- NFW DM profile / BIG propagation model
- Weakening of the bound = *slight excess*
- $\mu^+ \mu^-$ bound not competitive

Upper limits on the DM annihilation xs: **our results**




- Bounds for 5 representative annihilation channels
- NFW DM profile / BIG propagation model
- Weakening of the bound = *slight excess*
- $\mu^+ \mu^-$ bound not competitive

Upper limits on the DM annihilation xs: comparison with other works



- New propagation models, calibrated on AMS02, fit better HE pbars
- Cui et al. 16: agree with high masses, at low masses difference in propagation model + significance of the excess
- Cuoco et al. 16: same qualitative differences

Upper limits on the DM annihilation xs: comparison with photon constraints



- Three different dSph gamma ray constraints:
 - conservative → aggressive
- Large Magellanic Cloud (LMC):
 - no excess in synchrotron radiation from $e^+ e^-$
 - band = uncertainties in B field and DM profile
- Complementarity of the pbar bound

Outline

- 1 - Few words on positrons
- 2 - More words on antiprotons
- 3 - Conclusion and prospects

Cosmic antiparticles: Is there room for dark matter?

- e^+ Persistent excess in tension with present DM bounds
Better explained by local pulsars
 - HE γ -ray/radio signal and e^+ anisotropies data are coming (HESS, LHAASO, CTA, Fermi, AMS100?...)
- \bar{p} No significant excess reported until now
Refined treatment of errors is essential
 - Finer analysis needs: statistic does not help!
 - experimental data covariance matrix from AMS02 collab.
 - better pbar production xs (LHCb, AMBER, ..)
 - AMS02 2021 data bring new challenges: needs new CR models?
 - Meanwhile, constraints competitive with the best bound of the literature
- \bar{d} Measurements eagerly awaited from GAPS!
 - First flight at the end of this year?
- *anti-He* Few events presumably detected by AMS02...
 - Let's wait a published version

Backup slides



Statistical analysis

→ Likelihood ratio definition

$$LR(\mu_0) = -2 \ln \frac{\sup_{\lambda \in \Lambda} \mathcal{L}(\lambda, \mu_0)}{\sup_{\{\lambda, \mu\} \in \Lambda \cup M} \mathcal{L}(\lambda, \mu)}$$

$(L, K, \delta, V_a, V_C, \sigma_{CR}, \dots)$ CR-space

$(\langle \sigma v \rangle, m_\chi, channel)$ DM-space

→ With the following factorisation

$$-2 \ln \mathcal{L}(\lambda, \mu) \equiv \chi_{\text{LiBeB}}^2(\lambda) + \chi_{\bar{p}}^2(\lambda, \mu)$$



Constraints on the CR space

Tightest constraints on the DM space


→ Simplification of the likelihood

$$-2 \ln \mathcal{L}(\lambda, \mu) \equiv -2 \ln \mathcal{L}(L, \mu) = \left\{ \frac{\log L - \log \hat{L}}{\sigma_{\log L}} \right\}^2 + x_i (\mathbf{C}^{-1})_{ij} x_j$$

Upper limits on the DM annihilation cross section



Upper limits on the DM annihilation cross section





→ New data from 2021

Consequences for CR antiparticles

New definition of MIN/MED/MAX → generalisation to positrons

Y.G. et al., arxiv 2103.04108



Benchmark models for indirect DM search

| SLIM | L [kpc] | δ | $\log_{10} K_0$ [kpc ² Myr ⁻¹] | R_1 [GV] | δ_1 |
|------|--------------|----------|--|---------------|------------|
| MAX | 8.40 | 0.490 | -1.18 | 4.74 | -0.776 |
| MED | 4.67 | 0.499 | -1.44 | 4.48 | -1.11 |
| MIN | 2.56 | 0.509 | -1.71 | 4.21 | -1.45 |


Prospects for **Dark-Matter** exploration with Cosmic-rays

UPPER LIMITS ON DIPOLE AMPLITUDE: ELECTRONS AND POSITRONS



Electrons

1.3 x 10⁶ events, 16 GeV < E < 350 GeV




Upper limit on dipole amplitude:

$$\delta_{e^-} (>16 \text{ GeV}) < 0.5\% \text{ at } 95\% \text{ C.I.}$$

[Phys. Rev. Lett. 122, 101101]

Positrons

9.9 x 10⁴ events, 16 GeV < E < 350 GeV




Upper limit on dipole amplitude:

$$\delta_{e^+} (>16 \text{ GeV}) < 1.9\% \text{ at } 95\% \text{ C.I.}$$

[Phys. Rev. Lett. 122, 041102]

Prospects for **Dark-Matter** exploration with Cosmic-rays



(Manconi, S.+ 2019)

Prospects for **Dark-Matter** exploration with Cosmic-rays

TABLE I. Respective p -values for different sources of errors. We take dof= 57, i.e. the number of \bar{p} data. Total errors on data are defined to be $\sigma_{\text{tot}} = \sqrt{\sigma_{\text{stat}}^2 + \sigma_{\text{syst}}^2}$.

| Error considered | χ^2/dof | p-value (χ^2) | p-value (KS) |
|--|---------------------|----------------------|--------------|
| σ_{stat} | 23 | 0 | 0 |
| σ_{tot} | 1.69 | 8.3×10^{-4} | 0 |
| $\mathcal{C}^{\text{data}}$ | 0.84 | 0.79 | 0.98 |
| σ_{stat} and $\mathcal{C}^{\text{model}}$ | 1.32 | 0.05 | 0.99 |
| σ_{tot} and $\mathcal{C}^{\text{model}}$ | 0.37 | 1.0 | 0.04 |
| $\mathcal{C}^{\text{data}}$ and $\mathcal{C}^{\text{model}}$ | 0.77 | 0.90 | 0.27 |