


# Probing Top-quark Operators with Precision Electroweak Measurements

Yiming Liu


Institute of High Energy Physics, Chinese Academy of Sciences

LHC TOP WG meeting 2022/6/15


Based on 2205.05655 with Yuhao Wang, Cen Zhang, Lei Zhang,  
Jiayin Gu



- 1, At LEP1/2, low-energy precision measurements, and ee collider in the future, we can use loops to open up more possibilities. Loop factor suppression will be compensated for by the precision.
- 2, Our study is one of the many first steps towards a more complete loop-level SMEFT global analysis.




# Theoretical framework



$$\frac{1}{p^2 - M^2} = -\frac{1}{M^2} \left[ 1 + \left( \frac{p^2}{M^2} \right) + \left( \frac{p^2}{M^2} \right)^2 + \dots \right]$$

$$\mathcal{L}_{\text{Eff}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{C_i^{(6)}}{\Lambda^2} \mathcal{O}_i^{(6)} + \sum_i \frac{C_i^{(8)}}{\Lambda^4} \mathcal{O}_i^{(8)} + \dots$$




# Theoretical framework

$X^2$		$\varphi^6$ and $\varphi^4 D^2$		$\psi^2 \varphi^3$	
$Q_G$	$f^{ABC} G_{\mu\nu}^A G_{\rho\sigma}^B G_{\tau\delta}^C$	$Q_{\varphi^6}$	$(\varphi^1 \varphi)^2$	$Q_{\varphi\psi^2}$	$(\varphi^1 \varphi)(\bar{l}_p e_r \varphi)$
$Q_{\tilde{G}}$	$f^{ABC} \tilde{G}_{\mu\nu}^A G_{\rho\sigma}^B G_{\tau\delta}^C$	$Q_{\varphi^4 D^2}$	$(\varphi^1 \varphi) \square (\varphi^1 \varphi)$	$Q_{\psi^2 \varphi^3}$	$(\varphi^1 \varphi)(\bar{q}_p u_r \tilde{\varphi})$
$Q_W$	$\varepsilon^{IJK} W_{\mu\nu}^I W_{\rho\sigma}^J W_{\tau\delta}^K$	$Q_{\varphi^2 D^2}$	$(\varphi^1 D^\mu \varphi)^* (\varphi^1 D_\mu \varphi)$	$Q_{\psi^2 \varphi^3}$	$(\varphi^1 \varphi)(\bar{q}_p d_r \varphi)$
$Q_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_{\mu\nu}^I W_{\rho\sigma}^J W_{\tau\delta}^K$				
$X^2 \varphi^2$		$\psi^2 X \varphi$		$\psi^2 \varphi^2 D$	
$Q_{\varphi G}$	$\varphi^1 \varphi G_{\mu\nu}^A G^{A\mu\nu}$	$Q_{\psi W}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\psi^2 D^2}$	$(\varphi^1 \bar{D}_\mu \varphi)(\bar{l}_p \gamma^\mu l_r)$
$Q_{\varphi \tilde{G}}$	$\varphi^1 \varphi \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	$Q_{\psi B}$	$(\bar{l}_p \sigma^{\mu\nu} e_r) \varphi B_{\mu\nu}$	$Q_{\psi^2 D^2}$	$(\varphi^1 \bar{D}_\mu \varphi)(\bar{l}_p \tau^I \gamma^\mu l_r)$
$Q_{\varphi W}$	$\varphi^1 \varphi W_{\mu\nu}^I W^{I\mu\nu}$	$Q_{\psi G}$	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{\psi^2 D^2}$	$(\varphi^1 \bar{D}_\mu \varphi)(\bar{e}_p \gamma^\mu e_r)$
$Q_{\varphi \tilde{W}}$	$\varphi^1 \varphi \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	$Q_{\psi W}$	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	$Q_{\psi^2 D^2}$	$(\varphi^1 \bar{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi B}$	$\varphi^1 \varphi B_{\mu\nu} B^{\mu\nu}$	$Q_{\psi B}$	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{\varphi} B_{\mu\nu}$	$Q_{\psi^2 D^2}$	$(\varphi^1 \bar{D}_\mu \varphi)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$Q_{\varphi \tilde{B}}$	$\varphi^1 \varphi \tilde{B}_{\mu\nu} B^{\mu\nu}$	$Q_{\psi G}$	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) \varphi G_{\mu\nu}^A$	$Q_{\psi^2 D^2}$	$(\varphi^1 \bar{D}_\mu \varphi)(\bar{q}_p \gamma^\mu q_r)$
$Q_{\varphi WB}$	$\varphi^1 \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$	$Q_{\psi W}$	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I \varphi W_{\mu\nu}^I$	$Q_{\psi^2 D^2}$	$(\varphi^1 \bar{D}_\mu \varphi)(\bar{d}_p \gamma^\mu d_r)$
$Q_{\varphi \tilde{W} B}$	$\varphi^1 \tau^I \varphi \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	$Q_{\psi B}$	$(\bar{q}_p \sigma^{\mu\nu} d_r) \varphi B_{\mu\nu}$	$Q_{\psi^2 D^2}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{q}_p \gamma^\mu d_r)$

$(\bar{L}L)(\bar{L}L)$		$(\bar{R}R)(\bar{R}R)$		$(\bar{L}L)(\bar{R}R)$	
$Q_{ll}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	$Q_{ee}$	$(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	$Q_{le}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qq}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	$Q_{uu}$	$(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	$Q_{lu}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$Q_{qq}^{(2)}$	$(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	$Q_{dd}$	$(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{ld}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$Q_{qq}^{(3)}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	$Q_{uu}$	$(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	$Q_{qe}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$Q_{qq}^{(4)}$	$(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	$Q_{dd}$	$(\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{qu}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
		$Q_{ud}^{(1)}$	$(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qu}^{(2)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{u}_s \gamma^\mu T^A u_t)$
		$Q_{ud}^{(2)}$	$(\bar{u}_p \gamma_\mu T^A u_r)(\bar{d}_s \gamma^\mu T^A d_t)$	$Q_{qu}^{(3)}$	$(\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t)$
$(\bar{L}R)(\bar{R}L)$ and $(\bar{L}R)(\bar{L}R)$		$B$ -violating			
$Q_{lckd}$	$(\bar{l}_p^c e_r)(\bar{d}_s q_t^c)$	$Q_{duu}$	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{ijk} [(d_{\mu\nu}^{\alpha\beta})^T C u_{\mu\nu}^{\gamma\delta}] [(q_2^{\alpha\beta})^T C l_1^{\gamma\delta}]$		
$Q_{qqqd}^{(1)}$	$(q_{\mu\nu}^a)_{jk} \varepsilon_{ik} (q_{\mu\nu}^b)_d$	$Q_{quu}$	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{ijk} [(q_{\mu\nu}^{\alpha\beta})^T C q_{\mu\nu}^{\gamma\delta}] [(u_1^{\alpha\beta})^T C e_1^{\gamma\delta}]$		
$Q_{qqqd}^{(2)}$	$(\bar{q}_p^c T^A u_r) \varepsilon_{jk} (\bar{q}_s^c T^A d_t)$	$Q_{quq}$	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{ijk} [(q_{\mu\nu}^{\alpha\beta})^T C q_{\mu\nu}^{\gamma\delta}] [(q_1^{\alpha\beta})^T C l_1^{\gamma\delta}]$		
$Q_{lqqd}^{(1)}$	$(\bar{l}_p^c e_r) \varepsilon_{jk} (q_{\mu\nu}^b)_d$	$Q_{duu}$	$\varepsilon^{\alpha\beta\gamma} [(d_{\mu\nu}^{\alpha\beta})^T C u_{\mu\nu}^{\gamma\delta}] [(u_2^{\alpha\beta})^T C e_1^{\gamma\delta}]$		
$Q_{lqqd}^{(2)}$	$(\bar{l}_p^c \sigma_{\mu\nu} e_r) \varepsilon_{jk} (q_{\mu\nu}^b)_{\sigma\mu}$				

[Grzadkowski, Iskrzynski, Misiak, Rosiek, 2010]



$$Q_{\varphi Q}^{(3)} = i \left( \phi^\dagger \tau^I D_\mu \phi \right) \left( \bar{Q} \gamma^\mu \tau^I Q \right)$$

$$Q_{\varphi Q}^{(1)} = i \left( \phi^\dagger D_\mu \phi \right) \left( \bar{Q} \gamma^\mu Q \right)$$

$$Q_{\varphi t} = i \left( \phi^\dagger D_\mu \phi \right) \left( \bar{t} \gamma^\mu t \right)$$

$$Q_{\varphi b} = i \left( \phi^\dagger D_\mu \phi \right) \left( \bar{b} \gamma^\mu b \right)$$

$$Q_{\varphi tb} = i \left( \tilde{\phi}^\dagger D_\mu \phi \right) \left( \bar{t} \gamma^\mu b \right)$$


$$Q_{tW} = \left( \bar{q} \sigma^{\mu\nu} \tau^I t \right) \tilde{\phi} W'_{\mu\nu}$$

$$Q_{bW} = \left( \bar{q} \sigma^{\mu\nu} \tau^I b \right) \phi W'_{\mu\nu}$$

$$Q_{tB} = \left( \bar{q} \sigma^{\mu\nu} t \right) \tilde{\phi} B_{\mu\nu}$$

$$Q_{bB} = \left( \bar{q} \sigma^{\mu\nu} b \right) \phi B_{\mu\nu}$$

[Cen Zhang, Nicolas Greiner, Scott Willenbrock, 2012]



# Theoretical framework

Impose a  $U(2)_u \otimes U(2)_d \otimes U(2)_q \otimes U(3)_l \otimes U(3)_e$  flavor symmetry


$\psi^2 \varphi^3$	$X^3$	$\varphi^4 D^2$
$Q_{u\varphi}^{ij} = (\varphi^\dagger \varphi)(\bar{q}_i u_j \tilde{\varphi})$ $Q_{d\varphi}^{ij} = (\varphi^\dagger \varphi)(\bar{q}_i d_j \varphi)$	$Q_W = \epsilon^{IJK} W_\mu^I W_\nu^J W_\rho^K$	$Q_{\varphi D} = (\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D_\mu \varphi)$
$\psi^2 \varphi^2 D$	$\psi^2 X \varphi$	$X^2 \varphi^2$
$Q_{\varphi l}^{ij(1)} = \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{l}_i \gamma^\mu l_j)$ $Q_{\varphi l}^{ij(3)} = \left( \varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{l}_i \tau^I \gamma^\mu l_j)$ $Q_{\varphi e}^{ij} = \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{e}_i \gamma^\mu e_j)$ $Q_{\varphi q}^{ij(1)} = \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{q}_i \gamma^\mu q_j)$ $Q_{\varphi q}^{ij(3)} = \left( \varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi \right) (\bar{q}_i \tau^I \gamma^\mu q_j)$ $Q_{\varphi u}^{ij} = \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{u}_i \gamma^\mu u_j)$ $Q_{\varphi d}^{ij} = \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{d}_i \gamma^\mu d_j)$ $Q_{\varphi ud}^{ij} = i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_i \gamma^\mu d_j)$	$Q_{uW}^{ij} = (\bar{q}_i \sigma^{\mu\nu} u_j) \tau^I \tilde{\varphi} W_{\mu\nu}^I$ $Q_{uB}^{ij} = (\bar{q}_i \sigma^{\mu\nu} u_j) \tilde{\varphi} B_{\mu\nu}$ $Q_{dW}^{ij} = (\bar{q}_i \sigma^{\mu\nu} d_j) \tau^I \varphi W_{\mu\nu}^I$ $Q_{dB}^{ij} = (\bar{q}_i \sigma^{\mu\nu} d_j) \varphi B_{\mu\nu}$	$Q_{\varphi WB} = \varphi^\dagger \tau^I \varphi W_{\mu\nu}^I B^{\mu\nu}$
$(\bar{L}L)(\bar{L}L)$	$(\bar{R}R)(\bar{R}R)$	$(\bar{L}L)(\bar{R}R)$
$Q_{ll}^{prst} = (\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$ $Q_{lq}^{prst(1)} = (\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$ $Q_{lq}^{prst(3)} = (\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	$Q_{ee}^{prst} = (\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$ $Q_{eu}^{prst} = (\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$ $Q_{ed}^{prst} = (\bar{e}_p \gamma_\mu e_r)(\bar{d}_s \gamma^\mu d_t)$	$Q_{le}^{prst} = (\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$ $Q_{lu}^{prst} = (\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$ $Q_{ld}^{prst} = (\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$ $Q_{qe}^{prst} = (\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$

$$Q_{\varphi Q}^{(+)} \equiv Q_{\varphi Q}^{(1)} + Q_{\varphi Q}^{(3)}$$

$$Q_{lQ}^{(+)} \equiv Q_{lQ}^{(1)} + Q_{lQ}^{(3)}$$

$$Q_{\varphi Q}^{(-)} \equiv Q_{\varphi Q}^{(1)} - Q_{\varphi Q}^{(3)}$$

$$Q_{lQ}^{(-)} \equiv Q_{lQ}^{(1)} - Q_{lQ}^{(3)}$$



# Theoretical framework

	Experiment	Observables
Low Energy	CHARM/CDHS/ CCFR/NuTeV/ APV/QWEAK/ PVDIS	Effective Couplings
Z-pole	LEP/SLC	$\frac{\text{Total decay width } \Gamma_Z}{\text{Hadronic cross-section } \sigma_{had}}$ $\frac{\text{Ratio of decay width } R_f}{\text{Forward-Backward Asymmetry } A_{FB}^f}$ $\frac{\text{Polarized Asymmetry } A_f}{\text{Total decay width } \Gamma_W}$
W-pole	LHC/Tevatron/ LEP/SLC	$\frac{\text{Branch Ratio of W Decay } Br(W \rightarrow l\nu_l)}{\text{Mass of W Boson } M_W}$ $\frac{\text{Hadronic cross-section } \sigma_{had}}{\text{Ratio of cross-section } R_f}$
$ee \rightarrow qq$	LEP/TRISTAN	$\frac{\text{cross-section } \sigma_f}{\text{Forward-Backward Asymmetry for } b/c \text{ } A_{FB}^f}$
$ee \rightarrow ll$	LEP	$\frac{\text{Forward-Backward Asymmetry } A_{FB}^f}{\text{Differential cross-section } \frac{d\sigma_f}{d\cos\theta}}$
$ee \rightarrow WW$	LEP	$\frac{\text{cross-section } \sigma_{WW}}{\text{Differential cross-section } \frac{d\sigma_{WW}}{d\cos\theta}}$

[J Erler and A Freitas. Electroweak model and constraints on new physics.] [D Geiregat, Gaston Wilquet, U Binder, H Burkard, U Dore, W Flegel, H Grote, T Mouthuy, H Øverås, J Panman, et al. First observation of neutrino trident production.]


[Aielet Efrati, Adam Falkowski, and Yotam Soreq. Electroweak constraints on flavorful effective theories.]

[Morad Aaboud, Georges Aad, Brad Abbott, Jalal Abdallah, O Abdinov, Baptiste Abeloos, Syed Haider Abidi, OS AbouZeid, Nadine L Abraham, Halina Abramowicz, et al. Measurement of the W-boson mass in pp collisions at  $\sqrt{s} = 7\text{GeV}$  with the ATLAS detector.]

[Electroweak Measurements in Electron-Positron Collisions at W-Boson-Pair Energies at LEP.]

[A Combination of Preliminary Electroweak Measurements and Constraints on the Standard Model.]

[Measurement of the cross-section and forward-backward charge asymmetry for the b and c-quark in  $e^+e^-$  annihilation with inclusive muons at  $\sqrt{s} = 58\text{GeV}$ ]



# Theoretical framework

1, The first class involves the third generation quarks

$$Q_{\varphi Q}^{(1)}, Q_{\varphi Q}^{(3)}, Q_{\varphi t}, Q_{\varphi b}, Q_{\varphi \varphi}, Q_{tW}, Q_{tB}, Q_{bW}, Q_{bB}$$

2, The second class have tree-level contribution to  $e^+ e^- \rightarrow \bar{f}f (f \neq t), e^+ e^- \rightarrow W^+ W^-$ .


$$Q_{\varphi Q}^{(1)}, Q_{\varphi Q}^{(3)}, Q_{\varphi u}, Q_{\varphi d}, Q_{\varphi l}^{(1)}, Q_{\varphi l}^{(3)}, Q_{\varphi e}, Q'_{ll}, Q_{\varphi D}, Q_{\varphi WB}, O_W$$

3, The third class are 4-fermion operators that directly contribute to the  $e^+ e^- \rightarrow \bar{f}f (f \neq t)$  and several low energy scattering processes at tree level.

$$Q_{qe}, Q_{eu}, Q_{ed}, Q_{lq}^{(1)}, Q_{lq}^{(3)}, Q_{lu}, Q_{ld}, O_{ll}, Q_{ee}, Q_{le}$$

4, The fourth class are 4-fermion operators that directly contribute to the  $e^+ e^- \rightarrow b\bar{b}$  at tree level.




$$Q_{lQ}^{(1)}, Q_{lQ}^{(3)}, Q_{lb}, Q_{eQ}, Q_{eb}$$







# Theoretical framework

1, For observable without  $Zbb$  couplings, the nine operators modify the self-energies of  $W$ ,  $Z$ ,  $\gamma$  at loop-level, and therefore affect all measurements indirectly.



## Example

$$\begin{aligned}
 Q_{\varphi Q}^{(-)} &= Q_{\varphi Q}^{(1)} - Q_{\varphi Q}^{(3)} \\
 &= - \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{q}_3 \tau^1 \gamma^\mu q_3) + \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{q}_3 \gamma^\mu q_3) \\
 &= - \left( \varphi^\dagger i \overleftrightarrow{D}_\mu^1 \varphi \right) (\bar{q}_3 \tau^1 \gamma^\mu q_3) - \left( \varphi^\dagger i \overleftrightarrow{D}_\mu^2 \varphi \right) (\bar{q}_3 \tau^2 \gamma^\mu q_3) \\
 &\quad - \left( \varphi^\dagger i \overleftrightarrow{D}_\mu^3 \varphi \right) (\bar{q}_3 \tau^3 \gamma^\mu q_3) + \left( \varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) (\bar{q}_3 \gamma^\mu q_3) \\
 &= \frac{-igv^2}{\sqrt{2}} W_\mu^+ (\bar{b} \gamma_\mu t) + \frac{igv^2}{\sqrt{2}} W_\mu^- (\bar{t} \gamma_\mu b) + \frac{igZ_\mu}{\cos \theta_W} v^2 \bar{t} \gamma_\mu t + \dots
 \end{aligned}$$



2, For observable with  $Zbb$  couplings,  $Q_{\varphi Q}^{(3)}$ ,  $Q_{\varphi Q}^{(1)}$ ,  $Q_{\varphi b}$ ,  $Q_{bW}$ ,  $Q_{bB}$  modify the  $Z \rightarrow b\bar{b}$  measurements at tree-level.



$$Q_{\varphi Q}^{(3)} = i(\phi^\dagger \tau^I D_\mu \phi) (\bar{Q} \gamma^\mu \tau^I Q)$$

$$Q_{\varphi Q}^{(1)} = i(\phi^\dagger D_\mu \phi) (\bar{Q} \gamma^\mu Q)$$

$$Q_{\varphi b} = i(\phi^\dagger D_\mu \phi) (\bar{b} \gamma^\mu b)$$

$$Q_{bW} = (\bar{q} \sigma^{\mu\nu} \tau^I b) \phi W'_{\mu\nu}$$

$$Q_{bB} = (\bar{q} \sigma^{\mu\nu} b) \phi B_{\mu\nu}$$



3, For observable with  $Zbb$  couplings,  $Q_{\varphi Q}^{(3)}$ ,  $Q_{\varphi Q}^{(1)}$ ,  $Q_{\varphi t}$ ,  $Q_{tW}$ ,  $Q_{tB}$  not only modify the self-energies of  $W$ ,  $Z$ ,  $\gamma$  at loop-level, but also modify the  $Zb\bar{b}$  vertex at loop-level.

$$Q_{\varphi Q}^{(3)} = i \left( \phi^\dagger \tau^I D_\mu \phi \right) \left( \bar{Q} \gamma^\mu \tau^I Q \right)$$



$$Q_{\varphi Q}^{(1)} = i \left( \phi^\dagger D_\mu \phi \right) \left( \bar{Q} \gamma^\mu Q \right)$$

$$Q_{\varphi t} = i \left( \phi^\dagger D_\mu \phi \right) \left( \bar{t} \gamma^\mu t \right)$$

$$Q_{\varphi tb} = i \left( \tilde{\phi}^\dagger D_\mu \phi \right) \left( \bar{t} \gamma^\mu b \right)$$

$$Q_{tW} = \left( \bar{q} \sigma^{\mu\nu} \tau^I t \right) \tilde{\phi} W'_{\mu\nu}$$

$$Q_{tB} = \left( \bar{q} \sigma^{\mu\nu} t \right) \tilde{\phi} B_{\mu\nu}$$



# Theoretical framework




Diagram 15 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 16 NP<sub>3</sub>, QED<sub>4</sub>, QED<sub>3</sub>




Diagram 17 NP<sub>2</sub>, QED<sub>4</sub>, QED<sub>3</sub>




Diagram 18 NP<sub>2</sub>, QED<sub>4</sub>, QED<sub>3</sub>

Diagrams made by MadGraph5\_aMC@NLO




Diagram 19 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 20 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 21 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 22 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 23 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 24 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>

Diagrams made by MadGraph5\_aMC@NLO




Diagram 25 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 26 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 27 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 28 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>




Diagram 29 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>







Diagram 30 NP<sub>2</sub>, QCD<sub>4</sub>, QED<sub>3</sub>

Diagrams made by MadGraph5\_aMC@NLO




# Theoretical framework



To better understand the impacts of the 3rd-generation-quark operators, we trade  $\frac{c_{\varphi D}}{\Lambda^2} Q_{\varphi D}$  and  $\frac{c_{\varphi WB}}{\Lambda^2} Q_{\varphi WB}$  in the Warsaw basis for  $\frac{c_{D\varphi B}}{\Lambda^2} iD^\mu \varphi^\dagger D^\nu \varphi B_{\mu\nu}$  and  $\frac{c_{D\varphi W}}{\Lambda^2} iD^\mu \varphi^\dagger \sigma_a \varphi W_{\mu\nu}^a$ :

$$\begin{aligned} Q_{D\varphi B} &\equiv iD_\mu \phi^\dagger D_\nu \phi B^{\mu\nu} \\ &= -\frac{g'}{4} Q_{\varphi B} + \frac{g'}{2} \sum_\psi Y_\psi Q_{\varphi\psi}^{(1)} + \frac{g'}{4} Q_{\varphi\Box} + g' Q_{\varphi D} - \frac{g}{4} Q_{\varphi WB} \end{aligned}$$

$$\begin{aligned} Q_{D\varphi W} &\equiv iD_\mu \phi^\dagger \sigma_a D_\nu \phi W^{a\mu\nu} \\ &= \frac{g}{4} \sum_F Q_{\varphi F}^{(3)} + \frac{g}{4} \left( 3Q_{\varphi\Box} + 8\lambda_\phi Q_\phi - 4\mu_\phi^2 (\phi^\dagger \phi)^2 \right) + \\ &+ \frac{g}{2} \left( y_{ij}^e (Q_{e\varphi})_{ij} + y_{ij}^d (Q_{d\varphi})_{ij} + y_{ij}^u (Q_{u\varphi})_{ij} + \text{h.c.} \right) \\ &- \frac{g'}{4} Q_{\varphi WB} - \frac{g}{4} Q_{\varphi W}, \end{aligned}$$



Operator	$C_{\varphi t}$	$C_{\varphi Q}^{(+)}$	$C_{\varphi Q}^{(-)}$	$C_{\varphi tb}$	$C_{tW}$	$C_{tB}$	$C_{t\varphi}$
$\mu_{EFT} = 125 \text{ GeV}$	2.5	1.3	3.2	9.3	0.2	0.07	0.9
$\mu_{EFT} = 1000 \text{ GeV}$	1.3	0.5	4.3	1.3	0.6	0.08	0.9
Current	2.3	5.1	1.2	5.3	0.06	0.145	3.9
Our results	0.286	0.04	0.336	14.8	0.822	0.592	–


[Ethier J J, Magni G, Maltoni F, et al. Combined SMEFT interpretation of Higgs, diboson, and top quark data from the LHC]

[Alioli S, Cirigliano V, Dekens W, et al. Right-handed charged currents in the era of the Large Hadron Collider [J/OL].]

[Maltoni F, Vryonidou E, Zhang C. Higgs production in association with a top-antitop pair in the Standard Model Effective Field Theory at NLO in QCD [J/OL].]



[Buckley A, Englert C, Ferrando J, et al. Constraining top quark effective theory in the LHC Run II era [J/OL].]

[Vryonidou E, Zhang C. Dimension-six electroweak top-loop effects in Higgs production and decay]







# Results




# Results





Marginalized bound at  $1\sigma$





Individual bound at  $1\sigma$





# Results(tree level)




# Results(tree level+loop level)



# Results



- 1 The outstanding precision of these measurements (especially at future lepton colliders) could be sensitive to many important loop contributions of the new physics.
- 2 The tree-level contributions of the bottom dipole operators to the electroweak processes are non-negligible.
- 3 Our study is one of the many first steps towards a more complete loop-level SMEFT global analysis, for which many improvements are still needed.



Thank you for your attention!

