Quark Matter 2023



Contribution ID: 165

Type: Poster

Measurement of Ξ_c^0 production as a function of multiplicity via hadron decay in pp collisions at \sqrt{s} = 13 TeV with ALICE

Tuesday 5 September 2023 17:30 (2h 10m)

Measurements of the production of heavy-flavour hadrons in proton–proton (pp) collisions provide an important test of quantum chromodynamics (QCD). The heavy-flavour production cross section can be calculated using the factorisation approach as a convolution of three factors: the parton distribution functions, the partonic cross section, calculated with perturbative QCD calculations, and the fragmentation functions. Recent measurements of the baryon-to-meson $(\Lambda_c^+/D^0, \Sigma_c^{0,++}/D^0, \Xi_c^{0,+}/D^0$ and BR $\times {}^0_c/D^0$) p_T -differential yield ratios in pp collisions are inconsistent with corresponding measurements in e^+e^- collision, indicating that the hadronisation of charm quarks is not a universal process among different collision systems. From measurements of the pT-differential ${}^+_c/D^0$ yield ratio as a function of the event multiplicity in pp collisions, the modification for the hadronisation mechanism is found to be multiplicity dependent. The measurement of the multiplicity dependence of the ${}^0_c/D^0$ yield ratio can provide further constraints on the study of charm hadronisation.

In this contribution, the production cross section of ${}^{0}_{c}$ baryon reconstructed from the hadronic decay channel ${}^{0}_{c} \rightarrow {}^{-}\pi^{+}$ and the baryon-to-meson yield ratio ${}^{0}_{c}/D^{0}$ as a function of charged-particle multiplicity in pp collisions at \sqrt{s} = 13 TeV at midrapidity with the ALICE detector at the LHC will be discussed.

Category

Experiment

Collaboration (if applicable)

Primary authors: FANG, Tao (Central China Normal University CCNU (CN)); JIANHUI, ZhuPresenter: FANG, Tao (Central China Normal University CCNU (CN))Session Classification: Poster Session

Track Classification: Heavy Flavor