



Contribution ID: 278

Type: Poster

## Measurements of jet $v_2$ in medium-sized systems at STAR

*Tuesday 5 September 2023 17:30 (2h 10m)*

Hard partonic scatterings, occurring at the early stages of heavy-ion collisions, produce jets, which experience the full evolution of the quark-gluon plasma (QGP). As they traverse through the QGP, jets lose energy through collisional and radiative processes, collectively known as the jet quenching. In semi-central heavy-ion events, the QGP takes an approximately elliptical shape in the transverse plane whose mean in-plane and out-of-plane distances differ. This fact can be used to vary the average path length for jets traversing the QGP, and those traveling in-plane should experience less quenching effects than those traveling out-of-plane. This differential quenching manifests as a suppression of jet yield out-of-plane relative to in-plane, quantified by jet  $v_2$ , the second order Fourier coefficient. In this poster, jet  $v_2$  will be presented from Ru+Ru, Zr+Zr, and O+O collisions at  $\sqrt{s_{NN}} = 200$  GeV with multiple jet resolution parameters. Studying jet  $v_2$  in collision systems of varying sizes may help disentangle path-length dependent quenching effects and other effects which could give rise to anisotropies in systems even smaller than O+O collisions.

### Category

Experiment

### Collaboration (if applicable)

STAR Collaboration

**Authors:** REED, Rosi (Lehigh University); PROTZMAN, Tristan (Lehigh University)

**Presenter:** PROTZMAN, Tristan (Lehigh University)

**Session Classification:** Poster Session

**Track Classification:** Jets