

Elliptic and triangular flow of light nuclei in Au+Au collisions in the BES-II energies using the STAR detector

Rishabh Sharma
(for the STAR Collaboration)
Indian Institute of Science Education and Research (IISER) Tirupati



September 5, 2023

Supported in part by:






Outline


- Motivation
- The STAR experiment
- Analysis details
- Results
 - Elliptic flow of light nuclei
 - Triangular flow of light nuclei
- Summary and Outlook



Motivation



STAR, PRL 130, 202301 (2023)




STAR, PRC 94, 034908 (2016)

- **Coalescence model**: Light nuclei are formed in the later stages of heavy-ion collision by the coalescence of protons and neutrons
- Recent measurements indicate **coalescence model** can reproduce light nuclei yields/ratios
- v_2/A of light nuclei was observed to be close to v_2 of protons for $p_T/A < 1.5$ GeV/c in BES-I data → Supporting **coalescence model**
- Higher statistics dataset in BES-II program will allow us to revisit and better understand the production mechanism of light nuclei



Motivation



ALICE, PRC 102, 055203 (2020)


$$v_{2(3),d}(p_T) = \frac{2v_{2(3),p}(p_T/2)}{1 + 2v_{2(3),p}^2(p_T/2)}$$

- At LHC energies:
 - v_2 was observed to deviate from mass number scaling by 20-30%
 - v_3 was observed to exhibit mass number scaling although with large uncertainties
- High statistics data in BES-II energies enables us to study mass number scaling of higher order flow harmonics at STAR energies



The STAR Experiment

- Particle identification is performed using
 - dE/dx information from **Time Projection Chamber (TPC)**
 - m^2 information from **Time of Flight (TOF)**
- BES-II upgrades:
 - **iTPC**: Large pseudorapidity coverage ($|\eta| < 1.5$)
 - **Endcap Time of Flight (eTOF)**: $-1.6 < \eta < -1.1$
 - **Event Plane Detector (EPD)**: $2.1 < |\eta| < 5.1$
 - Improved particle detection capabilities
 - Better track momentum and event plane resolution
- Datasets
 - **BES-II**: Au+Au collisions at $\sqrt{s_{NN}} = 14.6, 19.6, 27, \text{ and } 54.4 \text{ GeV}$



JINST 15 C07040 (2020)



Analysis Method

- The particle azimuthal distribution can be written as:

$$E \frac{d^3N}{dp^3} = \frac{1}{2\pi} \frac{d^2N}{p_T dp_T dy} \left(1 + \sum_{n=1}^{\infty} 2v_n \cos[n(\varphi - \Psi_n)] \right)$$

$$v_n = \langle \cos[n(\varphi - \Psi_n)] \rangle$$

v_2 : Elliptic flow

v_3 : Triangular flow



- n^{th} harmonic plane is calculated using the Q-vector:

$$Q_n \cos(n\Psi_n) = \sum_i w_i \cos(n\phi_i)$$


$$Q_n \sin(n\Psi_n) = \sum_i w_i \sin(n\phi_i)$$

$$\Psi_n = \left(\tan^{-1} \frac{\sum_i w_i \sin(n\phi_i)}{\sum_i w_i \cos(n\phi_i)} \right) / n$$


- η -sub event plane method is used



CMS, PRC 87 014902 (2013)




Elliptic flow



Triangular flow



Science 337, 310 (2012)



- ~10% improvement of resolution from BES-I owing to higher TPC acceptance and track momentum resolution



Light nuclei identification



	Proton	Deuteron	Triton	Helium-3
Identification using	m^2 -distribution	z-distribution	z-distribution	z-distribution
To increase purity of the signal	$ \text{n}\sigma < 3.0$	$2.8 < m^2 < 4.2$ (GeV/c 2) 2	$6.3 < m^2 < 9.5$ (GeV/c 2) 2	--

$$z_i = \ln \left(\frac{\langle dE/dx \rangle_{\text{measured}}}{\langle dE/dx \rangle_{i,\text{theory}}} \right)$$


$$m^2 = p^2 \left(\frac{1}{\beta^2} - 1 \right)$$



Elliptic flow of light nuclei



Elliptic flow (v_2)




- v_2 of light nuclei increases with increasing p_T for all collision energies in the measured p_T range
- v_2 shows mass ordering at low p_T between 1-2 GeV/c
- Statistical errors have reduced significantly compared to BES-I results

Proton v_2 : PRC 93, 014907 (2016); PRC 88, 014902 (2013); PLB 827, 137003 (2022)




Centrality dependence of v_2



- v_2 of d shows a strong centrality dependence
- Peripheral collisions have relatively larger v_2 due to their larger initial spatial anisotropy



Mass number scaling of v_2




➤ Systematic deviation of around 20-30% from mass number scaling is observed for all light nuclei in measured energies

Proton v_2 : PRC 93, 014907 (2016); PRC 88, 014902 (2013); PLB 827, 137003 (2022)



Comparison with AMPT+Coal.



➤ AMPT-SM model with a coalescence afterburner is in good agreement with $v_2(p_T)$ of d

PRC 72, 064901 (2005)

Nucl. Phys. A 729 (2003) 809–834


Proton v_2 : Phys. Rev. C 93, 014907 (2016); Phys. Rev. C 88, 014902 (2013); Phys. Lett. B 827, 137003 (2022)



Triangular flow of light nuclei




Triangular flow (v_3)



- v_3 of p and d increases with increasing p_T for all collision energies in the measured p_T range
- v_3 shows mass ordering at low p_T between 1-2 GeV/c




Mass number scaling of v_3



➤ $v_3(p_T)$ of d shows a good agreement with mass number scaling within $\sim 10\%$



Comparison with AMPT+Coal.



➤ AMPT-SM model with a coalescence afterburner is in good agreement with $v_3(p_T)$ of d

PRC 72, 064901 (2005)

Nucl. Phys. A 729 (2003) 809–834




Summary

- $v_2(p_T)$ of d , t , and ${}^3\text{He}$ is measured in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 14.6, 19.6, 27, \text{ and } 54.4 \text{ GeV (COL)}$
 - Clear centrality dependence is observed for d for all collision energies
 - 20-30% deviation of light nuclei v_2 from mass number scaling is observed
 - AMPT+Coal. seems to well describe the v_2 of d
- $v_3(p_T)$ of p and d is measured in Au+Au collisions at $\sqrt{s_{\text{NN}}} = 14.6, 19.6, 27, \text{ and } 54.4 \text{ GeV (COL)}$
 - v_3 of d shows a good agreement with mass number scaling
 - AMPT+Coal. seems to well describe the v_3 of d

Elliptic and Triangular flow measurements suggest coalescence to be the dominant mechanism of light nuclei production in heavy-ion collisions



- We will extend the analysis to the remaining BES-II energies (**Au+Au**, $\sqrt{s_{NN}} = 7.7, 9.2, 11.5, \text{ and } 17.3 \text{ GeV}$)
- Stay tuned for more exciting results on light nuclei from BES II energies



Other light (hyper-)nuclei flow results from STAR at QM 2023

Chengdong's talk (#666): First observation of $v_1(y)$ and $v_2(y)$ of light (hyper-)nuclei in FXT energies

Xiaoyu's talk (#632), Sharang's poster (#414): Event plane correlated directed ($v_1\{\Psi_1\}$) and triangular ($v_3\{\Psi_1\}$) flow of light nuclei in FXT energies



Thank you
