


Electromagnetic effects on charged particles

Andrzej Rybicki
H. Niewodniczański Institute of Nuclear Physics
Polish Academy of Sciences



by I. Sputowska

- 1) Prologue ;
- 2) Do never agree with your boss ;
- 3) EM effects on charged particles ;
- 4) Space-time evolution of the system ;
- 5) EM effects in small systems ;
- 6) UPC's ?
- 7) No epilogue.




Collision energy in the c.m.s. (center-of-mass system), per one pair of colliding nucleons.

Heavy-ion collisions:

Pb+Pb

$$\sqrt{s}_{NN} = 17.3 \text{ GeV}$$

spectator system



**produced particles
(π mesons)**

participant zone


1) Prologue


by I. Sputowska

- Charged spectators generate **electromagnetic fields**.
- These modify charged pion spectra in the **final state**.
- We use this effect as a new source of information on the **space-time evolution of the system**.


NA49 @ CERN SPS

Pb+Pb




Pb beam
E = 158 GeV/N

Particle identification



Please note:

advantages of fixed-target w.r.t. collider experiments:

- typically better coverage of kinematically available phase-space (p_x, p_y, p_z): “forward” hemisphere of the collision.

- Full coverage of low transverse momentum starting from $p_T=0$;


$$(p_T = \sqrt{p_x^2 + p_y^2})$$

- Easier to develop (add new subdetectors) ;


- Cheaper (?)


NA49 @ CERN SPS



Pb+Pb



Particle track reconstruction
+
Momentum measurement


Pb beam
E = 158 GeV/N






$$x_F = \frac{p_L}{p_L^{beam}} \quad (\text{c.m.s.})$$

2) Do never agree with your boss (*)



A historical (?) question:

- what is a heavy-ion collision?

A simple consequence of nucleon-nucleon processes? "New" physics? Both?



(*) unless present in this room



$$x_F = \frac{p_L}{p_L^{beam}} \quad (\text{c.m.s.})$$

2) Do never agree with y boss (*)


Now I will discuss the **ratios** of these distributions:

$$\pi^+/\pi^- = \frac{dn/dx_F(\pi^+)}{dn/dx_F(\pi^-)}$$

A historical (?) question

- what is a heavy-ion collision?

A simple consequence of nucleon-nucleon processes? "New" physics? Both?




(*) unless present in this room


π^+/π^- ratios in Pb+Pb @ $\sqrt{s}_{NN} = 17.3$ GeV

- simple superposition of proton and neutron collisions?

p+p, n+p collisions



predictions for nucleus-nucleus reactions



π^+/π^- ratios remember the structure (p/n content) of the nucleus...


$$X_F = \frac{p_L}{p_L^{beam}}$$

(c.m.s.)


π^+/π^- ratios in Pb+Pb @ $\sqrt{s}_{NN} = 17.3$ GeV

- simple superposition of proton and neutron collisions?

p+p, n+p collisions




predictions for nucleus-nucleus reactions



π^+/π^- ratios remember the structure (p/n content) of the nucleus...




... but what is happening in the *peripheral* Pb+Pb collision ?



Hypothesis no. 1: the “neutron halo” ?


A.Trzcińska et al., PRL87, 2001
R.Schmidt et al., PRC67, 2003
S.Mizutori et al., PRC61, 2000

P.Pawłowski, A.Szczurek, PRC70, 2004




- **Analysis of collision geometry:**
 $b = 10.9 \pm 0.5$ fm
- **Not possible to obtain 75% n, 25 % p**



Hypothesis no. 1: the "neutron halo" ?



A.Trzcińska et al., PRL87, 2001
R.Schmidt et al., PRC67, 2003
S.Mizutori et al., PRC61, 2000




P.Pawłowski, A.Szczurek, PRC70, 2004



- **Analysis of collision geometry:**
 $b = 10.9 \pm 0.5$ fm
- **Not possible to obtain 75% n, 25 % p**

3) EM effects




A. R.,
Acta Phys. Polon.
B42 (2011) 867

↑
spectator
velocity:

$$x_F = 0.15 = m_\pi / m_p$$


$$x_F = \frac{p_L}{p_L^{beam}} \quad (\text{c.m.s.})$$


P.Pawłowski, A.Szczurek, PRC70, 2004



$$x_F = \frac{p_L}{p_L^{beam}} = \frac{m_\pi \gamma \beta}{m_p \gamma \beta} = \frac{m_\pi}{m_p} = 0.15$$

$$\gamma = \frac{1}{\sqrt{1-\beta^2}}, \quad \beta = \frac{v}{c}$$





A. R.,
 Acta Phys. Polon.
 B42 (2011) 867



**spectator
 velocity:**

$$x_F = 0.15 = m_\pi / m_p$$




$$x_F = \frac{p_L}{p_L^{beam}} \quad (\text{c.m.s.})$$

$$x_F = \frac{p_L}{p_L^{beam}} = \frac{m_\pi \gamma \beta}{m_p \gamma \beta} = \frac{m_\pi}{m_p} = 0.15$$

$$\gamma = \frac{1}{\sqrt{1-\beta^2}}, \quad \beta = \frac{v}{c}$$



NA49 preliminary




A. R.,
Acta Phys. Polon.
B42 (2011) 867




↑
 spectator
 velocity:
 $x_F = 0.15 = m_\pi/m_N$

$$x_F = \frac{p_L}{p_L^{beam}} \quad (\text{c.m.s.})$$

$d_E \approx 0.75 \text{ fm} !$

A. R. and A. Szczurek.,
Phys. Rev. C75 (2007)
054903






A. R.,
Acta Phys. Polon.
B42 (2011) 867




↑
**spectator
velocity:
 $x_F = 0.15 = m_\pi/m_N$**

$$x_F = \frac{p_L}{p_L^{beam}} \quad (\text{c.m.s.})$$

$d_E \approx 0.75 \text{ fm} !$



A. R. and A. Szczurek.,
Phys. Rev. C75 (2007)
054903








A. R.,
Acta Phys. Polon.
B42 (2011) 867

↑
**spectator
velocity:
 $x_F = 0.15 = m_\pi / m_N$**



A. R. and A. Szczurek.,
Phys. Rev. C75 (2007)
054903








Directed flow:
 $V_1 = \langle \cos(\phi - \Psi_{RP}) \rangle$
reflects **sideways collective motion** of emitted particles.





EM effects induce charge splitting of directed flow.
(A.R., A.S., PRC 87, 2013)


$$y = \frac{1}{2} \ln \left(\frac{E + p_z}{E - p_z} \right)$$

$$y = \tanh^{-1}(v_z/c)$$











A. R.,
 Acta Phys. Polon.
 B42 (2011) 867

- EM effects provide **their own estimate** for the time of pion creation, at $y=0$.



d_E [fm]



4) Space-time evolution of forward pion production

Plan:


- Formulation of a simple model ;
- Validation with exp. data on rapidity distributions ;
- Application to EM effects.

PRC 95 (2017) 024908
PRC 99 (2019) 024908 (*)

(*) Yes, both papers have page no. 024908


PRC102 (2020) 014901

d_E [fm]




A. R., A. Szczurek et al., APP Supp. 9 (2016) 303

Bricks collide ...



PRC 95 (2017) 024908
 Idea by A. Szczurek,
 See also:
 R. Hagedorn, CERN-71-12
 W.D. Myers, NPA 296, 1978, 177

... and form "fire streaks"



Each fire streak fragments independently into pions

available energy


fire streak rapidity

$$\frac{dn}{dy} = A \cdot (E_s^* - m_s) \cdot \exp\left(-\frac{[(y - y_s)^2 + \epsilon^2]^{\frac{n}{2}}}{n\sigma_y^n}\right)$$

total fire streak energy


sum of brick masses

d_E [fm]




y/y_{beam}

Bricks collide ...



PRC 95 (2017) 024908
 Idea by A. Szczurek,
 See also:
 R. Hagedorn, CERN-71-12
 W.D. Myers, NPA 296, 1978, 177

... and form "fire streaks"






Each fire streak fragments independently into pions

$$\frac{dn}{dy} = A \cdot (E_s^* - m_s) \cdot \exp\left(-\frac{[(y - y_s)^2 + \epsilon^2]^{\frac{n}{2}}}{n\sigma_y^n}\right)$$




total fire streak energy

sum of brick masses

NA49, π^- , Pb+Pb, $\sqrt{s}_{NN} = 17.3$ GeV



data points from: NA49, PRC 86 (2012) 054903



Each fire streak fragments independently




$$f(y) = A \cdot (E_s^* - m_s) \cdot \exp\left(-\frac{[(y - y_s)^2 + \epsilon^2]^{\frac{n}{2}}}{n\sigma_y^n}\right)$$

total fire streak energy

sum of brick masses

- The pion rapidity distribution from **one fire streak** in Pb+Pb collisions is **similar** to the pion rapidity distribution in **p+p** reactions ;
- The difference in absolute normalization (**0.748**) can be directly obtained from the different energy repartition in **p+p** and **Pb+Pb** reactions (see PRC 99 (2019) 024908).

NA49, π^- , Pb+Pb, $\sqrt{s}_{NN} = 17.3$ GeV




Each fire streak fragments independently




$$f(y) = A \cdot (E_s^* - m_s) \cdot \exp\left(-\frac{[(y - y_s)^2 + \epsilon^2]^{\frac{n}{2}}}{n\sigma_y^n}\right)$$

total fire streak energy

sum of brick masses



- The pion rapidity distribution from **one fire streak** in Pb+Pb collisions is **similar** to the pion rapidity distribution in **p+p** reactions ;
- The difference in absolute normalization (**0.748**) can be directly obtained from the different energy repartition in **p+p** and **Pb+Pb** reactions (see PRC 99 (2019) 024908).




Each fire streak fragments independently

$$f(y) = A \cdot (E_s^* - m_s) \cdot \exp\left(-\frac{[(y - y_s)^2 + \epsilon^2]^{\frac{n}{2}}}{n\sigma_y^n}\right)$$

total fire streak energy


sum of brick masses



Courtesy by Harsch Shah, S.Hoche, arXiv: 1411.4085



Application to EM effects



- Longitudinal evolution of the system → from our model ;
- Initial (before the action of the EM field) rapidity distribution of pions from our model ;



- Initial p_T distribution of pions → from UrQMD v3.4 ;
- Fragmentation (expansion) of the spectator charge → included ;
- Isospin (p/n) effects between π^+ and π^- → included → PRC 99 (2019) 024908 ;
- Azimuthal anisotropies (flow), vorticity, transverse expansion → included optionally ;
- The pion creation time τ (taken in the fire streak c.m.s.) → taken as free parameter .



NA49, π^- , Pb+Pb, $\sqrt{s_{NN}} = 17.3$ GeV





First quantitative description of the electromagnetic distortion of π^+/π^- ratios in Pb+Pb collisions at 158 GeV/nucleon beam energy ($\sqrt{s_{NN}}=17.3$ GeV) [PRC102 (2020) 014901].

- ➔ Reasonable agreement with experimental data for $x_F \geq 0.1$;
- ➔ Inclusion of spectator expansion improves the description of exp. data ;
- ➔ **Short pion creation times ($0.5 < \tau < 2$ fm/c, to be compared with ~ 5.5 fm/c at $y=0$).**




Summary (1)

- EM effects induced by spectators bring **new information** on the **space-time evolution** of the process of (fast) pion production ;
- We obtained this information and used it ;
- Results look reasonable.



→ Small systems ;

→ UPC (gamma-gamma) .


d_E [fm]



SHINE = SPS Heavy Ion and Neutrino Experiment



$$\sqrt{s_{NN}} = 5.1-17.3(27.4) \text{ GeV}$$




Strong interactions

- study the onset of deconfinement
- search for the critical point





Cosmic rays and neutrinos

Adapted from Antoni Marcinek, QM22

5) Small systems









First ever measurement of these effects in small systems at the CERN SPS



M. Kielbaso for NA61/SHINE, App Supp. 12 (2019) 353
S. Bhosale for NA61/SHINE, Pos EPS-HEP2021 (2022) 309



$$\sqrt{s}_{NN} = 8.8 \text{ GeV}$$



From: Sneha Bhosale for the
NA61/SHINE collaboration,
MESON2021, 18 May 2021

Modelling EM effects in new Ar+Sc data

A. Rybicki, A. Szczurek, Phys. Rev. C 75, 054903 (2007)



- Non-expanding spectator system cannot describe data (contrary to Pb+Pb, see A. Rybicki et al., APPB 46,737 (2015))





need significant expansion velocity β of the charge cloud

- Optimal description: charge cloud moves slower than spectator system → presence of participant charge?

Modelling EM effects in new Ar+Sc data

A. Rybicki, A. Szczurek, Phys. Rev. C 75, 054903 (2007)




Entangled interplay of several phenomena: EM effects induced by the (small) spectator charge, those induced by the participant charge, isospin effects, spectator fragmentation, and others.

Further experimental guidance is needed.


Xe+La ?

Modelling EM effects in new Ar+Sc data

A. Rybicki, A. Szczurek, Phys. Rev. C 75, 054903 (2007)



NA61/SHINE





Entangled interplay of several phenomena: EM effects induced by the (small) spectator charge, these induced by the participant charge, isospin effects, spectator fragmentation, and others.

Further experimental guidance is needed.

Xe+La ?

6) Comment: how about gamma-gamma processes in ultra-peripheral collisions ?

- Think about **Pb+Pb** ($\gamma \gamma$) \rightarrow **e^+e^-** ;
- What happens to leptons once created ?
- Subject of long discussions (here in Kraków) ;
- Conceptual difficulties ;
- Never measured.




See talk by
Kasia Mazurek
(Tuesday).

2107.13239

6) No epilogue

The electromagnetic (EM) fields resulting from the presence of charged spectators induce **distortions** in spectra of charged particles, and result in **charge splitting of directed flow**.



These **spectator-induced EM effects** can be used to study the **space-time evolution of particle production**.

Such studies have shown that in high energy heavy-ion collisions, **faster** pions are produced **closer** to the spectator system. They also provided an **independent estimate** for the time of pion creation, at $y=0$.



A first quantitative description of the EM distortion of charge ratios (π^+/π^-) of fast pions produced in Pb+Pb collisions at $\sqrt{s_{NN}}=17.3$ GeV has been obtained. This gives an indication of significantly **shorter** pion production time scales (shorter proper times τ) w.r.t. what was obtained at central rapidity.

First ever measurements of these effects in **small systems** at the CERN SPS are now available from NA61/SHINE.



Thank you !

Extra slides


- p+A collisions with our model from Sec. 4.




- p+A collisions with our model from Sec. 4.



- More on directed flow.



STAR:
 L. Adamczyk et al., PRL 112 (2014) 162301
 (redrawn)
 ALICE:
 S. Acharya et al., PRL 125 (2020) 2, 022301,
 arXiv:1910.14406 [nucl-ex]



- v_1 (“directed flow”) is the sideways deflection of pions in the reaction plane :

$$v_1 \equiv \langle \cos(\varphi \text{ w.r.t. reaction plane}) \rangle$$
- the spectator charge induces *charge splitting* of v_1 .