

Neutrino Non-Standard Interactions: a solution to the the NOvA and T2K tension

Sabya Sachi Chatterjee




DAE-BRNS HEP Symposium 2022

IISER Mohali

Based on PRL. 126 (2021) 5, 051802 by S S Chatterjee & A Palazzo

13.12.2022

Current status of 3ν parameters (3σ bound) in the Standard framework



ArXiv: 2006.11237 by P. Salas et al., arXiv: 2007.14792 by Esteban et al., and arXiv: 2107.00532 by F. Capozzi et al.

NSI and its presence in the oscillation framework

The presence of the neutral current non-standard interactions (NSI) in neutrino oscillation can be realized through the Lagrangian as,



$$-\mathcal{L}_{NSI} = \frac{G_F}{\sqrt{2}} \sum_{\alpha, \beta, f} \varepsilon_{\alpha\beta}^f [\bar{\nu}_\alpha \gamma^\mu (1 - \gamma^5) \nu_\beta] [\bar{f} \gamma_\mu (1 \pm \gamma^5) f]$$

$$\alpha, \beta = e, \mu, \tau \text{ and } f = e, u, d$$

$$\varepsilon_{\alpha\beta} \equiv \sum_{f=e,u,d} \varepsilon_{\alpha\beta}^f \frac{N_f}{N_e} \quad N \text{ is the number density of fermions}$$


$$\varepsilon_{\alpha\beta} \simeq \varepsilon_{\alpha\beta}^e + 3\varepsilon_{\alpha\beta}^u + 3\varepsilon_{\alpha\beta}^d$$

→ Strength of NSIs



Now, the time evolution equation for the neutrino flavor eigenstates in presence of NSI is given by

$$i \frac{d}{dt} \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \\ |\nu_\tau\rangle \end{pmatrix} = \left[\frac{1}{2E} U \begin{pmatrix} m_1^2 & 0 & 0 \\ 0 & m_2^2 & 0 \\ 0 & 0 & m_3^2 \end{pmatrix} U^\dagger + V + V_{NSI} \right] \begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \\ |\nu_\tau\rangle \end{pmatrix}$$



 \mathbf{H}_{NSI}




Where,

$$V = \begin{pmatrix} V_{CC} + V_{NC} & 0 & 0 \\ 0 & +V_{NC} & 0 \\ 0 & 0 & +V_{NC} \end{pmatrix}, \quad V_{NSI} = V_{CC} \begin{pmatrix} \varepsilon_{ee} & \varepsilon_{e\mu} & \varepsilon_{e\tau} \\ \varepsilon_{e\mu}^* & \varepsilon_{\mu\mu} & \varepsilon_{\mu\tau} \\ \varepsilon_{e\tau}^* & \varepsilon_{\mu\tau}^* & \varepsilon_{\tau\tau} \end{pmatrix}$$

$$\varepsilon_{\alpha\beta} |_{\alpha \neq \beta} = |\varepsilon_{\alpha\beta}| e^{i\phi_{\alpha\beta}} \quad \text{and} \quad \varepsilon_{\alpha\beta} = (\varepsilon_{\beta\alpha})^*$$

The probability for one flavor ν_α transforming to another flavor ν_β is calculated as

$$P(\nu_\alpha \rightarrow \nu_\beta) = |S_{\beta\alpha}(L)|^2 = |(e^{-iHL})_{\beta\alpha}|^2$$



Limits from the IceCube (90% C.L.)

$$|\epsilon_{e\mu}^{\oplus}| \leq 0.15$$

$$|\epsilon_{e\tau}^{\oplus}| \leq 0.17$$

$$|\epsilon_{\mu\tau}^{\oplus}| \leq 0.023$$





$$\epsilon_{ee}^{\oplus} - \epsilon_{\mu\mu}^{\oplus} \rightarrow [-.25, -.15] \& [-.06, .04]$$

$$\epsilon_{\tau\tau}^{\oplus} - \epsilon_{\mu\mu}^{\oplus} \rightarrow [-.04, .045]$$



See the talk by T. Ehrhardt presented at PPNT, Uppsala (2019)

For more details please see PRD104(Oct, 2021) 072006


Results from the Collaborations



68% and 90% C.L. contours at 2 d.o.f





Combined analysis of T2K and NOvA



IO is preferred over NO in standard oscillation

However NO is preferred over IO in Presence of NSI

See also **PRL. 126 (2021) 5, 051801** by P. B. Denton, J. Gehrlein, & R. Pestes

T2K+NO ν AT2K+NO ν A



$$\Delta\chi^2 = \chi_{SM}^2 - \chi_{SM+NSI}^2$$

NMO	NSI	$ \varepsilon_{\alpha\beta} $	$\phi_{\alpha\beta}/\pi$	δ_{CP}/π	$\Delta\chi^2$
NO	$\varepsilon_{e\mu}$	0.15	1.38	1.48	4.50
	$\varepsilon_{e\tau}$	0.27	1.62	1.46	3.75
IO	$\varepsilon_{e\mu}$	0.02	0.96	1.50	0.07
	$\varepsilon_{e\tau}$	0.15	1.58	1.52	1.01

$$\chi_{SM,NO}^2 - \chi_{SM,IO}^2 = 1.87$$

$$\chi_{e\mu,NO}^2 - \chi_{e\mu,IO}^2 = -2.56$$

$$\chi_{e\tau,NO}^2 - \chi_{e\tau,IO}^2 = -0.21$$

T2K+NO ν AT2K+NO ν A





$$\chi_{e\mu,NO}^2 - \chi_{SM,IO}^2 = -2.63$$

$$\chi_{e\mu,IO}^2 - \chi_{SM,IO}^2 = -0.07$$

$$\chi_{e\tau,NO}^2 - \chi_{SM,IO}^2 = -1.21$$


$$\chi_{e\tau,IO}^2 - \chi_{SM,IO}^2 = -1.01$$

NSI with e-mu sector (NO) is better preferred over e-tau sector (NO) !




IceCube bounds
are compatible !


SM



SM + NSI ($\epsilon_{e\mu}$)




SM + NSI ($\epsilon_{e\tau}$)



Tension resolved !


SM




NSI, NO is preferred over NSI, IO

No discrepancy !

SM + NSI ($\epsilon_{e\mu}$)



SM + NSI ($\epsilon_{e\tau}$)



Conclusion

- We have investigated the impact of NSI on the current data of T2K and NOvA.
- ➔ More than 90% C.L. disagreement between T2K and NovA in the measurement of the Standard Model CP-phase. It can be resolved if one considers the presence of NSI of type $\epsilon_{e\mu}$ or $\epsilon_{e\tau}$
- Future data from T2K and NOvA, and future experiments like T2HK, DUNE and atmospheric current and future data is expected to confirm the presence of NSI and also will help resolving this ambiguity.
- ★ Our work also evidences the importance of JUNO like experiment to determine NMO unambiguously, irrespective of the presence of NSI.
- ✓ **The current T2K and NOvA data might be a hint of Physics Beyond the Standard Model !**

Thank you for your kind attention!

Introduction to NSI

The effect of coherent forward scattering must be taken into account when considering the oscillations of neutrinos traveling through matter. In particular, for the case of massless neutrinos for which vacuum oscillations cannot occur, oscillations can occur in matter if the neutral current has an off-diagonal piece connecting different neutrino types.

L. Wolfenstein
Phys. Rev. D 17, 2369

Strong constraints on NC-NSI from the non-observation of charged lepton flavor violation

Possible to avoid these bounds:

1. Model with neutral light mediators
2. Heavy mediators models arising in radiative neutrino mass model
3. Models with two mediators in the framework of dimension-8 operators

For references please see:

Y. Farzan [1505.06906](#), Y. Farzan, I. Shoemaker [1512.09147](#),
Y. Farzan, M. Tortola [1710.09360](#),
M. Gavela, D. Hernandez, T. Ota, and W. Winter [0809.3451](#),
K.Babu, P. B. Dev, S. Jana, and A. Thapa [1907.09498](#),
D. Forero and W. Huang [1608.04719](#)
And many more.

For overview see, [1907.00991](#)

In presence of NSI, the $\nu_\mu \rightarrow \nu_e$ survival probability can be written approximately as,

$$P_{\mu e} \simeq P_0 + P_1 + P_2 .$$

NSI (e- μ) sector

$$P_0 \simeq 4s_{13}^2 s_{23}^2 f^2$$

$$P_1 \simeq 8s_{13} s_{12} c_{12} s_{23} c_{23} \alpha f g \cos(\Delta + \delta)$$

$$P_2 \simeq 8s_{13} s_{23} v |\varepsilon_{e\mu}| [s_{23}^2 f^2 \cos(\delta + \phi_{e\mu}) + c_{23}^2 f g \cos(\Delta + \delta + \phi_{e\mu})]$$

NSI (e- τ) sector

$$P_0 \simeq 4s_{13}^2 s_{23}^2 f^2$$


$$P_1 \simeq 8s_{13} s_{12} c_{12} s_{23} c_{23} \alpha f g \cos(\Delta + \delta)$$

$$P_2 \simeq 8s_{13} s_{23} v |\varepsilon_{e\tau}| [s_{23} c_{23} f^2 \cos(\delta + \phi_{e\tau}) - s_{23} c_{23} f g \cos(\Delta + \delta + \phi_{e\tau})]$$

$$\Delta = \frac{\Delta m_{31}^2 L}{4E}, \quad f \equiv \frac{\sin[(1-v)\Delta]}{1-v}, \quad g \equiv \frac{\sin v\Delta}{v}, \quad |v| = \left| \frac{2V_{CC}E}{\Delta m_{31}^2} \right|$$

Brief description of the experimental setup T2K

T2K (Tokai to Kamioka)	
Baseline	295 KM
Detector mass	22.5 Kt
Proton Energy	30 GeV



515 kW stable operation achieved

$\nu : 1.97 \times 10^{21} \text{ POT}$

$\bar{\nu} : 1.63 \times 10^{21} \text{ POT}$

Brief description of the experimental setup NOvA

NOvA (Fermilab to Minnesota)


Baseline 810 KM

Detector mass 14 Kt

Proton Energy 120 GeV







Beam Power

Typically ~ 700 kW



$$\nu : 1.6 \times 10^{21} \text{ POT}$$

$$\bar{\nu} : 1.3 \times 10^{21} \text{ POT}$$







$\alpha = e, \mu, \tau$

For antineutrinos (inverse beta-decay)





$$\bar{\nu}_l + p \rightarrow l^+ + n$$

In Liquid Ar detector


$$\nu_l + Ar \rightarrow l^- + K$$



Bievent plots for NO








Bievent plots for IO



Mass hierarchy sensitivity might get highly impacted in presence of large NSI coupling in DUNE!

Phys.Rev.Lett. 124 (2020) 11, 111801 by F Capozzi, *S S Chatterjee*, & A Palazzo

T2K Dataset



NOvA Dataset

