



Contribution ID: 575

Type: Poster

π_0 branching ratios from Quantum Impedance Model

Friday 16 December 2022 14:00 (1 hour)

One of the important concepts that governs the amplitude and phase of energy transmission is impedance. The other is the concept of geometric wavefunction that arises from geometric algebra. While Pauli sigma matrices form the basis of space in 3D, the Dirac matrices are basis vectors of space-time in the geometric representation. Wavefunction interactions are modeled by geometric products, which turns fermions into bosons and vice-versa. Physical manifestation of vacuum wavefunction interactions follows from assignment of appropriate quantized E and B fields to the eight vacuum wavefunction components. This is utilized to calculate quantized impedance network as a function of energy, with its nodes specified in powers of alpha, the em coupling constant. The particle lifetimes have been multiplied by speed of light to obtain their coherence lengths, which are in turn converted to corresponding energy units and certain particle lifetimes such as that of π_0 and η are seen to be matching with the nodes of impedance network. Utilizing the fact that impedances must be matched for the energy transmission essential in a decay, we determine the branching ratios for π_0 .

Session

Beyond the Standard Model

Primary author: AWASTHI, Ayushi (Central University of Himachal Pradesh)

Co-authors: SASTRI, O.S.K.S. (Central University of Himachal Pradesh, Dharamshala); CAMERON, peter

Presenter: AWASTHI, Ayushi (Central University of Himachal Pradesh)

Session Classification: Poster - 4