

# NNLO QCD corrections to $W+2$ b-jet production

---

Rene Poncelet

in collaboration with Bayu Hartanto, Andrei Popescu, Simone Zoia  
based on: [2102.02516], [2205.01687] and [2209.03280]

LEVERHULME  
TRUST



UNIVERSITY OF  
CAMBRIDGE





European Research Council

Established by the European Commission

# W + b - jets

Motivation: → testing perturbative QCD: large NLO QCD corrections, 4FS vs. 5 FS  
→ modelling of flavoured jets




Experiment: [D0,1210.0627,0410062] [ATLAS,1109.1470,1302.2929][CMS,1312.6608,1608.07561]

Theory W+1 b-jet: [Campbell et al,0611348,0809.3003][Caola et.al.,1107.3714]

Theory W+2 b-jet: mb=0 [Ellis et al,9810489] onshell W: [Cordero et al,0606102] W(lv)bb: [Campbell et al,1011.6647]  
NLO+PS: [Oleari et al,1105.4488][Frederix et al,1110.5502] W(lv)bb: [Luisoni et al,1502.01213]  
W(lv)bb+≤3j: [Anger et al, 1712.05721]

# NLO QCD corrections for W+2b-jet

[Anger et al, 1712.05721]




- Large NLO QCD corrections + scale dependence
- Opening of  $qg$ -channel



- NNLO QCD corrections required!  
Main challenges:
  - **Two-loop amplitudes** [Bager'21,Hartanto'22]
  - **Subtraction** for high-multiplicity processes → Stripper [Czakon'10'14'19]

# Twoloop amplitudes

Generate diagrams (contributing to leading-colour) with QGRAF



Factorizing decay:  $A_6^{(L)} = A_5^{(L)\mu} D_\mu P$

$$M_6^{2(L)} = \sum_{\text{spin}} A_6^{(0)*} A_6^{(L)} = M^{(L)\mu\nu} D_{\mu\nu} |P|^2$$

Projection on scalar functions (FORM+Mathematica):  
 → anti-commuting  $\gamma_5$  + Larin prescription


$$M_5^{(L)} = \sum_{i=1}^{16} a_i^{(L)} v_i^{\mu\nu}$$



$$a_i^{(L)} = a_i^{(L),\text{even}} + \text{tr}_5 a_i^{(L),\text{odd}}$$

$$a_i^{(L),p} = \sum_j c_{j,i}(\{p\}, \epsilon) \mathcal{I}(\{p\}, \epsilon)$$

# Amplitude reduction




Automated framework using finite fields to reconstruct the rational coefficients of Pentagon functions based on FiniteFlow [Peraro'19]

Independent method w.r.t. [Abreu et. al.'21] → cross check (projection and  $\gamma_5$ )

# NNLO QCD subtraction scheme

Cross section requires combination with real-radiation contributions



Credit: Bayu Hartanto

Sector-improved residue subtraction scheme → C++ implementation STRIPPER

[Czakon'10][Czakon,Heymes'14][Czakon,Hameren,Mitov,Poncelet'19]

(established framework: top-quark pairs, 2- & 3-jet, VV, V+jet, H, 3-photon)

Oneloop-amps from OpenLoops2 [Bucionni,Lang,Lindert,Maierhoefer,Pozzorini,Zhang,Zoller(2018,2019)]

Born-amps from AvH lib [Bury,van Hameren(2015)]

# Phenomenology

NNLO QCD corrections to  $Wb\bar{b}$  production at the LHC  
Hartanto, Poncelet, Popescu, Zoia 2205.01687

- LHC @ 8 TeV in 5 FS, NNPDF31, scale:  $H_T = E_T(l\nu) + p_T(b1) + p_T(b2)$
- Phasespace definition to model **[CMS, 1608.07561]**:  
 $p_T(l) \geq 30 \text{ GeV}$   $|y(l)| < 2.1$   $p_T(j) \geq 25 \text{ GeV}$ ,  $|y(j)| < 2.4$
- Inclusive (at least 2 b-jets) and exclusive (exactly 2 b-jets, no other jets) jet phase spaces (defined by the flavour-kT jet algorithm **[Banfi'06]**)

- Inclusive:

~ +20% corrections

~ 7% scale dependence

- Exclusive:

~ + 6% corrections

~ 2.5% scale dependence (7-pt)

Compare decorrelated model: **[Steward'12]**

~ 11% scale dependence


	inclusive [fb]	$\mathcal{K}_{\text{inc}}$	exclusive [fb]	$\mathcal{K}_{\text{exc}}$
$\sigma_{\text{LO}}$	$213.2(1)^{+21.4\%}_{-16.1\%}$	-	$213.2(1)^{+21.4\%}_{-16.1\%}$	-
$\sigma_{\text{NLO}}$	$362.0(6)^{+13.7\%}_{-11.4\%}$	1.7	$249.8(4)^{+3.9(+27)\%}_{-6.0(-19)\%}$	1.17
$\sigma_{\text{NNLO}}$	$445(5)^{+6.7\%}_{-7.0\%}$	1.23	$267(3)^{+1.8(+11)\%}_{-2.5(-11)\%}$	1.067

$$\sigma_{Wb\bar{b},\text{excl.}} = \sigma_{Wb\bar{b},\text{incl.}} - \sigma_{Wb\bar{b}j,\text{incl.}}$$


$$\Delta\sigma_{Wb\bar{b},\text{excl.}} = \sqrt{(\Delta\sigma_{Wb\bar{b},\text{incl.}})^2 + (\Delta\sigma_{Wb\bar{b}j,\text{incl.}})^2}$$

# Differential cross sections

## Transverse momentum of lepton



## Invariant mass b-jet pair






# Fixed order flavoured jets beyond NLO

What is the problem with FO flavoured jets?

Example NNLO: double real radiation and subtraction



- If  $F(n+2)$  does not treat the flavour pair appropriately:  
→ double soft singularity not subtracted
- **Implies correlated treatment of kinematics and flavour information**

# Solution: Modified jet algorithms

Flavour kT algorithm:

**Infrared safe definition of jet flavor,**  
Banfi, Salam, Zanderighi hep-ph/0601139

Pair distance:

$$d_{ij} = R_{ij}^2 \begin{cases} \max(k_{T,i}, k_{T,j})^\alpha \min(k_{T,i}, k_{T,j})^{2-\alpha} & \text{softer of } i,j \text{ is flavoured} \\ \min(k_{T,i}, k_{T,j})^\alpha & \text{else} \end{cases}$$

Beam distance:

$$d_{i,B} = \begin{cases} \max(k_{T,i}, k_{T,B}(y_i))^\alpha \min(k_{T,i}, k_{T,B}(y_i))^{2-\alpha} & i \text{ is flavoured} \\ \min(k_{T,i}, k_{T,B}(y_i))^\alpha & \text{else} \end{cases}$$

$$d_B(\eta) = \sum_i k_{T,i}(\theta(\eta_i - \eta) + \theta(\eta - \eta_i))e^{\eta_i - \eta}$$



$$d_{\bar{B}}(\eta) = \sum_i k_{T,i}(\theta(\eta - \eta_i) + \theta(\eta_i - \eta))e^{\eta - \eta_i}$$

Problem: Measurements are done with anti-kT

A proper comparison would require to  
**unfold experimental data**

→ (flavour-) kT and anti-kT cluster partonic jets differently

→ Non-trivial procedure.



# Old problem, new approaches

Renewed interest:

- Anti-kT + flv.-kT flavour matching:

**QCD-aware partonic jet clustering for truth-jet flavour labelling** Buckley, Pollard 1507.00508

**Practical Jet Flavour Through NNLO**

Caletti, Larkoski, Marzani, Reichelt 2205.01109

**A dress of flavour to suit any jet**

Gauld, Huss, Stagnitto 2208.11138

- Fixed-order fragmentation:

**B-hadron production in NNLO QCD: application to LHC ttbar events with leptonic decays**, Czakon, Generet, Mitov and Poncelet, 2102.08267

**A Fragmentation Approach to Jet Flavor**

Caletti, Larkoski, Marzani, Reichelt 2205.01117

- Modified anti-kT algorithm:

**Infrared-safe flavoured anti-kT jets**, Czakon, Mitov, Poncelet 2205.11879

Proposed modification:

A soft term designed to modify the distance of flavoured pairs.


$$d_{ij}^{(F)} = d_{ij} \begin{cases} \mathcal{S}_{ij} & i,j \text{ is flavoured pair} \\ 1 & \text{else} \end{cases}$$

$$\mathcal{S}_{ij} = 1 - \theta(1 - x) \cos\left(\frac{\pi}{2}x\right) \quad \text{with} \quad x = \frac{k_{T,i}^2 + k_{T,j}^2}{2ak_{T,\max}^2}$$

# W+2 bjets: flavour anti-kT

Flavour anti-kT algorithm applied to Wbb production at the LHC


Hartanto, Poncelet, Popescu, Zoia 2209.03280



## Comparison to data

Measurement of the production cross section of a W boson in association with two b jets in pp collisions at  $\sqrt{s} = 8$  TeV, CMS 1608.07561

(assumes small unfolding corrections → wip)



Significant differences between kT and anti-kT  
In small DeltaR(bb) region. → Beam-function?!

# Summary & Outlook

---

## Summary

- Independent computation of two-loop five-point amplitudes with external mass
- NNLO QCD corrections to  $W+2b$ -jet production at the LHC
- Comparison of flavour sensitive jet-algorithms

## Outlook

- Application of Stripper to further 5-point signatures
- Working towards non-planar contributions (also for 1 ext. mass)  
→ See Abreu's summary at (HP)<sup>2</sup>
- Flavour-tagging  
→ more studies and comparisons between different algorithms needed
- Comparison with 4FS computation