

Contribution ID: 23 Type: not specified

A simple model to include initial-state and hot-medium effects in the computation of quarkonium nuclear modification factor

Monday 28 November 2022 17:00 (15 minutes)

Quarkonium suppression is one of the more useful observables to obtain information about the hot medium created in ultrarelativistic heavy-ion collisions.

In this talk, we discuss a simple way to implement both the initial-state effects and the hot-medium evolution, and to compute the quarkonium nuclear modification factor if the survival probability for a bound state at a given energy density is known.

Based on the Glauber model, the temperature of the evolving medium and the centrality dependence of the nuclear modification factor will be analysed.

Declaration

I certify that I have checked that I am authorised to submit the abstract with the listed co-authors with their current affiliations

Change of Speaker

I understand that change of speaker is allowed provided that no participant gives more than one talk. Otherwise, we will ask the speaker to choose between one or the other abstract to be presented.

Primary author: ESCOBEDO ESPINOSA, MIGUEL ANGEL (Instituto Galego de Física de Altas Enerxías (IG-

FAE))

Co-author: FERREIRO, Elena (Instituto Galego de Física de Altas Enerxías (IGFAE))

Presenter: ESCOBEDO ESPINOSA, MIGUEL ANGEL (Instituto Galego de Física de Altas Enerxías (IGFAE))

Session Classification: Parallel A - WG4&6

Track Classification: WG4: Heavy-quark and Quarkonium Physics