

# Soft Logs in Processes with Heavy Quarks



Andrea Ghira

QCD@LHC 2022, Orsay

Based on [arXiv:2207.13567](https://arxiv.org/abs/2207.13567) in collaboration with D. Gaggero,  
S.Marzani and G.Ridolfi

# Introduction

We consider heavy flavour production from the decay of a Higgs boson (colour singlet)



We want to compute the differential decay rate over  $x = \frac{2p_1 \cdot q}{q^2} = \frac{2E_b}{\sqrt{q^2}}$  which tends to 1 in the limit  $k \rightarrow 0$

# Massless vs Massive Scheme Approach

## Massless Scheme:

- Quark mass used as a regulator
- Cross section computed as a convolution of a coefficient function times a fragmentation function
- Logs of  $\xi = \frac{m^2}{q^2}$  resummed through DGLAP

## Massive Scheme:

- All mass dependence taken into account
- Kinematics treated correctly at every order
- Large logs spoil the convergence of the series.

# What is our goal?

Matching resummed scheme with fixed order calculations gives better prediction in the study of differential decay rate in various regions of  $\xi$ :

$$\tilde{\Gamma}(N, \xi) = \underbrace{\tilde{\Gamma}^{(\text{F.O.})}(N, \xi)}_{\xi = \mathcal{O}(1)} + \underbrace{\tilde{\Gamma}^{(\text{RES}, \xi)}(N, \xi)}_{\xi \ll 1} - \text{d.c}$$

- This is the FONLL scheme, where we match massive calculation with the resummed one
- We want to resum also soft logs in both schemes at NLL  $\longrightarrow$  **FO(NLL)<sup>2</sup>**

# Problems with Merging

We want to merge the two different calculations of the differential decay rate resumming logs of  $N$  in the large  $N$  limit.

## Massless Scheme

- Double logs of  $N$  with mass independent coefficient  
(arxiv:0107138, arxiv:2207.10038).

## Massive Scheme

- Single logs of  $N$  with mass dependent coefficient
- If we perform the limit  $\xi \rightarrow 0$  after the large  $N$  limit, we do not recover the massless case

Different logarithmic structure of the bremsstrahlung radiation in the two cases, in the soft limit

# Example at Fixed order

If we compute the process with one emission at fixed order in the small mass and soft limit we will find: (see also arxiv:0311101v1)

## Massless Scheme

Performing the massless limit first :

$$\lim_{x \rightarrow 1} \lim_{\xi \rightarrow 0} \frac{1}{\Gamma_0} \frac{d\Gamma}{dx} = -\frac{\alpha_s C_F}{\pi} \left[ \frac{\log \xi}{1-x} + \frac{\log(1-x)}{1-x} + \frac{7}{4} \frac{1}{1-x} + \dots \right]$$



## Massive Scheme

Performing the soft limit first

$$\lim_{\xi \rightarrow 0} \lim_{x \rightarrow 1} \frac{1}{\Gamma_0} \frac{d\Gamma}{dx} = -\frac{2\alpha_s C_F}{\pi} \left[ \frac{1 + \log \xi}{1-x} + \dots \right]$$



**The two limits do not commute, comparing the accuracy of the soft logs can be confusing**

# Another Problem with Merging

The soft resummation formula in the massive scheme is the product of a coefficient function times a soft function:

$$\tilde{\Gamma}(N, \xi) = C(\xi, \alpha_s) e^{-2 \int_0^1 dx \frac{x^{N-1} - 1}{1-x} \gamma_{\text{soft}}(\beta, \alpha_s((1-x)^2 q^2))}$$

where  $\gamma_{\text{soft}}(\beta, \alpha_s)$  is the soft anomalous dimension.

If we perform the massless limit of the first order coefficient  $C^{(1)}(\xi)$  we find:

$$C^{(1)}(\xi) = C_F \left( \frac{1}{2} \log^2 \xi + \log \xi + \frac{\pi^2}{2} + \mathcal{O}(\xi) \right)$$

This term is not predicted by DGLAP!

# Our Strategy

In the same spirit of FONLL, we would like to define a matching scheme:


$$\tilde{\Gamma}(N, \xi) = \underbrace{\tilde{\Gamma}^{(\text{F.O.})}(N, \xi)}_{\xi = \mathcal{O}(1) \text{ and } N = \mathcal{O}(1)} + \underbrace{\tilde{\Gamma}^{(\text{RES soft})}(N, \xi)}_{\xi = \mathcal{O}(1) \text{ and } N \gg 1} + \underbrace{\tilde{\Gamma}^{(\text{RES soft}, \xi)}(N, \xi)}_{\xi \ll 1 \text{ and } N \gg 1} - \text{d.c}$$

- The problem is that we cannot identify an all-order subtraction term.
- We modify the resummed massless scheme expression in order to behave like the massive expression if  $\frac{1}{N} \ll \xi \ll 1$



# The origin of the Problem

At NLL in the FF calculation one can see the resummation as a product of two independent jet functions:



In the measured leg the double logarithmic structure cancel between the quark jet function and the fragmentation function.

# Towards a Solution

Only the recoil jet function exhibit double soft logs, this is the function to modify:

- We can modify it function so that:
  1. When  $\xi \ll (1 - x) \ll 1$  we recover Catani-Cacciari formula (arxiv:0107138).
  2. When  $(1 - x) \ll \xi \ll 1$  we recover the massive case in the small mass limit.

**This is achieved by computing the recoil jet function in the quasi-collinear limit.**

# Calculation in the quasi-collinear limit

In the quasi-collinear limit we keep  $\xi \simeq \theta^2 \ll 1$ . Setting  $\rho = \frac{1}{N}$

$$J_{\bar{b}} = \int_0^1 dx \int_0^1 \frac{d\theta^2}{\theta^2 + \xi} P_{qq}^{(m)}(x) \frac{\alpha_s((1-x)^2 \theta^2 q^2)}{\pi} \left[ \Theta\left((1-x)(\theta^2 + \xi) < \rho\right) - 1 \right]$$

↓  
Massive splitting  
function
↓  
Massive contribution  
to the jet mass

At fixed coupling we obtain:

$$\rho > \xi : \quad J_{\bar{b}} = -\frac{1}{2} \log^2 \frac{1}{\rho} + \frac{3}{4} \log \frac{1}{\rho} + \mathcal{O}(\xi^0, \rho^0),$$

$$\rho < \xi : \quad J_{\bar{b}} = \frac{1}{2} \log^2 \frac{1}{\xi} - \log \frac{1}{\rho} \log \frac{1}{\xi} + \frac{3}{4} \log \frac{1}{\xi} + \log \frac{\xi}{\rho} + \mathcal{O}(\xi^0, \rho^0)$$

# Resummed formula

Our resummed formula looks like:

$$\begin{aligned} \log \tilde{\Gamma}(N, \xi) = & \log \left( 1 + \alpha_s \delta C^{(1)} \right) + J_{\bar{b}, \xi} \left( N, \frac{m^2}{q^2}, \frac{\mu_R^2}{q^2}, \alpha_s(\mu_R^2) \right) \\ & + J_b \left( N, \frac{\mu_F^2}{q^2}, \frac{\mu_R^2}{q^2}, \alpha_s(\mu_R^2) \right) + E \left( N, \frac{\mu_F^2}{\mu_{0F}^2}, \alpha_s(\mu_F^2) \right) \\ & + D_0 \left( N, \frac{\mu_{0F}^2}{m^2}, \frac{\mu_{0R}^2}{m^2}, \alpha_s(\mu_{0R}^2) \right) - 2 \int_{1/\bar{N}}^1 \frac{dz}{z} \frac{\alpha_s(z^2 m^2)}{\pi} \tilde{\gamma}_{\text{soft}}(\xi) \end{aligned}$$

- In the first line there is the modified jet function of the  $\bar{b}$  quark which accounts for the mass effects .
- In the following lines there is quark jet function plus the evolved fragmentation function.
- The last contribution is the  $\mathcal{O}(\xi)$  contribution to the soft dimension.

# Conclusions and Outlook

- The merging of the massive and massless calculation is far from trivial because of the fact that the massless and massive limit do not commute.
- We build a joint resummation in such a way that if we are in the regime in which  $\frac{1}{N} \ll \xi \ll 1$  we recover the massive scheme resummation and if  $\xi \ll \frac{1}{N} \ll 1$  we have the resummed expression by CC at NLL accuracy.
- What is the numerical impact of these finite mass corrections?
- Is it possible to extend this framework at NNLL accuracy?

Thanks for your attention